

FIRST APPROACH TO THE TROPHIC ECOLOGY AND DIET OF THE RAINBOW RUNNER, *Elagatis bipinnulata* (QUOY & GAIMARD, 1825) (PISCES: CARANGIDAE), IN THE CENTRAL COLOMBIAN CARIBBEAN

Primera aproximación a la ecología trófica del salmón, *Elagatis bipinnulata* (Quoy & Gaimard, 1825) (Pisces: Carangidae), en el Caribe central colombiano

CAMILO BERNARDO GARCÍA RAMIREZ¹, Ph. D; CAMILA POSADA², M.Sc.

¹ Profesor Departamento de Biología, Universidad Nacional de Colombia, sede Bogotá. cbgarcia@unal.edu.co

² Universidad de Bogotá Jorge Tadeo Lozano. camila.posada@utadeo.edu.co

Autor de correspondencia: Camilo García, cbgarcia@unal.edu.co

Received 23 December 2013, first decision 19 January 2014, accepted 19 February 2014.

Citation / Citar este artículo como: GARCÍA CB, POSADA C. First approach to the trophic ecology and diet of the rainbow runner, *Elagatis bipinnulata* (Quoy & Gaimard, 1825) (Pisces: Carangidae), in the Central Colombian Caribbean. Acta biol. Colomb. 2014. 19(2):309-314.

ABSTRACT

The rainbow runner (*Elagatis bipinnulata*) scores high in trophic level (4.24) but it is not a voracious fish (Q/B = 10.8). Its diet is dominated by the dwarf herring (*Jenkinsia lamprotaenia*), eats more in the dry season, than in the rainy season and significantly more in the afternoon than in the morning suggesting preference for daylight feeding.

Keywords: Colombian Central Caribbean, diet, rainbow runner, trophic ecology.

RESUMEN

El salmón (*Elagatis bipinnulata*) marca alto en nivel trófico (4,24) pero no es un pez voraz (Q/B = 10,8). Su dieta está dominada por la sardina enana (*Jenkinsia lamprotaenia*), come más en la época seca que en la época de lluvia y come significativamente más en la tarde que en la mañana, lo cual sugiere preferencia por la alimentación diurna.

Palabras clave: Caribe central colombiano, dieta, ecología trófica, salmón.

The rainbow runner (*Elagatis bipinnulata*) is a medium sized, oceanic and neritic, epipelagic fish of circumtropical distribution with intrusions into subtropical areas (e.g. Romanov, 2001). It is a highly migratory (Saito *et al.*, 1999) and swift swimming fish (Brunnschweiler and Sazima, 2010).

Qualitative references to the diet of the rainbow runner describe it as a piscivore, eating small pelagic fish (Hiatt and Strasberg, 1960; Yukinobo, 1998). However, the diet is complemented by a long list of zooplankton and pelagic invertebrates including copepods, gastropods, amphipods, Sagittidae, Euphaciacea, Thaliacea, *Siphonophora*, shrimps and crabs, stomatopods, polychaeta, cephalopods, fish eggs and even insects like *Halobates* sp. (Senta *et al.*, 1993).

The rainbow runner is in turn prey of bigger pelagic fishes, including blue marlin (*Makaira nigricans*, Shimose *et al.*, 2006), black marlin (*Istiompax indica*, Shimose *et al.*, 2008), sail fish (*Istiophorus platypterus*, Arizmendi-Rodríguez *et al.*, 2006), dolphinfish (*Coryphaena hippurus*, Bocanegra, 2007), silky shark (*Carcharhinus falciformis*, Bocanegra, 2007) and mammals like Fraser's dolphin (*Lagenodelphis hosei*, Watkins, 1994). Its juveniles are predated by sooty terns and brown noddies (Hensley and Hensley, 1995). Bocanegra (2007) reports instances of cannibalism.

There are few truly quantitative descriptions of the diet of the rainbow runner. One exception is the study by Vaske *et al.*, (2006) in the mid-tropical Atlantic (the Saint Peter and Saint Paul archipelago). They found a diet that included 27 taxa, dominated by flyingfishes (*Cypselurus cyanopterus* and *Exocoetus volitans*). Vaske *et al.*, (2006) suggested that *E. bipinnulata* eats both in daylight and in the night, but did not address seasonality. Dambacher *et al.* (2010), estimated a trophic level for *E. bipinnulata* of 3.59, which places it as an intermediary predator, and Opitz (1996) estimated a consumption to biomass ratio (Q/B) of 9.8 (recalculated from data in Opitz (1996), Appendix 8.7, Table 8.7.1 with the same empirical model as used here, see below. Originally 7.9) which is relatively low.

In Colombian Caribbean waters the rainbow runner, locally known as "salmón", is a common member of pelagic-neritic assemblages (Garzón and Acero, 1983; Grijalba-Bendeck *et al.*, 2004; Reyes-Nivia *et al.*, 2004), is also associated with artificial reefs (Delgadillo *et al.*, 2004; Delgadillo-Garzón and García, 2009), and is targeted in the fishery (Criales-Hernández *et al.*, 2006; Castro *et al.*, 2007). In the study area (see below), the rainbow runner is fished actively year round with a notable increase in landings from January to April, which corresponds to the dry season. It is not, however, the main pelagic target species either regional or locally (García, 2010).

This study presents the diet of the rainbow runner, reports its Q/B and trophic level and determines whether season (dry vs. rainy season) and the time of capture (morning vs. afternoon) influence diet.

Sampling was done in Taganga Bay and the National Natural Park Tayrona (NNPT), situated on the Central Colombian Caribbean Sea (11°21'00"-11°15'33"N, 73°54'06"-74°12'33"W). This coastal zone is influenced by the Caribbean Current, generated by the trade winds ("Alisios"), and the Panama-Colombia Countercurrent, and is characterised by seasonal upwelling during the dry season that extends from December to April (Ramírez, 1990; Andrade, 2001).

Fishes were captured by artisanal fishermen with beach seines and by trolling from March 2003 to May 2004 at approximately monthly intervals, thus covering the dry (December to April) and the rainy (May to November) seasons. Times of capture were before and after noon, but no nocturnal samples were taken. Individuals were weighed, measured

(fork length in cm) and their stomachs were immediately removed and injected with a buffered solution of ethyl alcohol (70 %), neutralised formaldehyde (10 %) and marine water in proportion 4:1:3. The stomachs were refrigerated until analysis. Prey items were identified to the lowest possible taxonomic level and weighed (wet weight). Unidentifiable stomach material was excluded in percentage weight calculations and general categories like "teleosts" were distributed proportionally among identified fish species or genera (see Abarca-Arenas *et al.*, 2007 for a similar approach).

The consumption/biomass (Q/B) ratio was estimated by means of an empirical model proposed by Palomares and Pauly (1998) that for carnivorous fishes is:

$$Q/B = 10^{[7.964 - 0.204 * \text{LOG } W_{\infty} - 1.965 * (1000 / (T_c + 273.15)) + 0.083 * A]}$$

where W_{∞} is the asymptotic weight of the individuals (wet weight, in g), T_c is the mean annual habitat temperature for the fish population in question (in °C) and A is the aspect ratio of the caudal fin (unitless). The asymptotic weight (5814 g) was estimated considering the maximum size of rainbow runner found in the study area, using the relation proposed by (Pauly, 1986):

$$W_{\infty} = W_{\text{max}} / 0.86$$

Sea surface temperature was recorded between September 2002 and March 2004 in Taganga Bay ($n = 57$, mean 26.8 °C, mercury thermometer, precision ± 0.1 °C) to obtain mean annual habitat temperature. The aspect ratio (A) of the caudal fin, a measurement of fish activity (Pauly, 1986), was estimated as:

$$A = (h^2/s)$$

where h is the distance between the upper and lower lobule and s is the surface area of the caudal fin ($n = 22$, mean $A = 4.7$). Fins were projected on millimeter paper to estimate s . Trophic level (defined as the weighted mean trophic level of prey items plus one) of the rainbow runner was estimated using the application TrophLab (<http://www.fishbase/download/TrophLab2K.zip>). See García and Contreras (2011) for a review of Colombian Caribbean fishes trophic levels.

A permutation procedure (Good, 2006) was used to test whether mean weight of stomach content (including unidentifiable material and stomachs with zero content) was different (one tailed test) between seasons (dry vs. rainy season) and between time of capture (06 hours to 13 hours in the morning, vs. 13 hours to 18 hours in the afternoon). A routine for the effect was written with the program Statistics 101, v. 1.5.3 (<http://www.statistics101.net/>) whereby the data of the two groups to be contrasted are combined and resampled without replacement iteratively, the difference of the two resample means at each iteration is recorded and the percentage of

time that this difference is equal or exceed the observed difference is taken as the p-value. Data were also analyzed by a two-way analysis of variance with the program Past (Hammer *et al.*, 2001) that uses weighted means for unbalanced designs.

Because total length and/or fish weight may influence the contrasts in stomach content a correlation test (Pearson's product moment correlation) for the data set was run between those variables and stomach content. Bias corrected 95 % bootstrap confidence intervals were fitted to means.

A total of 42 stomachs were collected, of which 35 contained prey represented by a limited number of species (Table 1, one stomach was excluded as its stomach content weight was an outlier). The main diet item was dwarf herring (*Jenkinsia lamprotaenia*) (75.13 % of contents by weight of which about 80 % was concentrated in the dry season), a small pelagic fish. The diet of the rainbow runner (seven diet items, Table 1) in the Central Colombian Caribbean is sparse compared to that in the mid-tropical Atlantic, where 27 taxa were recorded (Vaske *et al.*, 2006), or in the Pacific Ocean with at least 22 species of prey (Bocanegra, 2007). One obvious reason is the difference in the number of stomachs with food analyzed, 35 in this study versus 192 in Vaske *et al.*, (2006) and 184 in Bocanegra (2007), as well as the much larger geographic area covered in Bocanegra (2007). Even so it is surprising that the local diet of *E. bipinnulata* was so heavily dominated by a single species (*Jenkinsia lamprotaenia*). The extent to which local biodiversity influences *E. bipinnulata* remains unclear.

Mean total length and mean fish weight were higher both in the rainy season and in the afternoon than in the dry season and in the morning, respectively (Table 2). However, mean stomach content in the dry season was not no statistically higher contrasted to mean stomach content in the rainy season ($p > 0.05$). Mean stomach content in the afternoon, on the other hand, was significantly higher than in the morning ($p < 0.05$). Pearson's product moment correlation between stomach content and total length and fish weight turned out to be close to zero (-0.005 with fish weight and -0.099 with total length). Thus, those variables do not exert an unduly influence in the above results.

Two-way anova confirms the results above: no significant difference was found in stomach content between seasons ($p > 0.05$) but a significant difference was found between times of capture ($p < 0.05$). Interestingly a marginally significant interaction between season and time of capture was found ($p = 0.08$). This was so because of a total of six empty stomachs five were found in the dry season thus generating a huge variance. On the other hand, these five stomachs corresponded to fishes captured in the morning hours, which explains the marginal interaction found between season and time of capture.

The fuller stomachs of *E. bipinnulata* in the dry season cannot be tracked back to increased primary production normally associated to upwelling phenomena. The supply of nutrients in the rainy season due to continental discharges is not much different than nutrients supplied by deep water (Ramírez, 1990). Thus the upwelling manifests just as lower sea temperatures and higher salinities compared to the rainy season, but does not significantly influence primary production (Franco-Herrera *et al.*, 2006).

On the other hand inspection of the diet matrix suggests that the main diet item, the dwarf herring (*J. lamprotaenia*), exhibits a seasonal presence in the stomachs of *E. bipinnulata* with maxima in the dry season. In balance, both the quantitative and qualitative observations of the diet of the rainbow runner suggest a correlation with climate.

The trophic level for *E. bipinnulata* found in this study (4.24) exceeded that reported by Dambacher *et al.*, (2010) for the Pacific Ocean (3.59), and may indicate that system-level biodiversity plays a role in both diet and trophic structure. A less diverse system might support a less complex trophic structure, which would allow *E. bipinnulata* to occupy a higher trophic niche (e.g. Duffy and Stachovicz, 2006).

Consumption to biomass ratio was 10.8, which means that *E. bipinnulata* eats its own weight 10.8 times a year. The difference in Q/B value between the original value in Opitz (1996), 7.9, and this study may be due in part to differences in the models used (a model by Palomares and Pauly, 1989, was used by Opitz; a model by Palomares and Pauly, 1998, was used in this study). In fact, García and Duarte (2002) found that the Q/B values estimated with the Palomares and

Table 1. Diet items of the rainbow runner, *Elagatis bipinnulata*, in the Colombian Central Caribbean. % weight is wet weight.

Diet item	Taxonomic group	% Weight
<i>Jenkinsia lamprotaenia</i>	Pisces: Clupeidae	75.13
Mysidacea	Crustacea: Peracarida	7.04
<i>Portunus anceps</i>	Crustacea: Decapoda	6.09
<i>Septoteuthis sepioidea</i>	Mollusca: Cephalopoda	5.43
<i>Coronis</i> sp.	Crustacea: Stomatopoda	2.21
<i>Cavolinia uncinata</i>	Mollusca: Gastropoda	2.07
Harpacticoida	Crustacea: Copepoda	2.01

Table 2. Stomach content (wet weight in g, including unidentifiable material), total length (cm) and body weight (wet weight, g) of the rainbow runner (*Elagatis bipinnulata*) by season and capture time in the central Colombian Caribbean. Confidence Intervals are bias corrected 95 % bootstrap confidence intervals.

	Number of Samples	Mean	Range	Confidence Interval
Stomach Content				
Dry Season	26	2.06	0.00-17.31	0.88-3.91
Rainy Season	15	0.94	0.00-4.83	0.46-1.72
Morning	24	0.60	0.00-4.00	0.27-1.07
Afternoon	17	3.14	0.09-17.31	1.46-5.77
Total length				
Dry Season	26	46.39	33.50-67.00	43.54-49.55
Rainy Season	15	52.99	43.30-65.70	49.95-56.25
Morning	24	46.93	40.90-65.70	45.04-49.52
Afternoon	17	51.46	33.50-67.00	46.48-55.84
Fish Weight				
Dry Season	26	773.80	375.00-2125.00	634.36-957.45
Rainy Season	15	943.75	531.25-2000.00	777.08-1162.50
Morning	24	705.95	468.75-2000.00	603.39-868.23
Afternoon	17	1020.96	375.00-2125.00	823.63-1248.90

Pauly (1989) model were consistently smaller than those estimated with Palomares and Pauly (1998) model. Once recalculated ($Q/B = 9.8$) the difference gets shorter, but Opitz (1996) value is still below the Q/B in this study. These values suggest that *E. bipinnulata* is not a voracious fish. García and Duarte (2002) found a mean trophic value for carangids of 12.3, at mid way in a list of over 60 families.

The rainbow runner eats significantly more in the afternoon than in the morning (Table 2). Fuller stomachs in the afternoon suggest that the rainbow runner is eating during day light, at least in our study area. Further observations are needed to discern among the web of variables and time dynamics involved and to decide whether causal relationships exists and which are them in the shaping of the diet of this fish.

ACKNOWLEDGMENTS

Comments by two anonymous reviewers helped to improve the manuscript.

REFERENCES

- Abarca-Arenas LG, Franco-López J, Peterson MS, Brown-Peterson NJ, Valero-Pacheco E. Sociometric analysis of the role of penaeids in the continental shelf food web off Veracruz, México based on by-catch. *Fish Res.* 2007; 87(1):46-57. DOI: <http://dx.doi.org/10.1016/j.fishres.2007.06.019>
- Andrade CA. Las corrientes superficiales en la cuenca de Colombia observadas con boyas de deriva. *Rev Acad Colomb Cien Fis Nat.* 2001;25(96):321-335.
- Arizmendi-Rodríguez DI, Abitia-Cárdenas LA, Galván-Magaña F, Trejo-Escamilla I. Food habits of sailfish *Istiophorus platypterus* off Mazatlán, Sinaloa, México. *Bull Mar Scie.* 2006;79(3):777-791.
- Bocanegra N. Relaciones tróficas de los peces pelágicos asociados a la pesquería del atún en el océano Pacífico oriental. [Tesis Doctorado]. Instituto Politécnico Nacional, La Paz, Baja California Sur, México; 2007.
- Brunnschweiler JM, Sazima I. Swift swimming free fish as host of small juvenile shark suckers. *Coral Reefs.* 2010; 29(4):843. DOI: 10.1007/s00338-010-0655-9
- Castro E, Grandas Y, García C. Conocimiento pesquero tradicional: Aplicación del Análisis de Consenso Cultural para la evaluación y el manejo de la pesquería artesanal del la Isla de San Andrés, Colombia. *Proc Gulf Carib Fish Inst.* 2007;58:117-122.
- Criales-Hernández MI, García CB, Wolff M. Flujos de biomasa y estructura de un ecosistema de surgencia tropical en La Guajira, Caribe colombiano. *Rev Biol Trop.* 2006; 54(4):1257-1282.
- Dambacher MJ, Young JW, Olson RJ, Allain V, Galván-Magaña F, Landsdell M.J, *et al.* Analyzing pelagic food webs leading to top predators in the Pacific Ocean: A graph-theoretic approach. *Prog Oceanogr.* 2010;86(1-2):152-165. DOI: <http://dx.doi.org/10.1016/j.pocean.2010.04.011>
- Delgadillo O, García CB, Correa J. Dinámica temporal de la asociación de peces en dos arrecifes artificiales del Golfo de Morrosquillo, Caribe colombiano. *Actual Biol.* 2004; 26(81):219-230.
- Delgadillo-Garzón O, García CB. Impacto de dos arrecifes artificiales en la pesca artesanal diurna del golfo de Morrosquillo, Caribe de Colombia. *Rev Biol Trop.* 2009; 57(4):993-1007.

- Duffy JE, Stachowicz JJ. Why biodiversity is important to oceanography: potential roles of genetic, species and trophic diversity in pelagic ecosystem processes. *Mar Ecol Progr Ser.* 2006;311:179-189.
- Franco-Herrera, A, Castro L, Tigrero P. Plankton dynamics in the South-Central Caribbean Sea: Strong seasonal changes in a coastal tropical system. *Carib J Scie.* 2006; 42(1):24-38.
- García CB, Duarte LO. Consumption to biomass (Q/B) ratio and estimates of Q/B-predictor parameters for Caribbean fishes. *Naga.* 2002;25(2):19-31.
- García CB. Conocimiento tradicional: Lo que los pescadores artesanales del Caribe colombiano tienen para decirnos. *Panam J Aquat Sci.* 2010;57(1):78-90.
- García CB, Contreras CC. Trophic levels of fish species of commercial importance in the Colombian Caribbean. *Rev Biol Trop.* 2011;59(3):1195-1203.
- Garzón J, Acero A. Notas sobre la pesca y los peces comerciales de la isla de Providencia (Colombia), incluyendo nuevos registros para el Caribe Occidental. *Caribb J Scie.* 1983;19(3-4):9-19.
- Good PI. Resampling methods: a practical guide to data analysis. 3rd Edition Boston: Birkhäuser; 2006.
- Grijalba-Bendeck M, Castañeda-Moya E, Acero A. Estructura de un ensamblaje íctico asociado a fondos duros en el Caribe colombiano empleando la técnica del censo visual estacionario (CVE). *Actual Biol.* 2004;26(81):197-211.
- Hammer Ø, Harper DAT, Ryan PD. PAST: Paleontological Statistics Software Package for Education and Data Analysis. *Palaeont Electron.* 2001;4(1):9.
- Hensley VI, Hensley DA. Fishes eaten by sooty terns and brown noddies in the Dry Tortugas, Florida. *Bull Mar Sci.* 1995;56(3):813-821.
- Hiatt RW, Strasberg DW. Ecological relationships of the fish fauna on coral reefs of the Marshall Islands. *Ecol Monogr.* 1960;30(1):65-127. DOI: <http://dx.doi.org/10.2307/1942181>
- Opitz S. Trophic interactions in Caribbean coral reefs. ICLARM Tech Rep. Manila, Philippines. 1996.43, 341p.
- Palomares ML, Pauly D. A multiple regression model for predicting the food consumption of marine fish populations. *Mar Freshw Res.* 1989;40(3):259-273.
- Palomares ML, Pauly D. Predicting fish consumption of fish populations as functions of mortality, food type, morphometrics, temperature and salinity. *Mar Freshw Res.* 1998;49(5):447-453.
- Pauly D. A simple method for estimating the food consumption of fish population from growth data and food conversion experiments. *Fish Bull.* 1986;84(4):827-840.
- Ramírez, G. Evaluación de parámetros hidrográficos y su relación con la surgencia en las aguas costeras. En: Díaz JM, editor. Estudio ecológico integrado de la zona costera de Santa Marta y Parque Nacional Natural Tayrona. INVEMAR, Informe Final, Santa Marta, Colombia; 1990, p. 19-54
- Reyes-Nivia MC, Rodríguez-Ramírez A, Garzón-Ferreira J. Peces asociados a formaciones coralinas de cinco áreas del Caribe colombiano: listado de especies y primeros registros para las áreas. *INVEMAR.* 2004;33:101-115.
- Romanov EV. Bycatch in the tuna purse-seine fisheries of the western Indian Ocean. *Fish Bull.* 2001;100(1):90-105.
- Saito H, Yamashiro R, Ishihara K, Xue C. Lipids of three highly migratory fishes: *Euthynnus affinis*, *Sarda orientalis* and *Elagatis bipinnulata*. *Bioscie Biotech Bioch.* 1999;63(11):2028-2030.
- Senta T, Kimura M, Kambara T. Predation of fishes on open-ocean species of sea-skaters (*Halobates* spp). *Gyoryugaku Zasshi.* 1993;40(2):193-198.
- Shimose T, Shono H, Yokawa K, Saito H, Tachihara K. Food and feeding habits of blue marlin, *Makaira nigricans*, around Yonaguni Island, southwestern Japan. *Bull Mar Sci.* 2006;79(3):761-775.
- Shimose T, Yokawa K, Saito H, Tachihara K. Seasonal occurrence and feeding habits of black marlin, *Istiompax indica*, around Yonaguni Island, southwestern Japan. *Ichthyol Res.* 2008;55:90-94. DOI: 10.1007/s10228-007-0004-3
- Vaske T, Vieira FH, Lessa RP. Pesca e hábitos alimentares do peixe-rei, *Elagatis bipinnulata* (Quoy and Gaimard, 1825) (Pisces: Carangidae) no Arquipélago de Sao Pedro e Sao Paulo, Brasil. *Arquiv Ciên Mar.* 2006;39:61-65.
- Watkins WA, Daher MA, Ristrup KF, Notabartolo di Sciara G. Fishing and acoustic behavior of Fraser's Dolphin (*Lagenodelphis hosei*) near Dominica, Southeast Caribbean. *Caribb J Scie.* 1994;30(1-2):76-82.
- Yukinobo I. Maturity and feeding habits of rainbow runner, *Elagatis bipinnulata* in the western Pacific Ocean. *J Sch Mar Sci Tech Tokai Univ.* 1998;46:33-40.

