

NOTA BREVE

BLOOMS OF EPHEMERAL GREEN ALGAE IN SAN ANDRES ISLAND, INTERNATIONAL BIOSPHERE RESERVE *SEAFLOWER*, SOUTHWESTERN CARIBBEAN

Florecimientos de algas verdes efimeras en la isla de San Andrés, Reserva Internacional de Biosfera *Seaflower*, Caribe suroccidental

Brigitte GAVIO¹, Jose Ernesto MANCERA PINEDA¹

¹Universidad Nacional de Colombia, sede Bogotá, Carrera 30 Calle 45. Departamento de Biología, edificio 421, oficina 106, Bogotá, Colombia.

For correspondence: bgavio@unal.edu.co

Received: 7 October 2014; **Returned for revision:** 8 November 2014; **Accepted:** 9 December 2014.

Associate Editor: Rafael Riosmena Rodríguez.

Citation / Citar este artículo como: Gavio B, Mancera Pineda JE. Blooms of ephemeral green algae in San Andres island, International Biosphere Reserve Seaflower, Southwestern Caribbean. Acta biol. Colomb. 2015;20(2):259-262. doi: <http://dx.doi.org/10.15446/abc.v20n2.46062>

ABSTRACT

We report the presence of persistent blooms of *Chaetomorpha linum* in San Andres island, Southwestern Caribbean, during the year 2013.

Keywords: blooms, *Chaetomorpha linum*, Colombia.

RESUMEN

Reportamos la presencia de florecimientos persistentes del alga verde *Chaetomorpha linum* en la isla de San Andrés, Caribe suroccidental.

Palabras clave: *Chaetomorpha linum*, Colombia, florecimientos.

Excessive biomass of macroalgae is a common symptom of nutrient enrichment in seagrass and coral reef ecosystems, and their frequency has increased over the last decades (Lapointe *et al.*, 2004; Smith *et al.*, 2005). Macroalgal blooms can physically overgrow seagrasses and adult corals, inhibit recruitment of juvenile corals, reduce light availability, lead to hypoxia and/or anoxia, and result in greatly diminished fisheries and biological diversity (Lapointe *et al.*, 2004, Thomsen and Wernberg 2009; Rasmussen *et al.*, 2012).

Because of these detrimental effects, excessive biomass of macroalgae is considered Harmful Algal Blooms (ECOHAB, 1995).

These ephemeral blooms usually involve one or more species of early successional, opportunistic green algae, belonging to the genera *Ulva*, *Chaetomorpha* or *Cladophora* (Smith *et al.*, 2005).

In the Caribbean, recurrent blooms of the green algal species *Chaetomorpha linum* have been reported in Jamaica, overgrowing the fore reef communities on fringing reefs in the Negril Marine Park, and on seagrass beds in the Florida Keys (Lapointe *et al.*, 2004). These blooms have been attributed to nitrogen enrichment of the coastal waters, due to land inputs from sewage or agricultural runoff (Linton and Warner 2003; Lapointe *et al.*, 2004; Thomsen and Wernberg, 2009). In Jamaica, blooms of *Chaetomorpha linum* normally occur during the summer months, and Lapointe *et al.*, (1999) attribute these blooms to groundwater NO₃⁻ enrichment of the fore reef.

San Andres island is part of the International Biosphere Reserve *Seaflower* in the Southwestern Caribbean, and, with a resident population of 70069 habitants (DANE, 2011), equivalent to 2802.76 hab/km², it is one of the most populated island in the Caribbean (Díaz *et al.*, 1996); furthermore, the tourist affluence is high and constant throughout the year, increasing the anthropogenic pressure on the coastal ecosystems nearshore. The sewage system covers only 8% of the resident population (SIGAM/CORALINA, 2004), without any treatment prior to its disposal into the ocean. The great majority of



the population wastewater is disposed in septic tanks, but its leaking is impacting heavily its nearshore waters and ecosystems (Marín and Cadavid, 2001).

From February to May 2013, and again from September to October of the same year, San Andres experienced unusual blooms of a filamentous green alga, identified as *Chaetomorpha linum* (Fig. 1). The alga presented the typical

appearance of the species (Littler and Littler, 2000) with filaments up to 50 cm long. *Chaetomorpha* smothered seagrass beds (Fig. 2) before being washed ashore (Fig. 3). The alga formed large mats or “pillows” with sediment trapped in it (Fig. 2, arrows), which attached and moved upon seagrass for days, before ending on the beaches. The sediment-coated mats (Fig. 2) may have a scouring effect on




Figure 1. Filament of *Chaetomorpha linum*. Scale bar: 200 mm



Figure 2. Mats of *Chaetomorpha linum* smothering seagrass leaves.



Figure 3 *Chaetomorpha linum* washed ashore. Spratt Bight, San Andres island, Colombia, May 10th 2013.

seagrass leaves. The algae were not floating on the surface, but they were always found tumbling along the substrate, sometimes attaching to seagrass leaves and shoots; this behavior has already been reported, for the species, in other locations (Flindt *et al.*, 2007). These mats, beside the scouring effect, reduce light availability for seagrass shoots (Krause-Jensen *et al.*, 1996; McGlathery, 2001), increase the sediment organic matter load, inducing the risk of anoxia and sulfide intrusion into meristematic areas of seagrasses, therefore restricting seagrass growth (Nelson and Lee, 2001; Thomsen and Wernberg, 2009; Han and Liu, 2014). On the adjacent coral reefs, the presence of the mats was not observed.

In Jamaica, the first observed macroalgal blooms in shallow coastal waters were mainly due to *Chaetomorpha linum*, which smothered coral reefs and seagrass beds (Lapointe and Thacker, 2002). According to Lapointe *et al.*, (2011) green tides composed of *Chaetomorpha linum* expanded in areas most influenced by sewage discharge.

Although we did not pursue water analysis on nutrient content, it is probable that the progressive nutrification of the coastal waters of the island, which has been in act for at least the past 15 years (Gavio *et al.*, 2010) is responsible for these conspicuous blooms. Eutrophication of shallow coastal bays typically causes a shift in dominance from seagrasses and perennial macroalgae to ephemeral, bloom-forming algae (McGlathery *et al.*, 2007).

In order to avoid the appearance of these blooms in the future and its negative effects on the nearshore ecosystems, it is urgent to improve the management of wastewater discharge and reduce the nutrient input on the coastal waters.

ACKNOWLEDGEMENTS

This contribution was supported by the Universidad Nacional de Colombia sede Caribe. We wish to thank Fredy Duque and Natalia Rincón for their valuable help.

REFERENCES

- DANE [Internet]. Estimaciones de población 1985-2005 y proyecciones de población 2005-2020 total municipal por área. 2011. [Consulted 12 May 2011]. Available from: <https://www.dane.gov.co/index.php/es/poblacion-y-registros-vitales/proyecciones-y-series-de-poblacion/proyecciones-de-poblacion>
- Díaz JM, Díaz-Pulido G, Garzón-Ferreira J, Geister J, Sanchez J, Zea S. Atlas de los arrecifes coralinos del Caribe colombiano. I complejos arrecifales oceánicos. INVEMAR, Santa Marta; 1996. p. 29.
- ECOHAB. Introduction. In: Anderson DM, editor. The Ecology and Oceanography of Harmful Algae Blooms A National Research Agenda. WHOI, Woods Hole, MA, USA; 1995. p. 7-10.
- Flindt MR, Pedersen CB, Amos CL, Levy A, Bergamasco A, Friend PL. Transport, sloughing and settling rates of estuarine macrophytes: Mechanisms and ecological implications. Cont Shelf Res. 2007;27(8):1096-1103. Doi: <http://dx.doi.org/10.1016/j.csr.2006.08.011>
- Gavio B, Palmer-Cantillo S, Mancera E. Historical analysis (2000–2005) of the coastal water quality in San Andrés Island, SeaFlower Biosphere Reserve, Caribbean Colombia. Mar Pollut Bull. 2010;60(7):1018–1030. Doi: <http://dx.doi.org/10.1016/j.marpolbul.2010.01.025>
- Han Q, Liu D. Macroalgae blooms and their effects on seagrass ecosystems. J Ocean Univ China. 2014;13(5):791-798. Doi: <http://dx.doi.org/10.1007/s11802-014-2471-2>
- Krause-Jensen D, McGlathery K, Rysgaard S, Christensen PB. Production within dense mats of the filamentous macroalga *Chaetomorpha linum* in relation to light and nutrient availability. Mar Ecol Progr Ser. 1996;134:207-216. Doi: <http://dx.doi.org/10.3354/meps134207>
- Lapointe BE. Simultaneous top-down and bottom-up forces control macroalgal blooms on coral reefs (reply)

- to the comments by Hughes *et al.*). Limnol Oceanogr, 1999;44(6):1586-1592.
- Lapointe BE, Thacker K. Community-based water quality and coral reef monitoring in the Negril Marine Park, Jamaica: Land-based nutrient inputs and their ecological consequences. In: Porter J, Porter K, editors. The Everglades, Florida Bay, and Coral Reefs of the Florida Keys: An Ecosystem Sourcebook. CRC Press, Boca Raton, FL, USA; 2002. p. 939-963.
- Lapointe BE, Barile PJ, Matzie WR. Anthropogenic nutrient enrichment of seagrass and coral reef communities in the Lower Florida Keys: discrimination of local versus regional nitrogen sources. J Exp Mar Biol Ecol. 2004;308:23-58. Doi: <http://dx.doi.org/10.1016/j.jembe.2004.01.019>
- Lapointe BE, Thacker K, Hanson C, Getten L. Sewage pollution in Negril, Jamaica: effects on nutrition and ecology of coral reef macroalgae. Chin J Oceanol Limnol. 2011;29(4):775-789. Doi: <http://dx.doi.org/10.1007/s00343-011-0506-8>
- Linton DM, Warner GF. Biological indicators in the Caribbean coastal zone and their role in integrated coastal management. Ocean Coast Manage. 2003;46(3-4):261-276. Doi: [http://dx.doi.org/10.1016/S0964-5691\(03\)00007-3](http://dx.doi.org/10.1016/S0964-5691(03)00007-3)
- Littler DS, Littler MM. Caribbean reef plants. Offshore Graphics. Washington D.C; 2000. p. 542.
- Marín B, Cadavid B. Descripción general de la calidad de las aguas marinas de Colombia. Caracterización de la calidad de las aguas marinas. Informe técnico. Santa Marta; 2001. p. 80.
- McGlathery KJ. Macroalgal blooms contribute to the decline of seagrass in nutrient-enriched coastal waters. J Phcol. 2001;37:453-456. Doi: <http://dx.doi.org/10.1046/j.1529-8817.2001.037004453.x>
- McGlathery KJ, Sundbäck K, Anderson IC. Eutrophication in shallow coastal bays and lagoons: the role of plants in the coastal filter. Mar Ecol Progr Ser. 2007;348:1-18. Doi: <http://dx.doi.org/10.3354/meps07132>
- Nelson TA, Lee A. A manipulative experiment demonstrates that blooms of the macroalga *Ulvaria obscura* can reduce eelgrass shoot density. Aquat Bot. 2001;71(2):149-154. Doi: [http://dx.doi.org/10.1016/S0304-3770\(01\)00183-8](http://dx.doi.org/10.1016/S0304-3770(01)00183-8)
- Rasmussen JR, Olesen B, Krause-Jensen D. Effects of filamentous macroalgae mats on growth and survival of eelgrass, *Zostera marina*, seedlings. Aquat Bot. 2012;99:41-48. Doi: <http://dx.doi.org/10.1016/j.aquabot.2012.01.005>
- SIGAM/CORALINA [on line]. Agenda ambiental de San Andrés Islas. 2004. Available at: http://www.coralina.gov.co/archivos/Sigam_CAPITULO%20_I_Perfil_Ambiental_1.pdf
- Smith JE, Runcie JW, Smith CM. Characterization of a large-scale ephemeral bloom of the green alga *Cladophora sericea* on the coral reefs of West Maui, Hawai'i. Mar Ecol Progr Ser. 2005;302:77-91. Doi: <http://dx.doi.org/10.3354/meps302077>
- Thomsen MS, Wernberg T. Drift algae, invasive snails and seagrass health in the Swan River: patterns, impacts and linkages. Report no. CMER-2009-02 from the Centre for Marine Ecosystems Research, Edith Cowan University; 2009. 105 p.