

Antifungal activity of acidic electrolyzed water against strawberry postharvest molds (*Fragaria x ananassa* Duch cv. *Camarosa*)

Actividad antifúngica del agua electrolizada ácida contra mohos postcosecha de fresa (*Fragaria x ananassa* Duch cv. *Camarosa*)

Beatriz Elena Guerra Sierra^{1*}, Adriana Ximena Sandoval Meza², Liliana Torcoroma García Sánchez³

^{1,2} Universidad de Santander, Facultad de Ciencias Exactas, Naturales y Agropecuarias. ^{1,2,3} Grupo de Biotecnología Agroambiente y Salud-Microbiota. Calle 70 No. 55-210, C.P. 680003, Bucaramanga (Santander).

*Autor para correspondencia: bguerra@udes.edu.co

Rec.: 2019-03-04 Acep.: 2019-07-03

Abstract

The aim of this work was to assess the in vitro antifungal activity of acidic electrolyzed water (AEW) and the required contact time to inhibit the development of post-harvest molds common in strawberry fruit. Ninety-four strawberry samples were used. A five-fold assay in a randomized design was carried out using standard inoculum of fungal conidia, which were subjected to an array of contact times and AEW concentrations, using a controlled agitation (120 r.p.m) at 23°C. The conidia treated with AEW were grown on potato dextrose agar and their germination was evaluated. Mycelial growth of the treated samples was measured and compared against conidia control samples treated with distilled water instead of AEW. *Botrytis cinerea* and *Rhizopus stolonifer* were the most commonly isolated species at 54.25% and 28.72%, respectively. AEW inhibited conidia germination after 7 minutes of exposure. The AEW concentration required to inhibit conidia germination was 10 ppm and 50 ppm for *R. stolonifer* and *B. cinerea* respectively, a variance possibly due to the constitution of their cellular walls. A short exposure of AEW is able to inhibit post-harvest fungi growth, and is thus recommended as an efficient, economical, and environmentally sustainable sanitation agent, with the ability of extending the strawberry post-harvest consumption period.

Keywords: Acidic electrolyzed water; antifungal; *Botrytis cinerea*; post-harvest molds; *Rhizopus stolonifer*.

Resumen

El objetivo de este trabajo fue evaluar la actividad antifúngica in vitro del agua electrolizada ácida (AEA) y el tiempo de contacto requerido para inhibir el desarrollo de mohos postcosecha comunes en fresa. Se utilizaron 94 frutos dañados y sanos. En un ensayo al azar con cinco repeticiones, utilizando un inóculo estándar de conidios fúngicos, tratados durante una serie de tiempos de contacto y diferentes concentraciones de AEA, utilizando una agitación controlada (120 r.p.m) a 23 °C. Los conidios tratados con AEA se cultivaron en agar papa-dextrosa para evaluar su germinación. Se midió el crecimiento micelial de las muestras tratadas y se comparó con las muestras de control de conidias tratadas con agua destilada en lugar de AEA. *Botrytis cinerea* y *Rhizopus stolonifer* fueron las especies más comúnmente aisladas con 54.25% y 28.72%, respectivamente. AEA inhibió la germinación de conidios después de 7 minutos de exposición. La concentración de AEA requerida para inhibir la germinación de los conidios fue de 10 ppm y 50 ppm para *R. stolonifer* y *B. cinerea*, respectivamente, las diferencias podrían estar relacionadas con la constitución de sus paredes celulares. Una exposición corta de AEA es capaz de inhibir el crecimiento de hongos poscosecha y, por tanto, con base en los resultados en este trabajo es posible recomendar al AEA como higienizante eficiente, económico y ambientalmente sostenible, para uso en frutos poscosecha y así prolongar el periodo de consumo.

Palabras clave: Agua ácida electrolizada; antifúngicos; *Botrytis cinerea*; hongos poscosecha; *Rhizopus stolonifer*.

Introduction

Strawberries can be damaged by a series of molds that attack the fruit, such as *Botrytis cinerea*, the main pathogen responsible for the disease known as gray rot. This condition can occur as a quiescent or latent infection wherever strawberries are grown, appearing as a light brown or yellowish stain on the calyx, which expands within a few days to cover the surface of the fruit with a gray, dusty appearance. Gray rot is one of the most significant strawberry diseases worldwide. The cost of *B. cinerea* damage is very difficult to determine, as it occurs at several stages of the production and distribution chain (Dean R et al., 2012). The annual cost of treatments to combat *B. cinerea* in all vulnerable crops and all countries, has been estimated at up to €540 million, which is about 10% of total world expenditure on fungicides; in Spain around €2 million are spent on fungicides for protecting strawberry crops (Fernández-Acero et al., 2007). The fungus grows best between 15 - 25°C, however, it is able to do so at temperatures as low as 0°C, making it an important pathogen of stored products. Strawberries are also attacked by *R. stolonifer*, a common wound pathogen which affects a wide range of fruits and vegetables, causing a rapidly-spreading watery soft rot. Once the pathogen has entered the wounded tissue, its mycelia spread rapidly (Holz, G et al., 2007). This fungus is naturally found in the soil as a saprophyte, consuming organic residues and air. Other common pathogens that affect strawberry fruit are *Mucor spp*, *Colletotrichum spp*, and *Penicillium spp*.

Current strategies to control phytopathogenic fungi include the use of acid, alkali, and chloride compounds, and/or ozone, but these compounds are less than ideal, as they are both costly and environmentally unfriendly, and because they may form carcinogenic compounds as chemical derivatives. Additionally, they are ineffective against some pathogen strains which are resistant to fungicides. The development of sanitizing agents to eliminate pathogens from harvested agricultural products has been identified by the food industry as one of the key factors in their risk assessment and critical points analysis, (Zacharia, et al., 2010).

A new concept in disinfection is electrolyzed water, which was first developed in Japan (Al-Haq et al., 2005) and is now recognized as an effective sanitation agent with a high antimicrobial capacity. Acidic electrolyzed water's primary components are salt and water, and is produced using specialized electrolysis equipment, passing the solution through an electrolytic cell where

the anode and cathode are separated by an electrolytic membrane. When continuous voltage is applied, the negatively charged ions (Cl⁻, OH⁻), move towards the anode, where oxygen molecules release electrons while producing O₂, hypochlorite ions, (ClO⁻), hypochlorous acid (HOCl), and hydrochloric acid (HCl). The positively charged ions move to the cathode, where H₂ and NaOH are produced (Huang et al., 2008).

The disinfection principle of electrolyzed water is based on three factors: its pH, the chloride concentration available, and its redox potential, of which the latter represents the main cause of microorganism inactivation due to the severe and irreversible damage it causes to external and internal cell membranes (Hao et al., 2012). Electrolytically generated hypochlorous acid from the NaCl solution is considered to be a side-effect-free wide spectrum antimicrobial alternative.

Evaluations of its efficacy at disinfection and pathogen reduction or elimination have found it to demonstrate high redox potential, which prevents the microorganism by entering the fruit cells, disrupts its protein synthesis, and causes oxidation of its amino acid sulfur groups. This last effect, in turn, forms toxic derivatives and leads to enzyme destruction as a result of the consequent metabolism imbalance (Bialka et al., 2004). Nevertheless, the electrolyzed water does not affect the strawberry's tissue, surface texture, color, or general appearance (Callejas et al., 2010) which is of great importance in maintaining the organoleptic properties of the fruit. In order to take advantage of these beneficial properties, the question arises: what is the minimum AEW concentration and contact time required to inhibit the growth of the *B. cinerea* and *R. stolonifer* strains found to represent the most common fungi in the samples of strawberries acquired from field crops and supermarkets and used in this research?

Materials and methods

Production of electrolyzed water solution

The acidic electrolyzed water was generated from a 5% NaCl solution using a membrane exchange device at LTec-LimpiooTec-Colombia. The resulting solution showed a pH of 6.6±0.1 (Thermo Scientific Orion 3 Star), a redox potential (ORP) of 760 mV (Thermo Scientific Orion Star A221), and a chlorine concentration as hypochlorous acid of 360 ppm (Faro-300 Merck). The electrolyzed water thus produced was stored in a sealed dark bottle at 25 °C, and utilized for this study within 8 hours of its production.

Fungal species identification

Samples of unripened strawberry fruit were obtained from farms around the city of Pamplona (Colombia) which operate using local farming practices. Another sample consisted of imported strawberry fruit purchased from a local market. Both samples were of the strawberry variety *Fragaria x ananassa* Duch cv. *Camarosa*.

Random subsamples of healthy fruit (free of pathogen symptoms and/or decay caused by fungi and bacteria) were chosen; from each group (divided by origin), and placed into humid chambers formed by covering sterile glass containers with plastic. The samples were stored at room temperature (± 25 °C), and exposed to natural daylight, 8 h light and 16 h darkness for three days. The fruit was checked daily to evaluate the presence of rot and mycelial growth. Next, 94 fruit samples of 0.5 cm x 0.5 cm each were removed and taken for assays. Both healthy and rotten samples were cultured at 25 °C in Potato Dextrose Agar, supplemented with 0.01% chloramphenicol. These samples were subcultured to obtain pure isolated fungi cultures. Morphological fungi identification was carried out using macroscopic characteristics as growth, morphology, colony color and appearance. The cultures were then mounted in drops of lactophenol on microscope slides for examination under a light microscope of their microscopic characteristics (conidiophores and conidia morphology, conidiogenesis type, and asexual structures. Preliminary morphological identification of the fungi was carried out with the support of different taxonomic keys (Ainsworth et al., 1973, Cary et al. 2003, Coley-Smith, 1980, Domsch, et al. 1980, Nelson et al., 1983, O'Donnell, 1979, Seifert and Gams, 2011 and Zhou, 2014).

DNA extraction and quantification

Mycelia were grown in malt extract broth (20 g malt extract, 1000 mL distilled water). Fungal genomic DNA was extracted from pure cultures using the as described by Barnes et al. (2001). DNA concentration was estimated by gel electrophoresis and adjusted to 1 ng μl^{-1} . The fungal ITS region was amplified using the primers ITS1 and ITS4 (White et al., 1990). The electrophoresis was carried out at 75 volts using 0.5X TBE as a buffer, until the dye had migrated. The DNA fragments generated were observed under UV light on an image documentation system (Enduro GDS-1365), and the DNA pellets were stored in 25 μl of distilled water and preserved at -20 °C for later analysis.

DNA PCR amplification

Specific primers based on internal transcribed spacer sequences (ITS) region of the rDNA was amplified using primer ITS1 (5'-CCGTTGGTGAACCAGCGG-3') and primer ITS4 (5'-TCCTCCGCTTATTGATATGC-3') for *Rhizopus* spp. White et al. (1990), for *Botrytis* spp., (5'-AGCTCGAGAGAGATCTCTGA-3'), and (5'-CTGCAATGTTCTGCGTGAA-3') (Rigotti et al., 2002).

The PCR amplifications were performed on a thermocycler (Eppendorf-Mastercycler, Germany), using the aforementioned initiators for each fungus, with a positive DNA control of *Mucor* spp. and a negative control which contained only the reactant. For the amplification, 5 microliters of each of the isolated DNA were used in 25 μl of the PCR mixture. The reaction was carried out in 10 mM of Tris -HCl (pH8.0), 2 mM MgCl, 0.2mM dNTPs, 20mM of each initiator, and 1.5 U of Taq DNA Polymerase (TaKaRa Taq™ DNA Polymerase) The amplifications incorporated an initial denaturation at 95°C for 3 minutes, 30 denaturation cycles at 95°C for 1 minute, alignment at 60°C for 1 minute, and elongation at 72°C for 2 minutes, followed by a final extension at 72°C for 5 minutes. The PCR products were passed through a 1.2% (w/v) agar gel electrophoresis chamber and colored by Ethidium bromide at 90 volts for 50 minutes before being observed under a UV transilluminator (Enduro GDS-1365).

The PCR-amplified DNA fragments were purified using the column method according to the Qiagen Kit manufacturer's specifications, then prepared and sent to a Korean laboratory service for sequencing. The strawberry fungi ITS sequences were analyzed using the BLASTn database and compared to the ones deposited in the GenBank at the National Center for Biotechnological Information (NCBI). The ITS1 and ITS4 sequences were aligned using ClustalW of Lasergene's MegAlign 7.0 (DNASTAR Inc. USA) software. In addition to identifying the fungi, the alignment analysis was useful in confirming the information generated by BLASTn, including highest score, (max score), best scope (query cover) and highest identity.

Preparation of inoculums

Conidia were obtained from twelve days' growth of PDA and Sabouraud cultures, which had been incubated at 25°C, in light and darkness cycles of 12 hours each. Conidia were collected from the culture dishes using a loop and transferred to 10 ml vials with sterile distilled water and 0.05 % of Tween 80. The resulting enriched conidia solutions of each fungal species were passed

through a glass filter to obtain a mycelium-free suspension and their concentrations were adjusted to 10^6 conidia/mL using a Neubauer chamber (adjusted of the methodology proposed by Newmeyer, D. (1990).

Preparation of the acidic electrolyzed water fungicidal test

In order to evaluate the fungicidal effect of AEW, solutions of 10, 25, 50, 100, and 200 ppm were prepared by dilution from a stock solution of 360-ppm chlorine as hypochlorous acid, pH 6.6 ± 0.1 . From each resulting concentration, 9 ml was then transferred to sterilized glass test tubes. This process was carried out in sets of five. Each test tube was inoculated with 1 ml of 1×10^6 conidia/ml suspension solution from one of the two fungal species isolated. The solutions were permitted to react for 1, 3, 5, 7, and 10 minutes under constant shaking at 100 rpm (Shaker orbital Mazzini), with 1.0% Tween solution (Science Med). One milliliter of each spore suspension thus obtained was plated on Sabouraud dextrose agar medium, with 0.01% cloranfenicol, in 9 cm medium petri dishes. These dishes were sealed and incubated at 25°C for 8 days, and were monitored, with their daily fungal growth measured in centimeters. An identical procedure was carried out using sterile distilled water as a fungal growth control for each of the evaluated strains.

Observation and measurement of radial mycelial growth

Observation and measurement was carried out for seven days, with radial growth recorded for each treatment at the same time each day. To determine the fungal radial growth, the linear distance from the inoculation point to the mycelial border was measured and compared to a mycelial control. The experiment defined 100% growth as the point at which the mycelial border reached the 9 cm edge of the dish. The mycelial growth control test was treated with sterile distilled water, and the percentage of mycelial growth inhibited was used to determine the effective concentration and duration of AEW treatment. The following equation was established to calculate the percentage of growth inhibited:

$$\text{Inhibition growth (\%)} = [(dc - dt)/dc] \times 100(\%) \quad \text{Ec. 1}$$

where, dc = the average diameter of the fungal colony in the control case, and dt = the average diameter of the fungal colony in the treatment group

Germination test and development of germ tube

To evaluate the length of the germ tube of *R. stolonifer* and *B. cinerea*, a conidial solution of 1×10^5 spores mL^{-1} was prepared in five concentrations of electrolyzed water (10, 25, 50, 100, 200 ppm). A contact time with the electrolyzed water of 1, 3, 5, 7 and 10 minutes was allowed (or left). Subsequently, 100 μL of each assay was taken and deposited in 4 excavations of the PDA plate medium culture; the plates will be incubated at 27°C and checked for 24 and 48 hours, to evaluate the growth of the germ tube under the light microscope (Nikon -eclipse Ni). Percent inhibition of spore germination was determined by counting 100 spores at ten randomly select points, in the clear field at 10x using Software Nis-elements 4.0 to estimate the averages of the germ tube measurements. A control in water sterile was carried out for each treatment.

Statistical design and analysis

This study adopted a randomized design to investigate the effects of various treatment parameters that is; the effect of five contact times and five concentrations of electrolyzed water on mycelial growth and development of germ tube as part of an antifungal treatment against *R. stolonifer* and *B. cinerea*. Experiments were independently conducted, in quintuplicate, for each portion of the study. Fungal growth following treatment with electrolyzed water (using an array of concentration and contact time parameters), was evaluated daily. Radial growth was reported in centimeters, and compared against a growth control for each fungal species, which used sterile water in place of the electrolyzed water applied to the conidia. The fungal growth phase was observed for a period of 8 days, after which time the control specimen had reached the full 9 cm diameter of the dish. Germ tube formation was evaluated during 24 and 48 hours and its measurement was expressed in microns. Data from each of the independently replicated trials was analyzed by variance tests performed using Statistical Analysis Systems (IBM® SPSS® Statistics V22.0). Tukey's range tests were used to determine if significant differences existed between the treatment groups, with values of $p < 0.05$ considered statistically significant.

Results and discussion

Fungal identification

There were obvious differences in the number of fungal isolates of rotten and healthy tissues, the 94 analyzed samples, *Botrytis* and *Rhizopus*

were the genera most commonly isolated (Table 1), therefore it was determined to perform its molecular identification. The DNA concentration after amplification was in general above 600 base pair for both fungi; a data, which matched the expected value. By alignment of the processed sequences using the BLASTn, it was possible to identify the relevant fungal species as *R. stolonifer* and *B. cinerea*, with a maximum identity of 100%.

Botrytis cinerea was isolated more frequently from fruit tissues in a decaying or rotten state, while *R. stolonifer* showed no significant differences in this regard. Besides the two species named, other fungi may affect the harvested fruit, as detailed in Table 1. Other fungi associated with post-harvest rotting of strawberries include: *Geotrichum candidum*, *Pilidium concavum*, *Pestalotia longisetula*, *Phoma sp*, *Sclerotinia sclerotiorum*, and *Cylindrocladium candelabrum* (Lopes, 2011). The type of fungi present would be influenced by the prevalent precipitation and temperature of the crop. Additionally, pathogens tend to survive if they reside within wounds in the host tissue, or if they are present as incipient, latent, or quiescent infections within it (Korsten and Wehner, 2003).

Table 1. Fungi isolated from harvested Strawberry fruit (*Fragaria x ananassa* Duch.

| Isolated fungi | Strawberry tissue | | | | Total number | Total percent |
|----------------------------|-------------------|-----|-----------------|-----|--------------|---------------|
| | Pamplona crops | | Farmers markets | | | |
| | R* | H** | R* | H** | | |
| <i>Aspergillus sp</i> | 2 | 1 | 1 | 1 | 5 | 5.31 |
| <i>Alternaria sp</i> | 1 | | 1 | | 2 | 2.12 |
| <i>Botrytis cinerea</i> | 26 | 5 | 13 | 7 | 51 | 54.25 |
| <i>Cladosporium sp</i> | 1 | — | — | 1 | 2 | 2.12 |
| <i>Geotrichum sp</i> | 2 | — | — | — | 2 | 2.12 |
| <i>Mucor spp</i> | — | — | — | 1 | 1 | 1.06 |
| <i>Colletotrichum</i> | 1 | — | — | — | 1 | 1.06 |
| <i>Penicillium sp</i> | | 2 | — | 1 | 3 | 3.19 |
| <i>Rhizopus stolonifer</i> | 8 | 7 | 5 | 7 | 27 | 28.72 |
| Total samples | 36 | 29 | 24 | 5 | 94 | 100.00 |

R*=rotten, H**=healthy. The data correspond to the average of independent replicas. The results show significant differences ($P < 0.05$) in the different settings and times.

Inhibition of fungal radial growth by electrolyzed water

The application of acidic electrolyzed water (pH: 6.6 ± 0.1) was found to be effective in decreasing the growth rate of *R. stolonifer* in each of the various treatments (Figure 1) compared to the control growth rate.

According to the results (Figure 1), significant differences ($P < 0.05$) in rate of growth between the various treatments can be ascribed to two factors, contact time and solution concentration. The time factor proved to play an important role in the three lowest AEW concentrations used in this study; however, it is noteworthy that as little as 10 ppm of AEW was sufficient to achieve the inhibition of conidial germination, and therefore the inhibition of mycelial development in vitro.




Figure 1 Percentage of growth (compared to control) of *R. stolonifer* subjected to AEW.

The results were somewhat different in the case of *B. cinerea*, in which AEW concentration was observed to exert a greater fungicidal effect than contact time. Statistical analysis showed that the growth inhibition percentage for *B. cinerea*, for any tested concentration of AEW, was not statistically different after an exposure of 7 minutes (Figure 2).




Figure 2 Percentage of growth (compared to control) of *B. cinerea* subjected to AEW.

Nonetheless, microbiological observation indicated that concentrations of at least 50 ppm were required to totally inhibit mycelial

growth within a contact time of 7 minutes. Although the analysis did not find statistically significant differences between the treatments, the microbiological results are important because even a minimal growth of mycelium is able to colonize and infect the vegetable tissues.

Statistical analysis of the contact time and AEW concentration factors showed a less marked decrease in the growth rate of *B. cinerea* in all the treatments tested (Figure 4). These results differed from those noted in the case of *R. stolonifer*, in which a rapid decrease in the growth rate was achieved using AEW; likely due to differences in the constitution of the fungal walls.

The evaluation of the germ tube for the two fungi of interest is shown in the Table 2, these results correlate with the mycelial growth of the two species (Figures 1 and 3). It is observed that for *B. cinerea*, the reduction of the germinal tube occurred using a concentration of 50 ppm of AEW.

These results coincide with some studies that have reported the sporicidal effect of AEW on Botrytis (Cravero et al., 2016; Feliziani et al., 2016; Guentzel et al. 2011). We found the total inhibition of fungal growth in *R. stolonifer* from 10 ppm AEW. Gomez et al., (2017), found a 100%

reduction of this fungal species, with 18 and 43 ppm of neutral electrolyzed water, however, studies related to concentration and sporicidal effectiveness are scarce.

The fungicidal effect of the electrolyzed water is due to the combined action of its high ORP, low pH, and concentration of available free chlorine in the form of molecular chlorine or hypochlorous acid (Al-Haq, 2005, Cui et al., 2009). The AEW used in this research was of pH 6.6. Other authors argue that chlorine is strongest in the HOCl form, and exhibits 80 times greater sanitizing power than OCl^- when the pH of the solution is between 5.0 and 6.5 (Cao et al., 2009). Other studies conducted with acid electrolyzed water in strawberry, have not reported elimination of fungi (Koseki et al., 2004), probably due to electrolyzed water conditions related to pH, free chlorine and ORP.

Contact time and concentration were two important factors that influenced the degree of antifungal activity achieved by acidic electrolyzed water, with a positive correlation for both. It is worth noting that, in this study, the specific contact times found for the AEW and fungal conidia could also be affected by the shaking




Figure 3 Interaction between acidic electrolyzed water factors (contact times and AEW concentration) for *R. stolonifer* growth rates.




Figure 4 Interaction between acidic electrolyzed water factors (contact times and AEW concentration) for *Botrytis cinerea* growth rates.

Table 2. Length of the germ tube (μm) of *B. cinerea* and *Rhizopus stolonifer* after treatments with AEW.

| AEW concentration | Contac time | <i>B. cinerea</i> | | <i>R. stolonifer</i> | |
|-------------------|-------------|-------------------|-------|----------------------|-------|
| | | 24 h | 48 h | 24 h | 48 h |
| Control | | 245.4 | 497.5 | 318.4 | 837.7 |
| | 1 | 244.2 | 478.4 | 315.2 | 800.1 |
| | 3 | 240.1 | 450.1 | 310.1 | 800.0 |
| | 5 | 220.4 | 126.4 | 77.4 | 72.1 |
| | 7 | 73.6 | 42.6 | 0 | 0 |
| 10 | 10 | 24.4 | 14.2 | 0 | 0 |
| | 1 | 425.4 | 216.4 | 394.7 | 385.4 |
| | 3 | 217.3 | 102.5 | 198.3 | 189.1 |
| | 5 | 192.8 | 98.4 | 45.1 | 40.1 |
| | 7 | 32.8 | 20.1 | 0 | 0 |
| 25 | 10 | 15.8 | 11.4 | 0 | 0 |
| | 1 | 412.4 | 208.4 | 315.7 | 308.2 |
| | 3 | 213.4 | 189.5 | 178.4 | 168.9 |
| | 5 | 96.4 | 89.4 | 48.1 | 46.2 |
| | 7 | 0 | 0 | 0 | 0 |
| 50 | 10 | 0 | 0 | 0 | 0 |
| | 1 | 301.4 | 289.4 | 298.4 | 279.6 |
| | 3 | 68.4 | 54.1 | 115.8 | 107.9 |
| | 5 | 81.4 | 79.2 | 0 | 0 |
| | 7 | 0 | 0 | 0 | 0 |
| 100 | 10 | 0 | 0 | 0 | 0 |
| | 1 | 298.7 | 278.7 | 297.4 | 289.4 |
| | 3 | 56.2 | 53.4 | 74.8 | 70.5 |
| | 5 | 0 | 0 | 0 | 0 |
| | 7 | 0 | 0 | 0 | 0 |
| 200 | 10 | 0 | 0 | 0 | 0 |

The data correspond to the average of three independent replicas.

method used to mix them, since agitation facilitates the entry of AEW into the test cells, and the well-mixed AEW resulting from the agitation allows the chlorine to react with the cells more efficiently (Park et al., 2002).

Strawberry fruits are extremely sensitive to injury both before and after harvest due to various environmental factors. The dangers of insects and fungi have led to the use of various pesticides throughout the fruit's growth cycle, however, increased concern among consumers about the presence of pesticides in food is driving the development of novel control measures to avoid deterioration of the fruit at harvest and during storage and packaging. The results obtained in this research suggest that, in addition to storage temperature management of the post-harvest strawberry, a pre-wash with electrolyzed water would be useful to avoid deterioration of the fruit. This technique would be especially advantageous given that electrolyzed water has been reported to have no negative effects on the organoleptic properties of the fruit (Al-Haq, 2005) and has also been reported as being safe for consumers' health. This last factor has become especially important, as the presence of pesticide residues in food continue to drive a search for non-chemical disease control measures.

Conclusion

The results obtained in this work demonstrate that AEW can be employed as a useful fungicide for strawberry molds (*B. cinerea* and *R. stolonifer*), and that it holds potential for the development of environmentally friendly products for the control of fungi in fruits in the storage phase. The treatment is an economical sanitization alternative against the phytopathogens that cause the majority of rot in the post-harvest period. Specifically, AEW (pH 6.6) could be used as a pre-wash treatment to reduce or eliminate the prevalence of fungal spores, and thereby help prolong the shelf-life of the fruit.

Acknowledgement

This research was supported by Universidad de Santander- Bucaramanga, Colombia. The authors wish to thank the firm of LimpiooTec for their technical support and donation of the electrolysis equipment used in this research.

References

- Ainsworth, G.C.; Sparrow, F.K.; Sussman, A.S. 1973. The Fungi: an advanced treatise. Academic Press, New York 4A:1-621.
- Al-Haq, M.I.; Sugiyama, J.; Isobe, S. 2005. Applications of Electrolyzed Water in Agriculture and Food Industries. Food Sci. Technol. Res. 11 (2):135-150. <https://doi.org/10.3136/fstr.11.135>
- Barnes, I.; Roux, J.; Wingfield, M.J.; Coetzee, MPA. 2001. Characterization of *Seiridium* spp. associated with cypress canker based on β -tubulin and histone sequences. Plant Dis.85:317-321. <https://apsjournals.apsnet.org/doi/10.1094/PDIS.2001.85.3.317>
- Bialka, K.L.; Demirci, A.; Knabel, S.J.; Patterson, P.H.; Puri, V.M. 2004 Efficacy of electrolyzed oxidizing water for the microbial safety and quality of eggs. J Poultr Sci. 83(12): 2071-2078. <https://doi.org/10.1093/ps/83.12.2071>
- Cao, W.; Zhu, Z.W.; Shi, Z.X.; Wang, C.Y.; Li, B.M. 2009. Efficiency of slightly acidic electrolyzed water for inactivation of *Salmonella* Enteritidis and its contaminated shell eggs. Int J Food Microbiol. 130(2):88-93. <https://doi.org/10.1016/j.ijfoodmicro.2008.12.021>
- Cary, J.W.; Beltz, S.B.; Bennett, CA. 2003 Identification of Common *Aspergillus* Species. Utrecht, Netherlands: Centraalbureau voor Schimmelcultures. 116 pp.
- Coley-Smith, J.R. 1980 Sclerotia and other structures in survival. In: Coley-Smith JR, Verhoeff K and Jarvis WR (eds) The Biology of *Botrytis*. 85-114. Academic Press, London, UK
- Callejas, T.A.; Martinez, G.B.; Artes, F.; Artes, H. F. 2010. Neutral and acid electrolyzed water as emergent sanitizer for fresh-cut mizuna baby levels. Postharvest Biol Technol. 59(3):298-306. <https://doi.org/10.1016/j.postharvbio.2010.09.013>
- Cui, X., Shang, Y.; Shi, Z.; Xin, H.; Cao, W. 2009. Physicochemical properties and bactericidal efficiency of neutral and acidic electrolyzed water under different storage conditions. J. Food Eng. 91, 582-586. <https://doi.org/10.1016/j.jfoodeng.2008.10.006>
- Cravero, F., Englezos, V.; Torchio, F.; Giacosa, S.; Segade, S. R.; Rantsiou, V. G. K.; Rolle, L. and Cocolin, L. 2016. Post-harvest control of wine-grape mycobiota using electrolyzed water. Inn. Food Sci. Emer. Technol. 35:21-28. <https://doi.org/10.1016/j.ifset.2016.03.010>
- Dean, R.; Van Kan, J.A.; Pretorius, Z.A.; Hammond-Kosack K.E.; Di Pietro, A.; Spanu, P.D.; Rudd, J.J.; Dickman, M.; Kahmann, R.; Ellis, J and Foster GD 2012. The Top 10 fungal pathogens in molecular plant pathology. Mol. Plant Pathol. 13(4):414-430. <https://doi.org/10.1111/j.1364-3703.2011.00783.x>
- Domsch, K.H.; Gams, W and Anderson, T.H. 1980. Compendium of Soil Fungi, 1-2: 1308. Acad Press, New York.
- Feliziani, E.; Lichter, A.; Smilanick, J. L. and Ippolito, A. 2016. Disinfecting agents for controlling fruit and vegetable diseases after harvest. Postharvest Biol Technol. 122:53-69. <https://doi.org/10.1016/j.postharvbio.2016.04.016>
- Fernández, A.F.J.; Carbú, M.; Garrido, C.; Vallejo, I.; Cantoral J.M. 2007 Proteomic advances in

- phytopathogenic fungi. *Curr Proteomics*. 4:79-88. <https://doi.org/10.3390/ijms12010795>
- Guentzel, J. L.; Callan, M. A.; Lam, K. L.; Emmons, S. A.; and Dunham, V. L. 2011. Evaluation of electrolyzed oxidizing water for phytotoxic effects and pre-harvest management of gray mold disease on strawberry plants. *Crop Prot.* 30(10):1274-1279. <https://doi.org/10.1016/j.cropro.2011.05.021>.
- Hao, J., Qui, S.; Li, H.; T, Cheng, Liu.H.; Li, L. 2012. Roles of hydroxyl radical in electrolyzed oxidizing water (EOW) for the inactivation of *Escherichia coli*. *Int. J. Food Microbiol.* 155(3):99-104. <https://doi.org/10.1016/j.ijfoodmicro.2011.12.031>
- Holz, G.; Coertze, S.; Williamson, B. 2007. *The ecology of Botrytis on plant surfaces*. Eds (Elad Y, Williamson B, Tudzynski P, Delen N) In: *Botrytis: Biology, Pathology and Control*. Heidelberg: Springer.9-27. https://doi.org/10.1007/978-1-4020-2626-3_2
- Huang, Y.R.; Hung, Y.C.; Hsu, S.Y.; Huang, Y.W.; and Hwang DF (2008). Application of electrolyzed water in the food industry. *Food Control.* 19(4):329-345. <https://doi.org/10.1016/j.foodcont.2007.08.012>
- Korsten, L.; Wehner, F.C. 2003. *Fungi*, In: Bartz, J.A., Brecht, J.K. (Eds.), *Postharvest Physiology and Pathology of Vegetables*. Second edition Marcel Dekker, Inc., New York, 485-518.
- Koseki, S.; Yoshida, K.; Isobe, S.; & Itoh, K. 2004. Efficacy of acidic electrolyzed water for microbial decontamination of cucumbers and strawberries. *J. Food Prot.* 67(6):1247-1251. <https://doi.org/10.4315/0362-028X-67.6.1247>.
- Lopes, U.P. 2011. Podridões em pós-colheita de morango: Etiologia e efeito de produtos alternativos. MSc Dissertation, Universidade Federal de Viçosa, Viçosa, MG. <http://locus.ufv.br/bitstream/handle/123456789/4399/texto%20completo.pdf?sequence=1&i>.
- Nelson, P.E.; Toussoun, T.A.; and Marasas, W.F.O. 1983. *Fusarium Species: An Illustrated Manual for Identification*. The Penn. St. Univ.Press, Univ. Park, Pennsylvania. 203.
- Newmeyer, D. 1990 Filtering small quantities of conidial suspensions to remove mycelial fragments, *Fungal Genetics Reports: Vol. 37, Article 15*. <https://doi.org/10.4148/1941-4765.1480>
- O'Donnell, K.L. 1979. *Zygomycetos in culture*. Univ. Georgia Press, Georgia EE. UU., 257
- Park, C.M.; Hung, Y.C. and Kim, C. 2002. Efectiveness of electrolyzed water as a sanitizer for treating diferent surfaces. *J. Food Prot.* 65:1276-1280 <https://doi.org/10.4315/0362-028X-65.8.1276>
- Rigotti S., K. Gindro.; H. Richter.; Viret, O. 2002. Characterization of molecular markers for specific and sensitive detection of *Botrytis cinerea* Pers.: Fr. in strawberry (*Fragaria x ananassa* Duch.) using PCR. *FEMS Microbiol Lett.* 209:169-174. [https://doi.org/10.1016/S0378-1097\(02\)00491-3](https://doi.org/10.1016/S0378-1097(02)00491-3)
- Seifert, K.A.; Gams, W. 2011 The genera of hyphomycetes. *Persoonia.* 27:119-129. <https://doi.org/10.3767/003158511X617435>
- White T.J.; Bruns, T.; Lee, S.; Taylor, J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: *PCR Protocols: a guide to methods and applications*. (Innis MA, Gelfand DH, Sninsky JJ, White TJ, eds). Academic Press, San Diego, U.S.A.: 315-322
- Y.J. Zhou.; J. Zhang.; X.D. Wang.; L. Yang.; D.H. Jiang.; G.Q. Li.; T. Hsiangand W.Y. Zhuang. 2014. Morphological and phylogenetic identification of *Botrytis sinoviticola*, a novel cryptic species causing gray mold disease of table grapes (*Vitis vinifera*) in China. *Mycologia* 106:1.43-56. <https://doi.org/10.3852/13-032>
- Zacharia, I.A.; Kamitani, Y.; Morita, K.; Iwasaki, K. 2010. Sanitization potency of slightly acidic electrolyzed water against pure cultures of *Escherichia coli* and *Staphylococcus aureus*, in comparison with that of other food sanitizers. *Food Control* 21(5):740-745. <https://doi.org/10.1016/j.foodcont.2009.11.002>