

# Effect of UV-C irradiation combined with a sodium alginate edible coating on postharvest quality and antioxidant potential of sour guava (*Psidium friedrichsthalianum* Nied.)

Efecto de la combinación de irradiación UV-C y un recubrimiento comestible de alginato de sodio en la calidad poscosecha y el potencial antioxidante de la guayaba agria (*Psidium friedrichsthalianum* Nied.)

Alba Mery Garzón García <sup>1,2</sup>.

<sup>1</sup>Universidad del Valle - Sede Regional Caicedonia. Caicedonia, Valle del Cauca, Colombia. <sup>2</sup>  [garzon.alba@correounivalle.edu.co](mailto:garzon.alba@correounivalle.edu.co)



DOI: <https://doi.org/10.15446/acag.v74n1.123919>

2025 | 74-1 p 55-62 | ISSN 0120-2812 | e-ISSN 2323-0118 | Rec.: 2025-11-16 Acep.: 2026-03-06

## Abstract

Sour guava is a promising fruit due to its nutritional value and phytochemical richness in the fresh state. However, reports on the postharvest behavior of this fruit are very limited. Therefore, the aim of this research was to evaluate the effect of emerging technologies on the quality attributes and antioxidant potential of sour guava. A completely randomized design with two factors (treatment type and storage time) was employed. Fruits were divided into three lots according to the treatment applied: control (C), UV-C irradiation for 5 minutes (UV), and a combined treatment consisting of UV-C irradiation for 5 minutes followed by the application of a 2 % sodium alginate edible coating (UV+EC). After treatment, the fruits were stored for 21 days at 8 °C and 70 % relative humidity. Physicochemical parameters (pH, total soluble solids, titratable acidity, firmness, and weight loss), color attributes ( $L^*$ ,  $a^*$ , and  $b^*$ ), and antioxidant activity (total phenolic content and antioxidant capacity determined by DPPH, ABTS, and FRAP assays) were evaluated on days 0, 7, 14, and 21. A two-way ANOVA followed by a Tukey's post hoc test was conducted to assess the effects of treatment type and storage time on physicochemical parameters, color attributes, and antioxidant activity ( $p < 0.05$ ). Results showed that UV+EC treatment effectively delayed ripening and helped preserve both surface color and antioxidant capacity during storage, whereas the UV treatment alone only increased antioxidant activity up to approximately day 14 of storage. Further research is required to evaluate sensory acceptance and to conduct a complete metabolomic characterization.

**Keywords:** Antioxidant capacity, color analysis, fruit quality, phenols, physicochemical parameters.

## Resumen

La guayaba agria es una fruta prometedora debido a su valor nutricional y riqueza fitoquímica en estado fresco. Sin embargo, los estudios sobre el comportamiento poscosecha de esta fruta son muy limitados. Por tanto, el objetivo de esta investigación fue evaluar el efecto de tecnologías emergentes sobre los atributos de calidad y el potencial antioxidante de la guayaba agria. Se empleó un diseño completamente al azar con dos factores (tipo de tratamiento y tiempo de almacenamiento). Las frutas se dividieron en tres lotes según el tratamiento aplicado: control (C), irradiación UV-C durante 5 minutos (UV) y un tratamiento combinado consistente en irradiación UV-C durante 5 minutos seguida de la aplicación de un recubrimiento comestible de alginato de sodio al 2 % (UV+EC). Tras los tratamientos, las frutas se almacenaron durante 21 días a 8 °C y con humedad relativa de 70 %. Se evaluaron parámetros fisicoquímicos (pH, sólidos solubles totales, acidez titulable, firmeza y pérdida de peso), atributos de color ( $L^*$ ,  $a^*$  y  $b^*$ ) y actividad antioxidante (contenido de fenoles totales y capacidad antioxidante mediante los ensayos DPPH, ABTS y FRAP) en los días 0, 7, 14 y 21. Se realizó un ANOVA de dos vías seguido de una prueba post hoc de Tukey para evaluar el efecto del tipo de tratamiento y el tiempo de almacenamiento sobre los parámetros fisicoquímicos, los atributos de color y la actividad antioxidante ( $p < 0.05$ ). Los resultados mostraron que el tratamiento UV+EC retardó eficazmente la maduración y ayudó a preservar tanto el color superficial como la capacidad antioxidante durante el almacenamiento, mientras que el tratamiento UV por sí solo incrementó la actividad antioxidante hasta aproximadamente el día 14 de almacenamiento. Se requiere investigación adicional para evaluar la aceptación sensorial y realizar una caracterización metabolómica completa.

**Palabras clave:** análisis de color, calidad de la fruta, capacidad antioxidante, fenoles, parámetros fisicoquímicos.

## Introduction

Sour guava or Costa Rican guava (*Psidium friedrichsthalianum* Nied.) is a fruit produced by a tree of the Myrtaceae family, distributed from southern Mexico to eastern Venezuela. In Colombia, this species grows naturally in the Cauca and Magdalena valleys, from sea level up to 800 m of altitude, as it is sensitive to low temperatures. Generally, the fruit is used to prepare beverages, jellies, and jams (Lim, 2012). The fruit is globose, measuring between 3 cm and 6 cm in diameter, with a fleshy and acidic pulp. During ripening, its exocarp changes color from green to yellowish hues (Muñoz-Arrieta *et al.*, 2021). Recent studies have highlighted the phytochemical and pharmacological potential of this species, mainly attributing it to its high antioxidant and antimicrobial activities (Santos *et al.*, 2023), as well as to the presence of flavonoids and ellagic acid derivatives as the main bioactive compounds (Cuadrado-Silva *et al.*, 2017).

According to Singh (2011), during guava ripening, the chlorophyll content generally decreases while carotenoid levels increase. Furthermore, fruit softening is accompanied by an increase in soluble pectin and a decrease in certain carbohydrates such as cellulose, hemicellulose, lignin, and starch. Therefore, several postharvest handling factors are crucial to ensure fruit quality, including temperature management ( $> 8^{\circ}\text{C} - 10^{\circ}\text{C}$ ), prevention of physical damage, control of water loss, and reduction of dry rot incidence. Although some studies have reported the effect of the application of 1-methylcyclopropene (1-MCP) on cv. 'Allahabad Safeda' (Singh and Pal, 2008), melatonin on cv. 'Zhenzhu' (Fan *et al.*, 2022), and edible coatings on cv. 'Lalit' and 'pear' guava (Vishwasrao and Ananthanarayan, 2016; Hong *et al.*, 2012), only one investigation has addressed the extension of sour guava shelf life using silver nitrate, mercuric chloride, and sodium hypochlorite at different doses and exposure times (Villalobos *et al.*, 1999).

Edible coatings are widely used and well accepted in fruit preservation as a barrier to gases, water vapor, and external contamination (Aljabary *et al.*, 2025). Nevertheless, the combined application of this technology with superficial and non-ionizing radiation, such as ultraviolet shortwave radiation (UV-C), remains very limited, particularly in species belonging to the Myrtaceae family. Previous studies suggest that the combined use of postharvest technologies may exert a positive effect on shelf life and bioactivity of sour guava. Therefore, the aim of this study was to evaluate the effect of combining UV-C and a sodium alginate edible coating on the postharvest quality and antioxidant potential of sour guava.

## Materials and methods

### Raw material

Approximately 1.7 kg of sour guava (*Psidium friedrichsthalianum* Nied.) was obtained from Génova, Quindío, Colombia. Guavas were transported to the RoastLab at Universidad del Valle, Caicedonia Regional Campus, where they were rinsed, washed, and air-dried under ambient conditions (Figure 1).

### Chemical reagents

Sodium acetate buffer, gallic acid, and Folin-Ciocalteu reagent used in this study were purchased from Alpha Chemika (Mumbai, MH, India). 2,4,6-tris(2-pyridyl)-s-triazine (TPTZ), 2,2-diphenyl-1-picrylhydrazyl (DPPH), and 2,2'-azino-bis-(3-ethylbenzothiazolinone-6-sulfonic acid (ABTS) radicals were obtained from Sigma-Aldrich (St. Louis, MO, USA). Hydrochloric acid was supplied by Panreac AppliChem (Castellar del Vallès, Barcelona, Spain). Ferric chloride ( $\text{FeCl}_3$ ) and potassium peroxodisulfate were purchased from Millipore Corporation (Darmstadt, Hesse, Germany). Absolute ethanol (95 %) was locally procured from San Jorge (Cali, Valle del Cauca, Colombia).

### Fruit treatments and storage

The fruits were divided into three lots according to the treatment applied: 1) control (C), 2) UV, and 3) UV+EC. The first lot consisted of guavas without treatment. The second lot was irradiated for 5 min with UV-C. For this purpose, sets of eight fruits were placed under a 20 W UV lamp (UVC20W, ANTIVID19, Ecolite S.A.S., Cali, VAC, CO) positioned 10 cm above the guavas (Acevedo-Daza *et al.*, 2024). The lamp intensity was  $58.549 \text{ W/m}^2$ , and the applied dose was  $2.196 \text{ kJ/m}^2$ , calculated according to the extensive source with spherical emission (ESSE) model (Gardner and Shama, 1999).

UV-C irradiation was performed away from laboratory personnel to prevent skin and eye exposure. The last lot was subjected to UV-C irradiation followed by the application of a sodium alginate edible coating (EC). The coating formulation contained 2 % (w/w) sodium alginate, 1.5 % (w/w) glycerol, 0.1 % (w/w) polysorbate 80, and 96.4 % (w/w) distilled water. It was applied with a soft brush to form a layer approximately 0.2 mm thick and then dried by forced convection. Finally, all fruits were stored under refrigeration ( $8^{\circ}\text{C}$  and 70 % relative humidity, RH) for 21 days.

### Physicochemical analysis

pH was measured using a potentiometer (HI 991001, HANNA Instruments, Woonsocket, RI, USA). Total soluble solids (TSS) were determined using a digital refractometer (HI96800, HANNA Instruments), while



**Figure 1.** Sour guava (*Psidium friedrichsthalianum* Nied.) employed during the study.

firmness was assessed using a digital penetrometer equipped with a 7.9-mm diameter probe (GY-M30, Shenzhen, GD, CN). Titratable acidity (TA) was determined following the AOAC 942.15 method, considering citric acid as the predominant organic acid (AOAC, 2022). Weight loss (WL%) was calculated as the percentage difference between the initial weight and the weight recorded during storage (Peralta-Ruiz *et al.*, 2020).

### Color evaluation

Color parameters in the CIE  $L^*a^*b$  system were obtained using a 3NS800 spectrophotometer (3NH, Zengcheng District, GZ, CN), calibrated with a D65 illuminant and a  $10^\circ$  standard observer.

### Antioxidant activity

#### Preparation of extracts

Sample extracts were prepared using 1 g of sour guava and 10 mL of absolute ethanol. The mixtures were subjected to an ultrasonic bath for 15 min and subsequently centrifuged for 15 min at 3600 rpm (Z 206 A; Hermle Benchmark, Franklin, WI, USA) (Garzón-García *et al.*, 2023). The extracts were diluted by a factor of 50 before being used in the subsequent tests.

#### Total phenolic content

Total phenolic content was determined according to Pérez-Perez *et al.* (2020). 200  $\mu$ L of extract and 500  $\mu$ L of 1 N Folin-Ciocalteu reagent were mixed and kept under refrigeration for 5 min. Subsequently, 500  $\mu$ L of 20 %  $\text{NaCO}_3$  and 2.8 mL of distilled water were added. The mixture was allowed to stand for 30 min, after which the absorbance was measured at 760 nm using an UV-Vis spectrophotometer (UV-1200; MAPADA

INSTRUMENTS, Songjiang District, SH, CN). A calibration curve was constructed using gallic acid as the standard, with concentrations ranging from 0 g/mL to 0.4 g/mL ( $R^2 = 0.990$ ). Results were expressed as milligrams of gallic acid equivalents per 100 g of fresh sample.

#### Antioxidant capacity by ABTS assay

The antioxidant capacity by the ABTS assay was determined according to González-Vega *et al.* (2021). Initially, a radical solution was prepared by dissolving 19.3 mg of ABTS in 5 mL of distilled water. Separately, 0.0378 g of potassium persulfate was dissolved in 1 mL of distilled water. Subsequently, 88  $\mu$ L of the persulfate solution was added to the initial solution. This resulting mixture was kept in the dark for 12 h, after which it was adjusted to an absorbance of  $0.7 \pm 0.01$  at a wavelength of 734 nm. For the measurements, 250  $\mu$ L of extract was mixed with 3375  $\mu$ L of the prepared radical cation solution, and the absorbance was measured at 734 nm after 30 min. Results were expressed as radical inhibition capacity (%) according to Equation 1, using distilled water as the control.

$$\% \text{Inhibition} = \frac{\text{Control absorbance} - \text{Extract absorbance}}{\text{Control absorbance}} \times 100 \quad (\text{Eq. 1})$$

#### Antioxidant capacity by DPPH assay

The antioxidant capacity by the DPPH assay was evaluated based on the methodology proposed by Pérez-Perez *et al.* (2020) with some modifications. A total of 1.5 mg of DPPH radical was dissolved in 50 mL of methanol. This solution was adjusted to an absorbance of  $0.7 \pm 0.01$  at a wavelength of 515 nm. Subsequently, 300  $\mu$ L of extract and 3 mL of the radical solution were mixed. The mixture was

allowed to stand in the dark for 30 min, after which the absorbance was measured at a wavelength of 515 nm. Results were expressed as radical inhibition capacity (%) according to Equation 1, using distilled water as the control.

### Antioxidant capacity by ferric reducing antioxidant power (FRAP)

Antioxidant capacity by the FRAP assay was determined following the methodology proposed by Garzón-García *et al.* (2023) with some modifications. Three stock solutions were prepared: a 300 mM sodium acetate buffer (pH 3.6), a 20 mM  $\text{FeCl}_3 \times 6\text{H}_2\text{O}$  solution, and a 10 mM TPTZ solution in 40 mM HCl. The working solution was prepared by mixing the solutions in a 10:1:1 (v/v/v) ratio (buffer,  $\text{FeCl}_3 \times 6\text{H}_2\text{O}$ , and TPTZ-HCl, respectively). 3.5 mL of the working solution was added to 250  $\mu\text{L}$  of extract, and the mixture was allowed to stand for 30 min. The absorbance of this mixture was measured at a wavelength of 638 nm. A calibration curve was constructed using gallic acid as the standard, with concentrations ranging from 0 mg/mL to 0.1 mg/mL ( $R^2 = 0.995$ ). The results were expressed as milligrams of gallic acid equivalents per 100 g of fresh sample.

### Statistical analysis

Results were presented as the mean of triplicate measurements  $\pm$  standard error. A completely randomized design with two factors was employed. The first factor was treatment type, with three levels (C, UV, and UV+EC), while the second factor was storage time, with four levels (0, 7, 14, and 21 days). A two-way ANOVA followed by Tukey's post hoc test was conducted to assess the effects of treatment type and storage time on the physicochemical parameters, color attributes, and antioxidant activity ( $p < 0.05$ ). All statistical analyses were performed in Minitab 17 (Minitab Inc, State College, PA, US).

## Results

### Physicochemical parameters


Treatment type, storage time, and their interaction had a significant effect on pH and TA ( $p < 0.05$ ). The pH values ranged from 2.763 to 2.947, 2.727 to 2.693, and 2.813 to 2.827 for C, UV, and UV+EC treatments, respectively. During storage, TA values decreased in control samples and increased for UV and UV+EC (C: 2.240 % - 1.173 %, UV: 1.920 % - 2.880 %, UV+EC: 2.027 % - 2.240 %).

Total soluble solids (TSS) were significantly affected by storage time and by the interaction between storage time and treatment type ( $p < 0.05$ ). TSS values increased rapidly in fruits from C and UV treatments (11.5 - 17.1 °Brix and 11.1 - 17.9 °Brix, respectively), whereas the combined UV+EC treatment delayed this increase (10.3 - 11.5 °Brix). These results indicate that the combined treatment effectively slowed metabolic activity during storage.

All factors and their interaction had a significant effect on firmness and weight loss ( $p < 0.05$ ). As shown in Figure 2, sour guavas from the UV treatment exhibited the highest firmness values at the end of storage, along with weight loss levels similar to those observed in the control fruits. Thus, weight loss likely contributed to firmer fruits during texture assessment. Conversely, fruits subjected to the combined UV+EC treatment showed lower firmness and weight loss values than those observed in the C and UV treatments.

### Color parameters

The coordinates  $L^*$ ,  $a^*$ , and  $b^*$  (where  $L^*$  represents luminosity,  $-L^*$  = black and  $+L^*$  = white;  $a^*$  indicates color from  $-a^*$  = green to  $+a^*$  = red; and  $b^*$  from  $-b^*$  = blue to  $+b^*$  = yellow) were assessed in sour guavas during 21 days of storage (Figure 3). According to the




**Figure 2.** Firmness (a) and weight loss (b) of sour guava subjected UV, and UV+EC treatments, as well as the control (C), during storage at 8 °C and 70 % relative humidity for 21 days.

Note. Bars represent the mean values, and the vertical lines above the bars correspond to the standard error. Different lowercase letters above the bars indicate significant differences among treatments during storage, according to Tukey's post hoc test ( $p < 0.05$ ).



statistical analysis, treatment type, storage time, and their interaction significantly affected the CIE  $L^*a^*b^*$  coordinates ( $p < 0.05$ ). Luminosity ( $L^*$ ) of fruits from C and UV treatments slightly decreased during storage, whereas  $L^*$  values in fruits from the combined UV+EC treatment remained nearly constant. Additionally,  $a^*$  values progressively increased over time, while  $b^*$  values remained relatively stable in fruits from the control and UV+EC treatments. These results indicate that postharvest treatments significantly contributed to preserving the visual appearance of the fruits during storage. In contrast, fruits from the control treatment exhibited darker and more yellowish tones at the end of storage, indicating accelerated ripening.



**Figure 3.** Color coordinates  $L^*$  (a),  $a^*$  (b), and  $b^*$  (c) of sour guava subjected to UV, and UV+EC treatments, as well as the control (C), during storage at 8 °C and 70 % relative humidity for 21 days.

Note. Bars represent the mean values, and the vertical lines above the bars correspond to the standard error. Different lowercase letters above the bars indicate significant differences among treatments during storage, according to Tukey's post hoc test ( $p < 0.05$ ).

## Antioxidant activity


Antioxidant activity was evaluated using different antioxidant capacity assays and by determining total phenolic content. For both the ABTS and DPPH assays, all factors and their interactions significantly influenced antioxidant capacity ( $p < 0.05$ ). As shown in Figure 4, the inhibition of ABTS radical cation tended to decrease at the end of the storage; however, the highest values were recorded in samples from the UV and UV+EC treatments. For DPPH radical inhibition, fruits exposed to the UV treatment showed an increase and maintained the highest values throughout storage.

All factors and their interaction had a significant effect on total phenolic content and antioxidant capacity determined by the FRAP assay ( $p < 0.05$ ). As shown in Figure 5, total phenolic content significantly decreased after day 14 of storage, whereas fruits subjected to the UV treatment exhibited the highest values, followed by those treated with the UV+EC treatment. The antioxidant capacity determined by the FRAP assay showed a trend similar to that observed for total phenolic content. A significant reduction in antioxidant capacity was detected after day 14 of storage, with the highest values recorded in UV-treated samples, followed by those subjected to the combined UV+EC treatment.

## Discussion


Physicochemical parameters provided a partial assessment of the metabolic response to treatments. In some cases, particularly in control samples, the observed decrease in pH may be attributed to the formation of free acids, ascorbic acid degradation, and pectin hydrolysis (Wagh *et al.*, 2022). Labib *et al.* (2025) reported that the application of edible coatings based on chitosan, cinnamon essential oil, aloe vera gel, propolis, and gum Arabic, applied individually or in combination on 'Thai 5' guava fruits, resulted in a slower conversion of organic acids into sugars. This effect delayed the increase in TSS and pH, and the reduction of TA during 12 days of storage.

The rapid increase in TSS after radiation treatments has already been widely reported. Yousaf *et al.* (2023) found that low doses of  $\gamma$ -irradiation ( $< 1.5$  kGy) caused a rapid rise in TSS in 'Gola' guava during 20 days of storage, as the treatment did not slow down ripening and starch was hydrolyzed into sugars and organic acids. In another study, the application of edible coatings based on 2 % sodium alginate and 0.2 % – 0.8 % *Cyclea barbata* leaves extract on 'Getas Merah' guavas maintained stable TSS levels until day 16 of a 20-day storage period. By the end of storage, TSS values had decreased to approximately 6 °Brix (Utama *et al.*, 2022).



**Figure 4.** ABTS (a) and DPPH (b) radical inhibition percentage in sour guava subjected to UV, and UV+EC treatments, as well as the control (C), during storage at 8 °C and 70 % relative humidity for 21 days.

Note. Bars represent the mean values, and the vertical lines above the bars correspond to the standard error. Different lowercase letters above the bars indicate significant differences among treatments during storage, according to Tukey's post hoc test ( $p < 0.05$ ).



**Figure 5.** Total phenolic content (a) and antioxidant capacity determined by the FRAP assay (b) in sour guava subjected to UV, and UV+EC treatments, as well as the control (C) during storage at 8 °C and 70 % relative humidity for 21 days.

Note. Bars represent the mean values, and the vertical lines above the bars correspond to the standard error. Different lowercase letters above the bars indicate significant differences among treatments during storage, according to Tukey's post hoc test ( $p < 0.05$ ).

The reduced firmness loss and weight loss at the end of storage observed in the UV+EC treatment can be attributed to the barrier generated by the edible coating. Hasan *et al.* (2022) reported that 2 % sodium alginate-based edible coatings reduced weight and firmness loss in guavas during 16 days of storage at 11 °C and 85 % RH, as alginate coatings effectively limit the activity of polyphenol oxidase, peroxidase, cellulase, polygalacturonase, and pectin methylesterase.

Regarding color parameters, fruits from the UV+EC treatment exhibited a more uniform and characteristic coloration. In contrast, control fruits showed a marked increase in  $b^*$  values along with reduced luminosity, with a more intense yellowish appearance indicating accelerated deterioration. This behavior is consistent with findings reported by Hasan *et al.* (2022), who observed a similar evolution of color in uncoated mature guavas stored for 16 days at 11 °C.

The inhibition of both the DPPH radical and ABTS radical cation was higher in samples from the UV treatment, followed by those from the UV+EC treatment. This effect could be attributed to UV-C radiation stimulating the shikimic acid pathway and phenylalanine ammonia-lyase activity, thereby enhancing the synthesis of hydroxycinnamic acids (Surjadinata *et al.*, 2017). In addition, during storage, lipophilic phenolic compounds, including flavonoids, may persist and stabilize free radicals by donating electrons. On the other hand, Yousaf *et al.* (2023) reported that  $\gamma$ -irradiation at doses between 0.2 kGy and 1.0 kGy significantly reduced DPPH radical inhibition in 'Gola' guava samples stored for 20 days, likely due to increased polyphenol oxidase activity, which promotes the production of free radicals.

Total phenolic content and FRAP assay results were expressed using gallic acid as the standard, according to the report by Bibi Sadeer *et al.* (2020). Total phenolic content of sour guavas was lower

than the values reported for *Psidium araca* and *Psidium guajava* L. cv. 'Thai' (Zapata *et al.*, 2013; Hasan *et al.*, 2022). The UV treatment also generated higher values for both the FRAP assay and total phenolic content. A similar behavior was observed in 'Barafkhana' guavas exposed to different UV-C doses (0, 1.6, 2.0, and 2.4 kJ/m<sup>2</sup>), stored under refrigeration for 20 days, where control fruits exhibited a reduction in total phenolic content toward the end of the storage period (Menaka *et al.*, 2024). Furthermore, edible coatings may help limit oxygen uptake and reduce the accumulation of reactive oxygen species (ROS) (Labib *et al.*, 2025).

This trend aligns with the findings of Hasan *et al.* (2022), who observed a decrease in antioxidant capacity determined by the FRAP assay at the end of 16 days of storage at 11 °C in guavas (*Psidium guajava* L.) treated with different formulations of 2 % sodium alginate-based edible coatings. The behavior observed for the FRAP assay also resembles that of ascorbic acid content, as the antioxidant capacity measured by this assay may reflect the presence of hydrophilic bioactive compounds capable of stabilizing free radicals through electron donation, such as ascorbic acid (Bibi Sadeer *et al.*, 2020; Yousaf *et al.*, 2024).

## Conclusions

This study enabled the evaluation and comparison of the effects of emerging postharvest technologies on a promising exotic fruit. The combined treatment consisting of UV-C irradiation followed by the application of a sodium alginate edible coating effectively delayed the ripening and senescence of sour guava, as demonstrated by improvements in physicochemical and color attributes. Nevertheless, this treatment did not promote an increase in antioxidant activity but rather contributed to its preservation during storage.

Conversely, UV-C irradiation alone resulted in an increase in antioxidant activity; however, it did not exert the same positive influence on overall fruit quality as the combined treatment. Further omics analyses are necessary to determine which bioactive compounds and metabolic pathways are involved in the response of sour guava to different postharvest treatments. Overall, this investigation corresponds to an initial step toward highlighting underutilized species with nutraceutical potential, as well as postharvest strategies that could be scalable and sustainable for the food industry in the region.

## References

- Acevedo-Daza, J.; Villegas-Ciro, J. A.; Pineda-Sepulveda, V.; Jaramillo González, J. C.; Restrepo-Cubillos, V.; Peña-García, J. T.; Soto-Bastidas, J. H.; Buitrago-Pinilla, L. T.; Arroyo-Merino, J. A.; Ñañez-Palacio, D. C.; Castro-Vargas, D. F.; Largo-Ávila, E. and Garzón-García, A. M. (2024). UV-C effect on postharvest quality of citrus fruits from the northeastern region of Valle del Cauca, Colombia. *Biotechnia*, 26, e2303. <https://doi.org/10.18633/biotechnia.v26.2303>
- Aljabary, A. M. A. O.; Awlqadr, F. H.; Altemimi, A. B.; Abdulrahman, A. B. M.; Saeed, M. N. and Hesarinejad, M. A. (2025). Advances in oil-based edible coatings for postharvest preservation of fruits and vegetables: A comprehensive review of biopolymer types, functional plant oils, nanoemulsion systems, and application techniques. *Journal of Agriculture and Food Research*, 24, 102425. <https://doi.org/10.1016/j.jafr.2025.102425>
- AOAC International. (2022). AOAC official method 942.15 - Acidity (titratable) of fruit products. In Latimer, G. W. (Ed.), *Official methods of analysis of AOAC International* (22nd Ed.). Oxford Academic.
- Bibi Sadeer, N.; Montesano, D.; Albrizio, S.; Zengin, G. and Mahomoodally, M. F. (2020). The versatility of antioxidant assays in food science and safety - Chemistry, applications, strengths, and limitations. *Antioxidants*, 9(8), 709. <https://doi.org/10.3390/antiox9080709>
- Cuadrado-Silva, C. T.; Pozo-Bayón, M. Á. and Osorio, C. (2017). Targeted metabolomic analysis of polyphenols with antioxidant activity in sour guava (*Psidium friedrichsthalianum* Nied.) fruit. *Molecules*, 22(1), 11. <https://doi.org/10.3390/molecules22010011>
- Fan, S.; Xiong, T.; Lei, Q.; Tan, Q.; Cai, J.; Song, Z.; Yang, M.; Chen, W.; Li, X. and Zhu, X. (2022). Melatonin treatment improves postharvest preservation and resistance of guava fruit (*Psidium guajava* L.). *Foods*, 11(3), 262. <https://doi.org/10.3390/foods11030262>
- Gardner, D. W. M. and Shama, G. (1999). UV intensity measurement and modelling and disinfection performance prediction for irradiation of solid surfaces with UV light. *Food and Bioprocess Processing*, 77(3), 232-242. <https://doi.org/10.1205/096030899532510>
- Garzón-García, A. M.; Ruiz-Cruz, S.; Dussán-Sarria, S.; Hleap-Zapata, J. I.; Márquez-Ríos, E.; Del-Toro-Sánchez, C. L.; Tapia-Hernández, J. A.; Canizales-Rodríguez, D. F. and Ocaño-Higuera, V. M. (2023). Effect of UV-C postharvest disinfection on the quality of fresh-cut 'Tommy Atkins' Mango. *Polish Journal of Food and Nutrition Sciences*, 73(1), 39-49. <https://doi.org/10.31883/pjfn/159290>
- González-Vega, R. I.; Cárdenas-López, J. L.; López-Elías, J. A.; Ruiz-Cruz, S.; Reyes-Díaz, A.; Pérez-Pérez, L. M.; Cinco-Moroyoqui, F. J.; Robles-Zepeda, R. E.; Borboa-Flores, J. and Del-Toro-Sánchez, C. L. (2021). Optimization of growing conditions for pigments production from microalga *Navicula incerta* using response surface methodology and its antioxidant capacity. *Saudi Journal of Biological Sciences*, 28(2), 1401-1416. <https://doi.org/10.1016/j.sjbs.2020.11.076>
- Hasan, K.; Islam, R.; Hasan, M.; Sarker, S. H. and Biswas, M. H. (2022). Effect of alginate edible coatings enriched with black cumin extract for improving postharvest quality characteristics of guava (*Psidium guajava* L.) fruit. *Food and Bioprocess Technology*, 15, 2050-2064. <https://doi.org/10.1007/s11947-022-02869-2>
- Hong, K.; Xie, J.; Zhang, L.; Sun, D. and Gong, D. (2012). Effects of chitosan coating on postharvest life and quality of guava (*Psidium guajava* L.) fruit during cold storage. *Scientia Horticulturae*, 144, 172-178. <https://doi.org/10.1016/j.scientia.2012.07.002>
- Labib, L. A.; Ahmed, S. and Hasan, F. (2025). Improving guava shelf life and preserving postharvest quality with edible

- coatings. *Food Science & Nutrition*, 13(6), e70491. <https://doi.org/10.1002/fsn3.70491>
- Lim, T. K. (2012). *Psidium friedrichsthalianum*. In *Edible medicinal and non medicinal plants: Volume 3, Fruits* (pp. 681-683). Springer. [https://doi.org/10.1007/978-94-007-2534-8\\_94](https://doi.org/10.1007/978-94-007-2534-8_94)
- Menaka, M.; Asrey, R.; Vinod, B. R.; Ahamad, S.; Meena, N. K.; Bhan, C. and Goswami, A. K. (2024). UV-C irradiation enhances the quality and shelf-life of stored guava fruit via boosting the antioxidant systems and defense responses. *Food and Bioprocess Technology*, 17, 3704-3715. <https://doi.org/10.1007/s11947-024-03338-8>
- Muñoz-Arrieta, R.; Esquivel-Alvarado, D.; Alfaro-Viquez, E.; Rodríguez-Salazar, M.; Álvarez-Valverde, V.; Rodríguez, G. and Reed, J. D. (2021). Nutritional and chemical composition of the Costa Rican guava (*Psidium friedrichsthalianum* [O. Berg] Nied): An underexploited edible fruit with nutritional and industrial potential. *ACS Food Science & Technology*, 1(10), 1970-1978. <https://doi.org/10.1021/acsfoodscitech.1c00262>
- Peralta-Ruiz, Y.; Tovar, C. D. G.; Sinning-Mangonez, A.; Coronell, E. A.; Marino, M. F. and Chaves-Lopez, C. (2020). Reduction of postharvest quality loss and microbiological decay of tomato "Chonto" (*Solanum lycopersicum* L.) using chitosan-e essential oil-based edible coatings under low-temperature storage. *Polymers*, 12(8), 1822. <https://doi.org/10.3390/polym12081822>
- Pérez-Pérez, L. M.; Del Toro Sánchez, C. L.; Sánchez Chavez, E.; González Vega, R. I.; Reyes Díaz, A.; Borboa Flores, J.; Soto-Parra, J. M. and Flores-Cordova, M. A. (2020). Bioaccessibility of antioxidant compounds from different bean varieties (*Phaseolus vulgaris* L.) in Mexico, through an in vitro gastrointestinal system. *Biotechnia*, 22(1), 117-125. <https://doi.org/10.18633/biotechnia.v22i1.1159>
- Santos, P. V. L.; Da Cruz, E. D. N. S.; Nunes, J. D. A.; Mourão, R. H. V.; Do Nascimento, W. M. O.; Maia, J. G. S. and Figueiredo, P. L. B. (2023). Seasonal influence on volatile composition of *Psidium friedrichsthalianum* leaves, sampled in the Brazilian Amazon. *Horticulturae*, 9(7), 768. <https://doi.org/10.3390/horticulturae9070768>
- Singh, S. P. (2011). 10 - Guava (*Psidium guajava* L.). In Yahia, E. M. (Ed.), *Postharvest biology and technology of tropical and subtropical Fruits* (pp. 213-246e). Woodhead Publishing. <https://doi.org/10.1533/9780857092885.213>
- Singh, S. P. and Pal, R. K. (2008). Response of climacteric-type guava (*Psidium guajava* L.) to postharvest treatment with 1-MCP. *Postharvest Biology and Technology*, 47(3), 307-314. <https://doi.org/10.1016/j.postharvbio.2007.08.010>
- Surjadinata, B. B.; Jacobo-Velázquez, D. A. and Cisneros-Zevallos, L. (2017). UVA, UVB and UVC light enhances the biosynthesis of phenolic antioxidants in fresh-cut carrot through a synergistic effect with wounding. *Molecules*, 22(4), 668. <https://doi.org/10.3390/molecules22040668>
- Utama, N. A.; Pranata, I. A. and Pramesi, P. C. (2022). Maintaining physicochemical and sensory properties of guava var. Getas Merah using alginate and *Cyclea barbata* leaves powder as edible coating. *Advances in Horticultural Science*, 36(2), 135-144. <https://oaj.fupress.net/index.php/ahs/article/view/12341>
- Villalobos, M. R.; León de Sierralta, S. and Urdaneta Fernández, A. (1999). Evaluation of surface disinfectants on the in vitro establishment of *Psidium guajava* L. and *Psidium friedrichsthalianum* (Berg) Nierdz. *Revista de la Facultad de Agronomía (LUZ)*, 16, 243-255. [https://www.revfacagronluz.org.ve/v16\\_3/v163z003.html](https://www.revfacagronluz.org.ve/v16_3/v163z003.html)
- Vishwasrao, C. and Ananthanarayan, L. (2016). Postharvest shelf-life extension of pink guavas (*Psidium guajava* L.) using HPMC-based edible surface coatings. *Journal of Food Science and Technology*, 53, 1966-1974. <https://doi.org/10.1007/s13197-015-2164-x>
- Wagh, A. V.; Patil, S. R.; Sonkamble, A. M.; Gedam, A. P. and Deshmukh, R. N. (2022). Studies on storability of guava pulp. *The Pharma Innovation*, 11(4), 2144-2147. <https://www.thepharmajournal.com/archives/?year=2022&vol=11&issue=4&ArticleId=12307>
- Yousaf, A. A.; Abbasi, K. S.; Ibrahim, M. S.; Sohail, A.; Faiz, M. and Khadim, M. (2024). Storage stability assessment of guava fruit (*Psidium guajava* L.) cv. 'Gola' in response to different packaging materials. *Sustainable Food Technology*, 2(1), 210-221. <https://doi.org/10.1039/D3FB00113J>
- Yousaf, A. A.; Sarfraz, K.; Ahmed, A.; Hassan, I.; Ali, H. and Mehmood, T. (2023). Storage stability assessment of indigenous guava fruits (*Psidium guajava* L.) cv. "Gola" in response to  $\gamma$ -irradiation. *Journal of Food Processing and Preservation*, 2023, 4546469. <https://doi.org/10.1155/2023/4546469>
- Zapata, K.; Cortés, F. B. and Rojano, B. A. (2013). Polyphenols and antioxidant activity of sour guava fruit (*Psidium araca*). *Información Tecnológica*, 24(5), 103-112. <http://doi.org/10.4067/S0718-07642013000500012>