

First record of predation on *Rhinella diptycha* (Anura, Bufonidae) by *Caiman latirostris* (Crocodylia, Alligatoridae)

Primer registro de depredación de *Rhinella diptycha* (Anura, Bufonidae) por *Caiman latirostris* (Crocodylia, Alligatoridae)

Rodney Murillo Peixoto-Couto^{1*}, Maiara Cabrera-Miguel², Zaida Ortega³

Abstract

Caimans of the species *Caiman latirostris*, which are widely distributed in South America, are opportunistic predators. They show ontogenetic variation in diet, with young individuals initially feeding on invertebrates and then gradually adding vertebrates in their diets as they grow up. However, due to inexperience, young individuals may end up eating dangerous prey, such as poisonous amphibians. The intoxication caused by eating an individual of *Rhinella diptycha* may produce local irritation to the caiman, and even lead it to death shortly after ingestion. To avoid this threat, some animals use particular feeding strategies, such as starting to eat the prey's posterior part, or eating only its viscera. In this note, we report the first case of a predation event of an adult individual of the poisonous toad *R. diptycha* by a young individual of *C. latirostris* that we observed in a pond of the Southeast of Brazil. The caiman waited until the toad was dead before starting swallowing slowly, from the anterior part of the body. We did not observe any sign that the caiman was affected by the prey ingestion, even some hours later after ingestion. Our register suggests that the caiman may be tolerant or adapted to eat this poisonous prey and encourage research on the causes and mechanisms of this tolerance.

Keywords: caiman, ecological interaction, poison, predation, toad

Resumen

Los caimanes de la especie *Caiman latirostris*, distribuidos ampliamente por América del Sur, son depredadores oportunistas. Estos animales presentan variación ontogenética en su dieta de forma que los individuos jóvenes, que se alimentan inicialmente de invertebrados, van incorporando vertebrados a la dieta de manera gradual, a medida que crecen. Sin embargo, la falta de experiencia lleva, en ocasiones, a los caimanes jóvenes a consumir presas peligrosas, como anfibios venenosos. La intoxicación causada por ingerir un individuo de *Rhinella diptycha* puede ocasionar irritaciones locales al caimán, e incluso llevarle a la muerte. Para evitar esta amenaza, algunos animales utilizan estrategias de alimentación especiales, como comenzar a ingerir la presa por la parte posterior o comer solamente las vísceras. En esta nota presentamos el primer registro de un evento de depredación de un individuo adulto del sapo venenoso *R. diptycha* por parte de un individuo joven de *C. latirostris*, que observamos en una charca del sudeste de Brasil. El caimán esperó a que el sapo estuviera muerto antes de comenzar a tragarlo, lentamente, comenzando por la parte anterior de su cuerpo. No observamos ninguna señal de que la ingestión de la presa afectara al caimán, incluso algunas horas después de haberla ingerido. Nuestro registro sugiere que el caimán podría ser tolerante o estar adaptado a comer este tipo de presas venenosas, por lo que recomendamos investigar las causas y mecanismos de este tipo de tolerancia.

Palabras claves: anuros, caimán, depredación, interacción ecológica, veneno

¹. Faculdade de Ciências Biológicas e Ambientais, Universidade Federal da Grande Dourados, Brazil.

². Programa de Pós-Graduação em Biologia Animal, Universidade Federal de Mato Grosso do Sul, Brazil.

³. Programa de Pós-Graduação em Ecologia e Conservação, Universidade Federal de Mato Grosso do Sul, Brazil.

* Author for correspondence: <couto.murillo@gmail.com>

Brazil holds the world's greatest diversity of caimans, with six known species (Costa and Bernils, 2018). Caimans are opportunistic predators, eating almost any capturable prey they find, sometimes even eating individuals of their respective own species (Magnusson, Da-Silva, and Lima, 1987; Piña, Larriera, and Cabrera, 2003; Santos et al, 1996). Caimans show an ontogenetic variation of diet, with young individuals preferring invertebrates and gradually incorporating vertebrates as they grow up (Ortiz, Charrau, and Reynoso, 2020). However, eating poisonous amphibians may be dangerous. For example, the population densities of *Crocodylus johnstoni* in Australia declined in almost 80% due to the invasive poisonous toad *Rhinella marina* (Letnic, Webb, and Shine, 2008). Amphibians are widely distributed, vary in size, and are potential prey for many vertebrate and invertebrate species (Duellman and Trueb, 1994). Most predatory events of anurans are reported during the reproductive season, where many animals gather together. Despite the importance of identifying the predators of amphibians, researchers have difficulties making observations of predation in nature (Pombal Jr., 2007). Toads of the family Bufonidae reproduce at the end of the cold and dry season, probably as a strategy for the tadpoles to develop during the wet season, when food is abundant (Oda, Bastos, and Lima, 2009; Santos, Rossa-Feres, and Casatti, 2007). Bufonids are considered true toads due to the presence of an agglomerate of poison glands on their body. Specifically, the individuals of *R. diptycha* have the parotid glands behind their eyes, and the paracnemid glands on their hind legs, as well as many other small poison gland clusters scattered throughout their bodies (Frost, 2009; Jared et al, 2009). These glands are a passive defense for the toad that releases the poison when the gland is pressed (Jared et al, 2009). The potential predator would pass from local irritation to even death in less than 15 minutes after the intoxication (Knowles, 1968; Micuda, 1968; Oehme, Brown, and Fowler, 1980; Otani, Palumbo, and Read 1969). The broad-snouted-caiman, *Caiman latirostris*, is one of the most widely distributed caimans of South America, being present from the northeast of Brazil, through Bolivia and Paraguay, to the northeast of Argentina and Uruguay (Verdade, Larriera, and Piña, 2010). Here we report the first event of an individual of *C.*

latirostris feeding on *R. diptycha*.

The event took place in September 2019, at 20:00 h, during a nighttime sampling of reptiles (SISBIO 64074), in a swamp (20° 35' 14.87"S and 51° 21' 29.73"W) at the municipality of Ilha Solteira, at the northwest of the state of São Paulo (Brazil). An individual of *C. latirostris* was detected by the reflection of the lantern's light by its eyes. As we approached, we perceived that the individual was eating an adult male of *R. diptycha*, biting it through the left anterior region. The individual of *C. latirostris* was about 80 cm (total length), and thus considered a young individual (Verdade, 1995). The *R. diptycha* toad was an adult male, of approximately 15 cm long. The whole predatory event, from the death of the prey to its complete ingestion, took approximately 45 minutes. From the time we began the observation, the toad remained swollen for approximately one minute, until it died, with the caiman taking from the mouth, partially submerged (figure 1a). Once the toad died, the caiman started moving its mouth, biting the prey until it managed to position it with the anterior part inside its mouth (figure 1b). The caiman moved about, adjusting the toad with its mouth four or five times, and then standing still for about five minutes. We noticed that the caiman was breathing very quickly. When first spotted, the caiman was in a shallow pond with muddy water and clay (figure 1c). Over the course of the observation, the individual moved about a meter in direction to the shore, due to interference by the flash of the photographic camera. Once the caiman ate the whole toad, without tearing parts or limbs, the crocodilian walked slowly to the marsh, where it remained at rest. When we finished the fieldwork, two hours later, we passed by the pond again to check for the presence of the individual. The caiman was still near the place, at rest. None of the animals were manipulated or collected, having as testimonial material the photographs, deposited in the Reference Zoological Collection ZUFMS (ZUFMS-AMP12942; ZUFMS-AMP12943 e ZUFMS-REP03457).

In general, anurans are part of the diet of several animal species. Animals eat poisonous amphibians by their posterior region to avoid contact with the poison (Haddad and Bastos, 1997; Loebmann, Solé,

and Kwet, 2008; Toledo, 2003). Amongst vertebrates, some species are adapted to eat bufonid toads, as is the case of snakes of the genus *Xenodon*, which have a modified adaptation to pierce the toad's lung and are little susceptible to its poison (Lavilla, Schrocchi, and Terán, 1979). Other animals, such as raccoons and otters, avoid intoxication by skinning the toad, taking the viscera out and avoiding ingestion of the dorsal skin (Morales, Ruiz-Olmo, Lizana, and Gutiérrez, 2016). The Australian rodent *Hydromys chrysogastes* avoids intoxication by making a small incision in the toad's ventral area and only eating some internal organs (Parrott, Doody, McHenry, and Clulow, 2019). Similarly, some raptor birds eat the viscera and avoid eating the dorsal skin (Crozarior, and Gomes, 2009; Röhe, and Pinassi-Antunes, 2008).

Despite the different strategies used to eat bufonid toads, unsuccessful predation has been reported, such as an individual of *Ceratophrys aurita*, which ingested a specimen of *R. diptycha* and died within a day and a half after ingestion (Silva-Soares, Mônico, Ferreira, and De Castro, 2016). Records of predation and ingestion of poisonous amphibians are seldom reported in caimans. Among those reported, there are the predatory events of *R. granulosa* and *R. marina* by *Caiman crocodilus crocodilus* (Gorzula, 1978; Morato, Batista, and Paz, 2011); *R. marina* by *Paleosuchus trigonatus* (De Assis, and Dos Santos, 2007) and *R. diptycha* by *P. palpebrosus* (Toledo, Ribeiro, and Haddad, 2007). Although the presence of frogs in the diet of *C. latirostris* is already known, its actual frequency may be underestimated in the




Figure 1. Predation of *Rhinella diptycha* by *Caiman latirostris* in a pond of Ilha Solteira (São Paulo, Brazil). a. The moment we spotted the predatory event; b. *R. diptycha* is positioned to be ingested by *C. latirostris*; c. The pond where the event was observed; d. An adult female of *R. diptycha* was found a few meters away from the predatory event.

published literature due to the rapid digestion of this type of prey (Borteiro, Gutiérrez, Tedros, and Kolenc, 2009; Melo, 2002).

We think that the caiman might have been attracted to the prey by its vocalization, since we found a female *R. diptycha* nearby (figure 1d). Even though mating vocalization has significant advantages for anurans to attract sexual mates, it increases exposition to predators (Toledo, 2003). This is the first record of *C. latirostris* preying upon a specimen of the poisonous anuran *R. diptycha*. Unlike the case of *C. johnstoni*, who died by eating introduced *R. marina* toads in Australia (Letnic et al, 2008), our register suggests the caiman was unaffected by the poisonous toad. This may be explained because *R. diptycha* is a native species, probably coexisting for millions of years with *C. latirostris* at the study area. We recommend future research confirming the potential tolerance of *C. latirostris* to *R. diptycha*'s poison and assessing the causes and mechanisms that make it possible.

ACKNOWLEDGEMENTS

Maiara Cabrera Miguel and Zaida Ortega are grateful to Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) by funding two of the authors. Zaida Ortega was funded by a postdoctoral fellowship PNPd/CAPES (#1694744).

CONFLICTS OF INTEREST

The authors express that there is no conflict of interest.

REFERENCES

- Borteiro, C., Gutiérrez, F., Tedros, M., and Kolenc, F. (2009). Food habits of the broadsnouted caiman (*Caiman latirostris*: Crocodylia, Alligatoridae) in northwestern Uruguay. *Studies on Neotropical Fauna and Environment*, 44(4), 31-36. doi:10.1080/01650520802507572
- Costa, H.C., and Bérnils, R.S. (2018). Répteis do Brasil e suas Unidades Federativas: Lista de espécies. *Herpetologia Brasileira*, 7(1), 11-57.
- Crozariol, M.A., and Gomes F.B.R. (2009). Predação de *Leptodactylus fuscus* (Schneider, 1799) (Anura: Leptodactylidae) por *Caracara plancus* (Miller, 1777) (Aves: Falconidae), com nota sobre história natural, no Vale do Paraíba do Sul, SP. *Atualidades Ornitológicas*, 154, 4-5. Retrieved from <http://www.ao.com.br/ao149.htm>
- De Assis, V.B., and Dos Santos, T. (2007). *Paleosuchus trigonatus* predation. *Herpetological Review*, 38(4), 445. <https://ssarherps.org/herpetological-review-pdfs/>
- Duellman, W.E., and Trueb, L. (1994). *Biology of amphibians*. (2nd ed.) Baltimore, Maryland, USA: The Johns Hopkins University Press.
- Frost, D.R. (2009). *Amphibian species of the world: an online reference*. Version 5.3. New York, USA: American Museum of Natural History [Electronic database]. Retrieved from <http://research.amnh.org/herpetology/amphibia/>
- Gorzula, S.J. (1978). An ecological study of *Caiman crocodilus crocodilus* inhabiting savanna lagoons in the Venezuelan Guayana. *Oecologia*, 35(1), 21-34. doi:10.1007/BF00345539
- Haddad, C.F.B., and Bastos, R.P. (1997). Predation on the toad *Bufo crucifer* during reproduction (Anura: Bufonidae). *Amphibia-Reptilia*, 18, 295-298. doi:10.1163/156853897x00170
- Jared, C., Antoniazzi, M.M., Jordão, A.E.C., Silva, J.R.M., Greven, C.H., and Rodrigues M.T. (2009). Parotoid macroglands in toad (*Rhinella jimi*): Their structure and functioning in passive defence. *Toxicon*, 54, 197-207. doi:10.1016/j.toxicon.2009.03.029
- Knowles, R.P. (1968). Toad poisoning in dogs. *Journal of the American Veterinary Medical Association*, 153, 1202.
- Lavilla, E.O., Schrocchi, G.J., and Terán, E.M.T. (1979). Sobre algunos aspectos del comportamiento em cautiverio de *Xeneodon merremi* (Wagler) (Ophidia: Colubridae). *Acta Zoologica Lilloana*, 35, 287-293.
- Letnic, M., Webb, J.K., and Shine, R. (2008). Invasive cane toads (*Bufo marinus*) cause mass mortality of freshwater crocodiles (*Crocodylus johnstoni*) in tropical Australia. *Biological Conservation*. 141(7), 1773-1782. doi:10.1016/j.biocon.2008.04.031
- Loebmann, D., Solé, M., Kwet, A. (2008). Predation on spawn and adults of *Chaunus Dorbignyi* (Duméril and Bibron, 1841) (Amphibia, Anura) by leeches (Hirudinea) in southern Brazil. *Amphibia*, 7, 31-34. doi:10.1590/S1519-69842001000300012
- Magnusson, W.E., Da-Silva, E.V., and Lima, A.P. (1987). Diets of Amazonian crocodiles. *Journal of Herpetology*, 21(2), 85-95. doi:10.2307/1564468
- Melo, M.T.Q. (2002). Dieta do *Caiman latirostris* no sul do Brasil. In L.M. Verdate, and A. Larriera (Eds.). *Conservação e manejo de jacarés e crocodilos da América Latina* (2nd edition, pp 116-125). Piracicaba (Brazil): C.N. Editoria.
- Micuda, J. (1968). Toad poisoning. In H.P. Hoskins (Ed.). *Canine medicine* (1st edition, p. 260). Santa Barbara: American Veterinary Publications.
- Morales, J., Ruiz-Olmo, J., Lizana, M., and Gutiérrez, J. (2016). Skinning toads is innate behaviour in otter (*Lutra lutra*) cubs. *Ethology, Ecology and Evolution*, 28, 414-426. doi:10.1080/03949370.2015.1076525
- Morato, S.A.A., Batista, V.B.G.V., and Paz, A. (2011). *Paleosuchus trigonatus* (smooth-fronted caiman): diet and movement. *Herpetological*

- Bulletin*, 115: 34-35. <https://www.thebhs.org/publications/the-herpetological-bulletin/issue-number-115-spring-2011/104-08-natural-history-notes-2/file>
- Oda, F.H., Bastos, R.P., and Lima, M.A.C.S. (2009). Anuran assemblage in the Cerrado of Niquelândia, Goiás State, Brazil: diversity, local distribution and seasonality. *Biota Neotropical*, 9, 219-232. doi:10.1590/S1676-06032009000400022
- Oehme, F.W., Brown, J.F., and Fowler, M.E. (1980). Toxins of animal origin. In L.J., Casarett, C., Klaassen, J., Watkin, and J., Doull (Eds.). *Casarett and Doull's toxicology: The basic science of poisons* (pp. 557-77). New York, USA: McGraw-Hill Professional.
- Ortiz, G.J.S., Charruau, P., and Reynoso, V. (2020). Variation in diet of hatchlings, juveniles, and subadults of *Caiman crocodilus chiapasius* in La Encrucijada, Chiapas, Mexico. *Revista Mexicana de Biodiversidad*, 91, e91852. doi:10.22201/ib.20078706e.2020.91.2852
- Otani, A., Palumbo, N.E., and Read, G. (1969). Pharmacodynamics and treatment of mammals poisoned by *Bufo marinus* toxin. *American Journal of Veterinary Research*, 30(10), 1865-1872. <https://europepmc.org/article/med/10200112>
- Parrott, M.L., Doody, J.S., McHenry, C., and Clulow, S. (2019). Eat your heart out: choice and handling of novel toxic prey by predatory water rats. *Australian Mammalogy*, 42, 235-239. doi:10.1071/AM19016
- Piña, C.I., Larriera, A., and Cabrera, M.R. (2003). The effect of incubation temperature on hatching success, incubation period, survivorship and sex ratio in *Caiman latirostris* (Crocodylia, Alligatoridae). *Journal of Herpetology*, 7, 199-202. doi:10.1670/0022-1511(2003)037[0199:EOITOI]2.0.CO;2
- Pombal Jr, J.P. (2007). Notas sobre predação em uma taxocenose de anfíbios anuros no sudeste do Brasil. *Revista Brasileira de Zoologia*, 24, 841-843. doi:10.1590/S0101-81752007000300034
- Röhe, F., and Pinassi-Antunes, A.P. (2008). Barred Forest Falcon (*Micrastur ruficollis*) Predation on relatively large prey. *Wilson Journal of Ornithology*, 120, 228-230. Retrieved from <http://hdl.handle.net/11449/39793>
- Santos, S.A., Stoll, M., Silva, M., Campos, Z., Magnusson, W.E., and Mourão, G. (1996). Diets of *Caiman crocodilus yacare* from different habitats in the Brazilian Pantanal. *Herpetological Journal*, 6, 111-117.
- Santos, T.G., Rossa-Feres, D.C., and Casatti, L. (2007). Diversidade e distribuição espaço-temporal de anuros em região com pronunciada estação seca no sudeste do Brasil. *Iheringia. Série Zoologia*, 97, 37-49. doi:10.1590/S0073-47212007000100007
- Silva-Soares, T., Mônico, A.T., Ferreira, R.B., and De Castro, T.M. (2016). *Ceratophrys aurita* (Sapo-de-chifre: Brazilian Horned Frog). Mortality. *Herpetological Review*, 47(4), 641. <https://ssarherps.org/herpetological-review-pdfs/>
- Toledo, L.F. (2003). Predation on seven South American anuran species by water bugs (Belostomatidae). *Phyllomedusa: Journal of Herpetology*, 2, 105-108. doi:10.11606/issn.2316-9079.v2i2p105-108
- Toledo, L.F., Ribeiro, R.S., and Haddad, C.F.B. (2007). Anurans as prey: an exploratory analysis and size relationships between predators and their prey. *Journal of Zoology*, 271, 170-177. doi:10.1111/j.1469-7998.2006.00195.x
- Verdade, L. M. (1995). Biología reproductiva do jacaré-de-papo-amarelo (*Caiman latirostris*) em São Paulo, Brasil. In A, Larriera and L.M, Verdade (Eds.). *Conservación y manejo de los Crocodylia de America Latina* (pp. 57-79). Santo Tomé, Santa Fe, Argentina: Fundación Banco Bica.
- Verdade, L.M., Larriera, A., and Piña, C.I. (2010). Broad-snouted Caiman *Caiman latirostris*. In S.C, Manolis and C, Stevenson (Eds.). *Crocodyles. Status survey and conservation action plan* (3rd edition, pp. 18-22). Gland, Switzerland: UICN/SSC Crocodile Specialist Group.