

Effects of phosphorus and potassium levels on the yield of the tuber variety Criolla Colombia in the department of Cundinamarca

Efecto de niveles de fósforo y potasio sobre el rendimiento de la variedad Criolla Colombia en el departamento de Cundinamarca

Yohana Carolina Rozo M.¹ and Carlos Eduardo Núñez L.^{1,2}

ABSTRACT

In the locations of Zipaquirá and Cogua located in the department of Cundinamarca, the effects of applying different levels of phosphorus and potassium on tuber yield and specific gravity in the diploid potato variety Criolla Colombia were evaluated. The design used a completely randomized block with three replications and a 4 x 4 factorial structure, where the first factor corresponds to phosphorus (0, 50, 100 and 150 kg ha⁻¹ P₂O₅), and the second, potassium (0, 50, 100 and 150 kg K₂O ha⁻¹). The variables evaluated were specific gravity and tuber yield in the categories: first (PT1), second (PT2), third (PT3) and total (PTT). In Zipaquirá, there were differences in the phosphorus factor for the yield variables PT1, PT2 and PTT, while in the town of Cogua, the only difference for this factor was found in the variable PTT. In these variables, the levels of 50, 100 and 150 kg ha⁻¹ of P₂O₅ produced higher yields that were equal and above that of the 0 kg ha⁻¹ P₂O₅ level. In the potassium factor, differences were found only for the yield variable PT3 in the town of Cogua. The specific gravity did not respond to the P or K factors evaluated.

Key words: fertilization, specific gravity, *Solanum phureja*, diploid potato.

RESUMEN

En las localidades de Zipaquirá y Cogua de departamento de Cundinamarca, se evaluó el efecto de aplicación de diferentes niveles de fósforo y potasio sobre el rendimiento de tubérculo y la gravedad específica en la variedad de papa diploide Criolla Colombia. Se utilizó el diseño de bloques completamente al azar con tres repeticiones y estructura factorial 4 x 4, donde el primer factor corresponde a fósforo (0, 50, 100 y 150 kg ha⁻¹ de P₂O₅), y el segundo, potasio (0, 50, 100 y 150 kg ha⁻¹ de K₂O). Las variables evaluadas fueron gravedad específica y rendimiento de tubérculo en las categorías primera (PT1), segunda (PT2), tercera (PT3) y total (PTT). En Zipaquirá se encontró diferencias en el factor fósforo para las variables de rendimiento PT1, PT2 y PTT, mientras que en la localidad de Cogua sólo se encontró diferencias en este factor para la variable PTT. En estas variables los niveles 50, 100 y 150 kg ha⁻¹ de P₂O₅ fueron iguales entre sí y superiores al nivel 0 kg ha⁻¹ de P₂O₅. En el factor potasio sólo se encontró diferencias para la variable de rendimiento PT3 en la localidad de Cogua. La gravedad específica no presentó respuesta a los factores de P y K evaluados.

Palabras clave: fertilización, gravedad específica, *Solanum phureja*, papa diploide.

Introduction

The species *Solanum phureja* (2x) Juz. et Buk. is classified within the Solanaceae family, genus *Solanum*, section Petota series Tuberosa, and, together with the species *S. tuberosum* subsp. *andigena*, constitutes the diversity of potatoes grown in Colombia (López, 1993). Its name derives from the Aymara word “*phureja*” meaning early. This diploid has a single branched stem, slightly bright leaves; the tubers have no rest period and have more protein than the tubers of the species *S. tuberosum* (Luján, 1996). Its center of diversity is located in the department of Nariño, in southern Colombia, to the north of Ecuador. In addition,

the widely used taxonomic classification of Hawkes (1990) recognizes it as one of seven cultivated species (*Solanum phureja* with three subspecies); Huaman and Spooner (2002), based on a study of morphological characters, designated it as Group Phureja within a simple designation of the *Solanum tuberosum* class. Spooner *et al.* (2007), with data from molecular studies, suggested that *S. phureja* remain included within the classification of *Solanum tuberosum* Group Andigenum.

The species *S. phureja* is a genetic resource of national importance because its diverse genotypes have been identified with high protein content (Rodríguez *et al.*, 2006),

Received for publication: 18 May, 2009. Accepted for publication: 2 June, 2011.

¹ Department of Agronomy, Faculty of Agronomy, Universidad Nacional de Colombia. Bogota (Colombia).

² Corresponding author. cenuztezl@unal.edu.co

and because of its sources of resistance to *Phytophthora infestans* (Estrada, 2000) and PVV (Vargas, 2010), pleasant flavor and texture, easy preparation, good acceptance in the market and high export potential in various forms of processing (Rivera *et al.*, 2006).

The criolla potato is produced in different regions of Colombia, mainly in the valleys. Its quality depends on the microclimate, altitude, solar radiation and humidity during the production cycle (Carmona *et al.*, 1998). Despite its phenotypic heterogeneity, in Colombia, a small fraction of the criolla potato is processed into a precooked-frozen or pickled form for the domestic or export market (Rodríguez *et al.*, 2009).

According to the first national census of the potato crop conducted by DANE in Cundinamarca in the agricultural year 2001-2002, the criolla potato ranked third in harvest area after the varieties Parda Pastusa and Diacol Capiro with an average yield of 15.3 t ha⁻¹ (Fedepapa, 2003). Colombia is the largest producer, consumer and exporter of diploid potatoes in the world, with the competitive advantages of being a center of diversity and having wide consumer acceptance due to its organoleptic characteristics (Rodríguez *et al.*, 2009).

To exploit the genetic potential of crops, the optimum levels of potassium, nitrogen and phosphorus are required to achieve better quality and yield of the tuber. After nitrogen, phosphorus is the element that is most often limited in soils. It is absorbed primarily as the monovalent phosphate anion (H₂PO₄⁻) and less rapidly as the divalent anion (HPO₄²⁻). Soil pH controls the relative abundance of these two forms (H₂PO₄⁻ is favored at a pH below 7 and HPO₄²⁻ above). Much of the phosphate is converted into organic forms as it enters the root or after it is transported through the xylem to the stem or leaves. Phosphorus is never reduced in plants, where it remains as phosphate, either free or bound in organic forms such as esters (Salisbury and Ross, 1994).

Phosphorus performs functions in plants, such as a structural element forming part of the macromolecular structures such as nucleic acids (RNA and DNA) and in the phospholipids of cell membranes. It also has an important function in energy transfer such as in the esters of energy-rich phosphate present in the metabolic mechanisms of the cells, within which ATP participates in the main metabolic processes such as photosynthesis and respiration. Optimum plant growth is accompanied by a demand for P to meet the demands for the functions described above (Marschner, 2002) and so depends on the availability of P

in the soil, and the ability of the plant to absorb P in the soil (Álvarez-Sánchez *et al.*, 1999). Many factors influence the supply of P to plants. Among the most important are the availability of P in the soil, temperature, root density and root efficiency in P uptake (Pessarakli, 2001). Potato fertilization usually requires high amounts of phosphate fertilizer (60 to 80 kg ha⁻¹ P) to achieve economically acceptable yields, necessitated by the low root density of this plant (Álvarez-Sánchez *et al.*, 1999) and because P diffusion moves very slowly in soil, which causes difficulties in P absorption by the plant (Barber, 1994).

Young plants with a P deficiency are bluish green in color in the early stages of growth. Older leaves with deficiency symptoms usually appear dark green (Mengel *et al.*, 2001). Phosphorus deficiency in the potato slows apical growth, resulting in small, rigid plants; reduces the formation of starch in the tubers causing necrotic spots distributed in the tuber (Pumisacho and Sherwood, 2002); decreases CO₂ absorption capacity of leaf photosynthesis; results in an inadequate supply of phosphates which prevents the export of triose phosphate from the chloroplast, and therefore, affects the synthesis of sucrose (Mengel *et al.*, 2001), and causes delayed development of the tubers (McCollum, 1978).

In a soil clay-sandy with medium to high concentrations of P, the effects of different doses of fertilizer and sources of P on the variety Russet Burbank were analyzed. The numbers and yield of small tubers increased as the dose of P was increased, indicating that P may play an important role in regulating the number of tubers per plant (Rosen and Bierman, 2008). In soil of Turkey, poor in organic matter, slightly alkaline, with moderate levels of P and high levels of K, three doses of N with three doses of P (0, 90 and 180 kg ha⁻¹) with the Agria variety were evaluated. In this study, an effect of phosphorus levels on the oil content in fried potato chips was observed, lessening as the dose of P increased. Doses had no effect on dry matter, starch content or protein content. In studies conducted in different localities with the variety Diacol Capiro in Colombia, a response to P fertilization up to the level of 200 kg ha⁻¹ has been reported; no response was reported in the variable specific gravity (Núñez *et al.*, 2006).

The potato (*Solanum tuberosum*) has a relatively high potassium requirement (K), which has led to the suggestion that high doses of the element are needed for potato production (Panique *et al.*, 1997). Potassium is a monovalent cation and its capture is highly selective; is coupled with metabolic activity. It is characterized by high mobility in plants at all levels; is the most abundant cation in the cytoplasm

and along with its accompanying anions makes a high contribution to the osmotic potential of cells and tissues. This element has an important role in water relations of the plant, furthermore K is not metabolized and forms easily interchangeable weak complexes (Marschner, 2002). It stimulates the activity of the enzyme associated with starch synthesis (starch synthase) (Mengel and Kirkby, 1987), in addition, it facilitates the translocation of assimilates to the tubers, which ultimately increases the bulking capacity of the tuber and its biomass.

Potassium deficiency can result in a decrease in yield and tuber size McDole *et al.* (1978). Some quality factors such as dry tuber, specific gravity, sugar content, flesh color and hollow heart are affected by K fertilization (Chapman *et al.*, 1992). Numerous studies have reported reduction in specific gravity or dry matter content by the increasing application of potassium (Laboski and Kelling, 2007). Soils deficient in this element reportedly increase the specific gravity of tubers (Chapman *et al.*, 1992). Davenport and Bentley (2001) concluded that the source of K does not affect the specific gravity of the tuber. Contrary results were reported in a later study by Kumar *et al.* (2007) with varieties of India where the specific gravity and dry matter were higher with a sulfate and nitrate source than a chloride source, while the quality of processing, tuber yield and biomass were similar for the chloride and sulfate sources but lower for the nitrate source.

According to the ICA (1992), critical levels of available P for potato determined by the Bray II method for the Cundiboyacense plateaus are: a) low below 40 mg kg⁻¹, b) medium between 40 and 60 mg kg⁻¹ and c) high over 60 mg kg⁻¹. Critical levels of K in the Cundiboyacense plateaus are: a) low under 0.30 cmol kg⁻¹, b) medium: between 0.30 and 0.60 cmol kg⁻¹ and c) high: over 0.60 cmol kg⁻¹ (ICA, 1992). The above critical levels were determined for *Solanum tuberosum* subsp. *andigena*, no guide exists for the diploid varieties of *S. phureja*.

The fertilizer use in criolla potato cultivation in the cundiboyacense plateaus during semester A-2007 accounted for 37.7% of the total input costs, 16.9% of the direct costs and 13.7% of the total costs of the crop (Villarreal *et al.*, 2007). Its high production costs necessitate the need for research in the area of crop nutrition and fertilization, to determine the needs of the different varieties and different growing areas, and thus find a more efficient production system.

In Colombia, research on the species *Solanum phureja* has been carried out, however very few are related to the area

of plant nutrition, so, little is known about its fertilization needs. Pérez *et al.* (2008) reported for the Criolla Colombia variety that fertilization without N-P-K-Mg fractionation favored foliar development and yield potential, in addition, the application of boron is found to invoke positive responses in tuber yield. Arias *et al.* (1996) evaluated the performance of the criolla potato under different planting densities on the Bogota Plateau, in the Centro Agropecuario Marengo -CAM (Mosquera, Colombia at 2,570 m a.s.l.) in a soil with high levels of P and K, using a fertilizer with a 13-26-6 compound in doses of 500 kg ha⁻¹ (130 kg ha⁻¹ of P₂O₅ and 30 kg ha⁻¹ K₂O). The study reported yields of 6.8 t ha⁻¹ in the first category, 3.5 t ha⁻¹ in the second category, 2.5 t ha⁻¹ in the third category and 12.5 t ha⁻¹ in overall yield.

The objective of this study was to evaluate the effects of applying different levels of phosphorus and potassium on tuber yield and specific gravity on the local variety Criolla Colombia (*Solanum phureja* Juz *et. Buk*).

Materials and methods

Location

The present study was carried out at two localities in the department of Cundinamarca, the first in the town of Zipaquira (2,590 m a.s.l.) and the second in the municipality of Cogua (2,690 m a.s.l.); the soil analysis of these two locations is shown in Tab. 1. The tests were carried out in the second half of 2004.

Area precipitation data

The locations of the evaluations were in the municipalities of Cogua and Zipaquira which belong to the Sabana Centro area in the department of Cundinamarca, characterized by a cold semi-humid climate (dry forest, low mountain), an average temperature range of between 12 and 18°C and an average annual rainfall of between 500 to 1,000 mm with a similar historic distribution in the first and second halves of the year (SADC, 2008).

Experimental design

The experiment was performed using a completely randomized block design with three replications and a structural factorial of 4x4. The first factor corresponds to phosphorus (0, 50, 100 and 150 kg ha⁻¹ of P₂O₅) and the second to potassium (0, 50, 100 and 150 kg K₂O ha⁻¹). The experimental unit was 32 m², with row spacing of 0.8 m and 0.30 m between sites. The sources of fertilizers used were urea, triple superphosphate and potassium sulphate.

TABLE 1. Soil analysis of the locations Zipaquirá and Cogua (Colombia).

Location	pH	CO	N	Interchangeable bases				Exchangeable acidity		CEC
				Ca	K	Mg	Na	Al	H	
			(cmol kg ⁻¹)							
Zipaquirá	4.89	3.88	0.33	7.68	1.16	1.89	0.10	0	0.15	11.9
Cogua	5.11	5.90	0.51	5.05	1.72	1.56	0.31	0.65	0.27	34.4

Location	P	Microelements					Ar	L	A	Texture
		Cu	Fe	Mn	Zn	B				
			(mg kg ⁻¹)							
Zipaquirá	14.2	1.33	695	13.35	22.80	0.36	23.2	47.6	29.2	Loam
Cogua	23.9	2.48	373	10.25	1.83	0.23	27.7	34.0	38.3	Clay Loam

Methods of analysis. Organic Carbon (CO): Walkley-Black. Ca, Mg, Na, K: Extraction with 1N ammonium acetate pH 7. Al (interchangeable acids) Extraction: with 1M KCl. Cation Exchange Capacity (CEC): Displacement of exchanged NH₄ with 1M NaCl. Useful P: Bray II. Mn, Zn, Fe, Cu: Extraction with DTPA. Boron (B) Extraction with phosphate monobasic (Azomethine-H).

Variables evaluated

Tuber yield: The tubers were classified into three categories according to the diameter of the tuber: the first category (PT1:> 4 cm), second (PT2: 2 to 4 cm) and third (PT3: <2 cm). At harvest, the tubers were weighed by category and added to obtain the total, this variable was analyzed as t ha⁻¹.

Specific Gravity (SG): 10 tubers were sampled per plot from the first category to determine the specific gravity using the method of weight in water (Ww) and weight in air (Wa). The SG was calculated by Talburt and Smith (1975).

$$[SG = Wa / (Wa - Ww)] \quad (1)$$

Statistical analysis was performed using analysis of variance and multiple comparison test of Tukey, using the SAS software version 9.2.

Results and discussion

Zipaquirá

At this location, differences were found for the phosphorous factor for the yield variables in the categories: first (PT1),

second (PT2) and total (PTT) (Tab. 2). No differences were observed for the potassium factor or the phosphorus x potassium interaction for any of the variables evaluated (Tab. 2).

In the variables PT1, PT2 and PTT, phosphorus levels 50, 100 and 150 kg ha⁻¹ P₂O₅ produced higher yields that were equal and above that of the level 0 kg P₂O₅ ha⁻¹ (Figs. 1, 2 and 3). Keep in mind that the level of phosphorus found in this location (Zipaquirá) was 14.2 mg kg⁻¹ (Tab. 1), which is considered a low level of this element, which would explain the positive response to the application of P levels greater than 0 kg ha⁻¹ P₂O₅. However, the application of levels greater than 50 kg ha⁻¹ P₂O₅ did not present differences in tuber production of these yield variables, which can mean that the variety requires a low level of P or is very efficient at taking in the nutrient.

There were no differences for the variable specific gravity in any of the factors evaluated or for the potassium x phosphorus interaction (Tab. 2). At this site, the values of this variable were between 1.063 and 1.068, relatively low values for the variety, taking into account the reports from

TABLE 2. Analysis of variance (P) in the yield components for ‘Criolla Colombia’ in Zipaquirá and Cogua.

LOCATION	VARIABLE	BLOCK	P	K	P x K
Zipaquirá	PT1	0.8561	<.0001*	0.1903	0.2106
	PT2	0.5742	0.0156*	0.7843	0.3073
	PT3	0.7904	0.9575	0.8798	0.5374
	PTT	0.5618	<.0001*	0.1991	0.1510
	GE	0.0695	0.4051	0.2845	0.0978
Cogua	PT1	0.5195	0.1163	0.6043	0.5114
	PT2	0.4618	0.0758	0.2461	0.0718
	PT3	0.0166*	0.3439	0.0064*	0.2532
	PTT	0.7432	0.0185*	0.8061	0.3864
	SG	0.0524	0.4927	0.3124	0.4775

PT1, the first category; PT2, the second category; PT3, the third category; PTT, total yield; P phosphorus factor levels; K, potassium factor levels; P x K interaction of P and K.

* Difference statistically significant (P≤0.05).




FIGURE 1. Yield for potato 'Criolla Colombia', first category (PT1) with different levels of P in the town of Zipaquirá. Means with different letters indicate significant difference according to Tukey test ($P \leq 0.05$).




FIGURE 2. Yield for potato 'Criolla Colombia', second category (PT2) with different levels of P in the town of Zipaquirá. Means with different letters indicate significant difference according to Tukey test ($P \leq 0.05$).




FIGURE 3. Total yield (PTT) for potato 'Criolla Colombia' with different levels of P in the town of Zipaquirá. Means with different letters indicate significant difference according to Tukey test ($P \leq 0.05$).




FIGURE 4. Yield for potato 'Criolla Colombia', third category (PT3) with different levels of K in the town of Cogua. Means with different letters indicate significant difference according to Tukey test ($P \leq 0.05$).

Rodríguez *et al.* (2009) in evaluations in the department of Antioquia.

In this location, the level of soil K was 1.16 cmol kg⁻¹, which is high in these soils for potato cultivation in the departments of Cundinamarca and Boyacá (ICA, 1992), this explains the absence of a response to K including up to the level of 150 kg ha⁻¹. Similarly, Becerra *et al.* (2007) reported that in the local variety Criolla Guaneña, in soil with an average level of K in the municipality of Pasto (Nariño), there was no response to two levels of K (50 and 100 kg ha⁻¹ K₂O), and 50 kg ha⁻¹ K₂O was enough for proper growth and high yield.

Cogua

The differences found in the phosphorus factor for the variable total tuber yield (PTT) are explained by the difference between the 100 kg ha⁻¹ of P₂O₅ (34.8 t ha⁻¹) level and the 0 kg ha⁻¹ of P₂O₅ (26.8 t ha⁻¹) level (Tab. 2 and Fig 5). No interaction (P x K) was found for any of the yield variables evaluated (Tab. 2). In the potassium factor, the only significant difference in a yield variable found was in the third




FIGURE 5. Total yield (PTT) for potato 'Criolla Colombia' with different levels of P in the town of Cogua. Means with different letters indicate significant difference according to Tukey test ($P \leq 0.05$).

category (PT3), explained by the difference between the 150 kg ha⁻¹ of K₂O (0.72 t ha⁻¹) level and the 0 kg ha⁻¹ of K₂O (1.36 t ha⁻¹) level (Tab. 2 and Fig 4), this means that without a significantly increased application of K in proportion to the small sized potato, which is generally unfavorable to the farmer, the value and weight of the higher-priced commercial potato will be reduced (PT1 or PT2). On the other hand, if the aim of production is processing (potato for liquor), the absence of K fertilization may be favorable.

The variable specific gravity did not respond to the application of P and K, and had no P x K interaction (Tab. 2), the range observed for this variable in this locality was between 1.081 and 1.087, higher than observed in Zipaquirá and similar to the reports from Rodríguez *et al.* (2009) in evaluations of the variety in the department of Antioquia. The level of P in the soil of this town was 23.9 mg kg⁻¹ (a slightly higher value than that found in the town of Zipaquirá Tab. 1), which is low in soils of Cundinamarca and Boyacá for potato cultivation (*Solanum tuberosum* subsp. *andigena*) (ICA, 1992). In the town of Cogua, there was no response to application of P in the yield variables of the categories: first (PT1), second (PT2) and third (PT3) (Tab. 2), a result related to the amount of the element in the soil, which for the observed response is sufficient for the variety Criolla Colombia. The results suggest the need for future studies on critical levels of P and K for varieties of *Solanum phureja*, because the variety Criolla Colombia has shown, similar to the reports of Becerra *et al.* (2007) for the Guaneña variety, requirements well below those reported for tetraploid potato varieties, which favors the production cost structure for the cultivation of diploid varieties and therefore their competitiveness.

In the total yield (PTT) variable, differences were observed for the P factor, that is, the levels 50, 100 and 150 kg ha⁻¹ of P₂O₅ were equal and above the 0 kg ha⁻¹ P₂O₅ level, with values higher than 32 t ha⁻¹, well above the average production of six localities for this variety as reported by Rodríguez *et al.* (2009) in Antioquia (15.62 t ha⁻¹). As in the town of Zipaquirá, this variety responded to P application up to 50 kg ha⁻¹ P₂O₅, but did not differ above this level, which shows a low requirement of this element, or an efficient ability to absorb it, allowing one to gather that, taking into account that this variety is a clonal selection from the native yellow population, it is, no doubt, a natural strategy for survival not known in detail (Fig. 5).

The soil of this town presented a soil K level of 1.72 cmol kg⁻¹, a high level for the potato (ICA, 1992), this explains the lack of response to K application in the key yield categories, PT1 and PT2 (Tab. 2). Similar results were reported by Becerra *et al.* (2007), who found no differences in yield variables or specific gravity for phosphorus or K factors, nor for the interaction between them in 'Criolla Guaneña' grow at two locations in Pasto (Nariño).

The results of this work with respect to the variable specific gravity are similar to those reported by other research in potato fertilization, in which the absence of a variable

response: to the application of increasing levels of P has been noted in the variety Diacol Capiro (Núñez *et al.*, 2006), at different levels of K in soils of Saudi Arabia in the variety of the subspecies *tuberosum* (Abdelgadir *et al.*, 2003), at different levels and sources of K in the variety of Washington's Columbia Basin (Davenport and Bentley, 2001). Although the specific gravity did not respond to the application of different levels of K in the towns of Zipaquirá and Cogua for the variety Criolla Colombia, it is important to further studies to determine what level and source of K produces a reduced specific gravity, as reported by studies on *S. tuberosum* (McDole, 1978; Panique *et al.*, 1997; Kumar *et al.*, 2007).

Similarly, the results with respect to tuber yield are in line with those made by Dubetz and Bole (1975) who reported in experiments in Canada that K fertilization did not affect the number or weight of tubers in soils with high K, by Panique *et al.* (1997) who found no yield response to K application in six of 11 sites tested in Wisconsin, by Davenport and Bentley (2001) at different levels and sources of K in the variety Washington's Columbia Basin, and by Abdelgadir *et al.* (2003) in soils and varieties of Saudi Arabia.

Phosphorus fertilization research on the variety Diacol Capiro conducted in the municipalities of Villapinzón (Cundinamarca) and Toca and Soracá (Boyacá) showed a tuber yield response to the level of 200 kg ha⁻¹ of P₂O₅ (Núñez *et al.*, 2006.) The results of this research show that the variety Criolla Colombia (*Solanum phureja*) has a P requirement lower than that of the variety Diacol Capiro (*Solanum tuberosum*), bearing no response above the 50 kg ha⁻¹ P₂O₅ level, which can be explained by its lower ploidy, reduced cycle time and lower crop yield potential, resulting in lower consumption of this nutrient.

Conclusion

The results of this study showed that the variety Criolla Colombia responds to P application, and that yield does not increase with the implementation of levels above 50 kg ha⁻¹ P₂O₅. In the case of K, this variety did not respond to the application of different levels of this element, however, given the conditions of the locations where the tests were performed, where the level of this element was high, similar studies in soils with low contents of K are recommended, which would allow a better understanding of the fertilization needs of this variety that is highly important to Colombia.

Cited literature

- AbdelGadir, A., M. Errebhi, H. Al-Sarhan, and M. Ibrahim. 2003. The effect of different levels of additional potassium on yield and industrial qualities of potato (*Solanum tuberosum* L.) in an irrigated arid region. *Am. J. Pot. Res.* 80(3), 219-222.
- Alvarez-Sánchez, E., J. Etchevers, J. Ortíz, R. Núñez, V. Volke, L. Tijerina and A. Martínez. 1999. Biomass production and phosphorus accumulation of potato as affected by phosphorus nutrition. *J. Plant Nutr.* 22(1), 205-217.
- Barber, S. 1994. Root growth and nutrient uptake. pp. 95-99. In: Boote, K., J. Bennet, and M. Paulsen (eds.). *Physiology and determination of crop yield.* American Society of Agronomy, Madison, FL.
- Becerra, L.A., S.L. Navia, and C.E. Núñez. 2007. Efecto de niveles de fósforo y potasio sobre el rendimiento del cultivar 'Criolla Guaneña' en el departamento de Nariño. *Rev. Latinoam. Papa* 14(1), 51-60.
- Bustos, P., V. Arias, and C. Núñez. 1996. Evaluación del rendimiento en papa criolla (*Solanum phureja* Juz. et. Buk) variedad 'Yema de huevo', bajo diferentes densidades de siembra en la Sabana de Bogotá. *Agron. Colomb.* 13(2), 162-168.
- Carmona, G., H. Fano, M. Ordinola, and G. Scott. 1998. Experiencias de exportación de la papa amarilla peruana. International Potato Center (CIP); Asociación de Exportadores (ADEX), Lima.
- Chapman, K., L. Sparrow, P. Hardman, D. Wright, and J. Thorp. 1992. Potassium nutrition of Kennebec and Russet Burbank potatoes in Tasmania: Effect of soil and fertiliser potassium on yield, petiole and tuber potassium concentrations, and tuber quality. *Aust. J. Exp. Agr.* 32(4), 521-527.
- Davenport, J. and E. Bentley. 2001. Does potassium fertilizer form, source, and time of application influence potato yield and quality in the Columbia basin? *Am. J. Pot. Res.* 78(4), 311-318.
- Dubetz, S. and J. Bole. 1975. Effect of nitrogen, phosphorus, and potassium fertilizers on yield components and specific gravity of potatoes. *Am. J. Pot. Res.* 52(12), 399-405.
- Estrada, N. 2000. La biodiversidad en el mejoramiento genético de la papa. Fundación Promoción e Investigación de los Productos Andinos (PROINPA); International Potato Center (CIP); Centro de Información para el Desarrollo (CID), La Paz.
- Fedepapa, Federación Colombiana de Productores de Papa. 2003. Censo nacional de la papa departamento de Cundinamarca. *Rev. Papa* 23, 52.
- Hawkes, J.G. 1990. *The potato: evolution, biodiversity and genetic resources.* Belhaven Press, London.
- Huaman, Z. and D. Spooner. 2002. Reclassification of landrace populations of cultivated potatoes (*Solanum* sect, *Petota*). *Am. J. Bot.* 89(6), 947-965.
- ICA, Instituto Colombiano Agropecuario. 1992. Fertilización en diversos cultivos. Quinta aproximación. Manual de Asistencia Técnica No. 25. Bogotá.
- Kumar, P., S. Pandey, B. Singh, S. Singh, and D. Kumar. 2007. Influence of source and time of potassium application on potato growth, yield, economics and crisp quality. *Pot. Res.* 50(1), 1-13.
- Laboski, C. and K. Kelling. 2007. Influence of fertilizer management and soil fertility on tuber specific gravity: a review. *Am. J. Pot. Res.* 84(4), 283-290.
- López, L. 1993. Las papas silvestres de Colombia. *Rev. Papa* 7(1), 4-13.
- Luján, L. 1996. Historia de la papa. *Rev. Papa* 16(1), 4-25.
- Marschner, H. 2002. *Mineral nutrition of higher plants.* 2nd ed. Academic Press, Londres.
- McCollum, R. 1978. Analysis of potato growth under differing P regimes. I. Tuber yield and allocation of dry matter and P. *Agron. J.* 70(1), 51-57.
- McDole, R. 1978. Potassium fertilizer trials with potatoes on coarse-textured soils in southeastern Idaho. *Am. Pot. J.* 55(3), 161-170.
- Mengel, K. and E. Kirkby. 1987. *Principles of plant nutrition,* 4th ed. International Potash Institute. Bern.
- Mengel, K., E. Kirkby, H. Kosegarten, and T. Appel. 2001. *Principles of plant nutrition.* 5th ed. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Núñez, C.E., M. Santos, S.L. Navia, and J.M. Cotes. 2006. Evaluación de la fertilización fosfórica foliar y edáfica sobre el rendimiento de la variedad de papa 'Diacol Capiro' (*Solanum tuberosum* L.). *Agron. Colomb.* 24(1), 111-121.
- Panique, E., K. Kelling, E. Schulte, D. Hero, W. Stevenson, and V. James. 1997. Potassium rate and source effects on potato yield, quality, and disease interaction. *Amer J. Pot. Res.* 74(6), 379-398.
- Pérez, L., L.E. Rodríguez, and M.I. Gómez. 2008. Efecto del fraccionamiento de la fertilización con N, P, K y Mg y la aplicación de los micronutrientes B, Mn y Zn en el rendimiento y calidad de papa criolla (*Solanum phureja*) variedad Criolla Colombia. *Agron. Colomb.* 36(3), 477-486.
- Pessaraki, M. 2001. *Handbook of plant and crop physiology.* 2nd ed. Marcel Dekker, New York.
- Pumisacho, M. and S. Sherwood. 2002. *El cultivo de la papa en Ecuador.* Instituto Nacional Autónomo de Investigaciones Agropecuarias (INIAP); International Potato Center (CIP), Quito.
- Rivera, J.E., A. Herrera, and L.E. Rodríguez. 2006. Evaluación sensorial en productos procesados de papa criolla (*Solanum phureja*) y su importancia para el fitomejoramiento. *Fitotec. Colomb.* 6(2), 9-25.
- Rodríguez, D., L.E. Rodríguez, and C.E. Núñez. 2006. Heredabilidad y evaluación del contenido de proteínas totales de la colección de papa criolla (*Solanum phureja* juz et buk) de la Universidad Nacional de Colombia. In: *Memorias del XXII Congreso de la Asociación Latinoamericana de la Papa (ALAP).* Toluca, México.
- Rodríguez, L.E., C.E. Núñez, and N. Estrada. 2009. Criolla latina, Criolla Paisa y Criolla Colombia, nuevos cultivares de papa criolla para el departamento de Antioquia (Colombia). *Agron. Colomb.* 27(3), 287-301.
- Rosen, C. and P. Bierman. 2008. Potato yield and tuber set as affected by phosphorus fertilization. *Am. J. Pot. Res.* 85(2), 110-120.
- Salisbury, F.B. and C.W. Ross. 1994. *Fisiología vegetal.* 4th ed. Grupo Editorial Iberoamericano. México DF.
- SADC, Secretaría de Agricultura Departamento de Cundinamarca. 2008. Determinación de las zonas con ventajas competitivas para las principales cadenas agroproductivas mediante la implementación de un sistema experto basado en los Sistemas de Información Geográfica Agropecuario para el departamento de Cundinamarca. Convenio Interadministrativo

- No. 053. Gobernación de Cundinamarca; Universidad de Cundinamarca. En: <http://es.scribd.com/doc/32084576/43/Provincia-Sabana-Centro>; consulted: June, 2011.
- Spooner, D.M., J. Núñez, G. Trujillo, M. del R. Herrera, F. Guzmán, and M. Ghislain. 2007. Extensive simple sequence repeat genotyping of potato landraces supports a major reevaluation of their gene pool structure and classification. *PNAS* 104, 19398-19403.
- Talbert, W.F. and O. Smith. 1975. *Potato processing*. 3th ed. The AVI Publishing Company, Westport, CT.
- Vargas, A.M. 2010. Respuesta de la colección central colombiana de *Solanum phureja* de la Universidad Nacional de Colombia a la infección con *Potato yellow vein virus* (PYVV). Tesis de maestría. Facultad de Agronomía, Universidad Nacional de Colombia, Bogotá.
- Villarreal, H., P.D. Porras, A. Santa, J. Lagoeyte, and D. Muñoz. 2007. Costos de producción de papa en las principales zonas productoras de Colombia. In: http://papaunc.com/img_upload/f243f82ed8fde644849683ee48a08ef8/ESTUDIO_DE_COSTOS_DE_PROD._PAPA.pdf; consulted: June, 2011.