

Screening of microbial culture filtrates, plant extracts and fungicides for control of mango anthracnose

Screening de filtrados de cultivos microbianos, extractos vegetales y fungicidas para el control de la antracnosis del mango

Jairo A. Osorio¹, Erika P. Martínez¹, and Juan C. Hío¹

ABSTRACT

Anthrachnose (*Colletotrichum gloeosporioides*) induces losses of up to 40% in most mango growing regions in Colombia. A series of exploratory experiments were carried out to evaluate 14 potentially antagonistic microorganisms, six plant extracts and 10 fungicides for disease control based on their ability to inhibit spore germination, mycelium growth and development of symptoms on detached fruits. *In vitro* experiments showed that spore germination of *C. gloeosporioides* was strongly inhibited ($P \leq 0.0001$) by all microbial culture filtrates, all fungicides and two plant extracts, suggesting that some of these treatments may hold promise for disease control. Similarly, all microbial filtrates, most fungicides and two plant extracts caused a significant ($P \leq 0.0001$) inhibition in mycelium growth of the pathogen. Results from this study indicated small effects of fungicides or plant extracts on anthracnose development on inoculated fruits, while microbial filtrates induced a significant reduction ($P \leq 0.05$) in lesion size. These results are very promising since current control practices rely on fungicides or costly physical treatments.

Key words: *Colletotrichum gloeosporioides*, fruit protection, non-conventional control methods.

RESUMEN

La antracnosis (*Colletotrichum gloeosporioides*) induce pérdidas de hasta 40% en la mayoría de zonas productoras de mango en Colombia. Se realizó una serie de experimentos exploratorios para evaluar 14 antagonistas potenciales, seis extractos vegetales y 10 fungicidas en el control de la enfermedad, con base en su capacidad para inhibir la germinación de esporas, el crecimiento del micelio y el desarrollo de síntomas en frutos desprendidos. Los experimentos *in vitro* mostraron que la germinación de esporas de *C. gloeosporioides* fue inhibida ($P \leq 0.0001$) por todos los filtrados crudos microbianos, todos los fungicidas y dos de los extractos vegetales; indicando que algunos de estos tratamientos pueden ser promisorios para el control de la enfermedad. Asimismo, todos los filtrados microbianos, varios fungicidas y dos extractos vegetales causaron una inhibición significativa ($P \leq 0.0001$) en el crecimiento del micelio del patógeno. Los resultados de este estudio mostraron un bajo efecto de los fungicidas ó los extractos vegetales en el desarrollo de antracnosis en frutos inoculados, mientras que los filtrados microbianos mostraron una reducción significativa ($P \leq 0.05$) en el tamaño de lesión. Esto es muy promisorio pues las prácticas actuales de control se basan en fungicidas ó tratamientos físicos costosos.

Palabras clave: *Colletotrichum gloeosporioides*, protección de frutos, control no convencional.

Introduction

Several varieties of mango (*Mangifera indica*, L.) such as Haden, Irwin, Keitt and Tommy Atkins are grown in Colombia, as well as local varieties that are used in the industry (MADR, 2009). Mango orchards in all regions are affected by anthracnose (*Colletotrichum gloeosporioides* [Penz.] Penz. & Sacc., teleomorph: *Glomerella cingulata* [Stoneman] Spaulding & Von Schrenk); this disease induces field losses of up to 40% and postharvest losses of up to 70% in the most susceptible cultivars (Páez-Redondo, 1995; Afanador *et al.*, 2003; MADR, 2009) in Mexico, India, Australia and South Africa (Chowdhury *et al.*, 2007; Kumar

et al., 2007; Rojas-Martínez *et al.*, 2008). Typical symptoms of anthracnose are sunken, necrotic, dark brown to black lesions on mango fruit (Bailey *et al.*, 1992). The size and shape of these lesions vary with the cultivar, amount of quiescent infections and age of the infected organs (Arauz, 2000; Beno-Moualem and Prusky, 2000; Wharton and Diéguez-Uribeondo, 2004; Osorio, *et al.*, 2006).

Various anthracnose control methods are used by farmers including fungicides, calcium carbonate, antagonists and sanitation (Benítez *et al.*, 2004; Sundravadana *et al.*, 2006; Korsten, 2006; Korsten *et al.*, 1997; Stockwell and Stack, 2007). In Colombia, fungicide sprays and occasional

Received for publication: 23 May, 2011. Accepted for publication: 29 June, 2012.

¹ Plant Pathology Laboratory, Tibaitatá Research Center, Corporación Colombiana de Investigación Agropecuaria (Corpoica). Mosquera (Colombia). josorio@corpoica.org.co.

sanitation are used by mango growers once symptoms of the disease are visible, but the effectiveness of these measures is limited (Osorio *et al.*, 2000; Osorio *et al.*, 2006; Castro *et al.*, 2007; Moral *et al.*, 2009).

Fungicide tolerance in populations of *C. gloeosporioides* was reported more than 20 years ago (Liyanage *et al.*, 1992), and has resulted in reduced biological activity and increased fungicide residues on harvested fruits. This has prompted efforts to develop alternate methods for anthracnose control on various crops (Cheng *et al.*, 2006), including prediction models based on weather information (Dodd, 1991; Arauz, 2000). Methods used for post-harvest anthracnose control include fungicides, biocides, calcium carbonate, chemical inducers, microorganisms (Benítez *et al.*, 2004), hyperbaric pressure treatments, antioxidants, radiation, heat shock and, -at the experimental stage- genetic manipulation (Korsten, 2006).

Three recent studies carried out by Corpoica examined the survival strategy and population structure of the anthracnose pathogen (*C. gloeosporioides*) that affects mango orchards in Colombia (Ospina and Osorio, 2005; Cadavid *et al.*, 2007; Martínez *et al.*, 2009) and tamarillos (Reyes *et al.*, 2007); the results suggested the need to explore diverse alternative control methods for this disease. Here, we report the results from laboratory studies carried out to determine the biological activity of microbial antagonists, plant extracts and fungicides against *C. gloeosporioides*; focusing on the source of the inoculum and symptom suppression in mature fruits.

Materials and methods

Two sets of studies were conducted at the Plant Pathology Laboratory of Corpoica (Tibaitata Research Center, Mosquera, Colombia) to determine the ability of microbial culture filtrates (in 2007), plant extracts and commercial fungicides (in 2008) to inhibit spore germination and mycelium growth of *C. gloeosporioides* and prevent symptom development in fruits.

Pathogen isolates. Two single spore isolates (Cg-917 and Cg-935), obtained from “Hilacha” mango fruits harvested in Coello (Tolima, Colombia) and identified as *C. gloeosporioides* by morphology, growth on selective media and species-specific rDNA amplification profiles, were used in this study. The two isolates were similar in spore morphology, colony growth habit, pathogenicity and SSR (microsatellites) profiles, according to previous studies (Cadavid *et al.*, 2007). The two isolates also showed similar

germination and mycelium growth responses to fungicides, plant extracts and culture filtrates in exploratory trials.

Microorganisms and liquid culture filtrates. Isolates of potential antagonists were obtained from field samples (leaves, flowers, fruits) collected between October/2006 and March/2007 at two locations (Sevilla, Magdalena and Espinal/Guamo, Tolima). The plant material was processed for microbial isolation according to published protocols (Paterson and Bridge, 1994; Barnett *et al.*, 2000; Schaad, 2001). The fungal, yeast and bacterial colonies were purified and stored at -20°C in 20% glycerol stock solutions. Identification of fungal and bacterial isolates was made by morphological comparisons and standard biochemical tests, respectively.

Five liquid media were prepared to obtain crude culture filtrates: reelse and mandels medium for *Trichoderma*, Czapeck medium for *Verticillium* and *Paecilomyces* growth, TSA medium for bacteria, YEG medium for actinomycetes and NYBB medium for yeasts. Pure colonies of each isolate were used to inoculate 500 mL flasks containing 250 mL of a specific liquid media, before incubation in an orbital shaker, following specific protocols (Tab. 1) (Paterson and Bridge, 1994; Donadio *et al.*, 2002; Abril *et al.*, 2008). Following incubation, microbial biomass was separated by

TABLE 1. Liquid media and incubation conditions for microbial groups used in this study.

Microbial group	Liquid medium	Incubation time (d)	Incubation temperature (°C)
<i>Trichoderma</i> sp.	Reelse & Mandels	3	28
<i>Paecilomyces</i> sp.	Czapeck	12	28
<i>Verticillium</i> sp.	Czapeck	7	28
<i>Bacillus</i> sp., <i>Pseudomonas</i> sp.	TSA	5	28
Actinomycetes	YEG	3	37
Yeasts	NYDB	5	28

Filtration through cheesecloth, centrifugation (8000 g, 20 min) and filtration through millipore membrane filters (0.1 or 0.3 micrometers in diameter, according to the type of microorganism).

Plant extracts. Five commercially available plant extracts were obtained from local suppliers: CapsiAlil® (Sulfur compounds of Liliaceas+Protoalcaloides of Solanaceous), Eco-Swing® (terpenoids from *Swinglea glutinosa*), Desfan-100® (Citrus seed extract), Xplode SL (capsaicin+alisin), swinglea extract (Terpenoids). Additionally, an aqueous extract of the xeric lefaria cactus (*Cereus deficiens*) was prepared by grinding and drying tissue from the middle portion of fully

extended branches. A 20 g sample was fully homogenized in 400 mL of distilled water and filtered through cheesecloth before use. Four different concentrations of each extract were evaluated.

Fungicides. Ten commercial fungicides were used in this study: Score® 250EC (difenoconazole), Kocide® 101 (copper hydroxide), Alarm® 80WP (mancozeb), Amistar® 50WG (azoxystrobin), Mertec® 500SC (thiabendazole), Elosal® 720SC (sulphur), Carbendazim AC® (carbendazim), Kasumin® 2SL (kasugamycin), Validamycin (validamycin) and Rovral® 50WP (iprodione). Four concentrations of each fungicide were evaluated.

Spore germination assays. Exploratory tests were performed with isolates Cg-917 and Cg-935 in order to determine spore germination and standardize incubation conditions (Cerón *et al.*, 2005). Fresh colonies of the two isolates were produced on Marthur's medium incubated at 28°C in darkness for 10 d. Spore suspensions ($1 \cdot 10^5$ spores/mL) were prepared on either a) sterile water amended with Tween®-80 (0.05% v/v) or b) a 10% aqueous sucrose solution amended with Tween-80 (0.05% v/v); and incubated at 28°C and 100% relative humidity on glass slides. Germination counts were carried out every 2 h and for up to 24 h, germination was higher on water (10-15%) as compared to the sucrose solution (3%). Based on these results, all germination assays used 1.5 mL of spore suspensions of $1 \cdot 10^5$ spores/mL in water + Tween®-80 (0.05% v/v), using Eppendorf tubes incubated at 27°C for 18 h.

A completely randomized design with three repetitions was used to compare the inhibitory activity of crude filtrates from liquid cultures of 14 microbial isolates (six fungi, four bacteria, two actinomycetes, and two yeasts) on spore germination of the isolate Cg-917. For this, spore suspensions were mixed at a 1:1 ratio with the culture filtrates in 2 mL Eppendorf tubes, and incubated as described above. Following incubation, 50 µL were taken from each tube and poured into a Neubauer chamber to determine germination.

A randomized block design displayed in a factorial arrangement with three repetitions was used to compare six plant extracts in four concentrations (Tab. 2) by their inhibitory activity on spore germination of the isolate Cg-935 of the mango pathogen. Spore suspensions prepared as described were mixed at a 1:1 ratio with each one of the plant extract treatments, incubated and evaluated as described above.

To test the efficacy of ten fungicides, at different concentrations (Tab. 2), to inhibit spore germination of the

TABLE 2. Plant extracts and fungicide treatments used in this study.

Plant extract	Concentration (mg L ⁻¹)			
	D1	D2	D3	D4
CapsiAlil®	0.950	0.700	0.450	0.200
EcoSwing®	2.000	1.500	1.000	0.500
Desfan-100®	0.250	0.200	0.150	0.100
Xplode	5.000	4.000	3.000	2.000
Swinglea extract	20.000	15.000	10.000	5.000
Lefaria extract	2.000	1.000	0.500	0.250
Difenoconazole	0.375	0.250	0.125	0.075
Copper Hydroxide	1.500	1.000	0.500	0.250
Mancozeb	0.300	0.200	0.100	0.050
Azoxystrobin	0.450	0.300	0.150	0.075
Thiabendazole	0.675	0.450	0.225	0.115
Sulphur	1.440	0.720	0.360	0.072
Carbendazim	2.000	1.000	0.50	0.100
Kasugamycin	3.000	1.500	0.750	0.370
Validamycin	3.000	1.500	0.750	0.370
Iprodione	2.000	1.000	0.500	0.100

D2 is a reference concentration equivalent to the manufacturer's recommendation. D1 is a higher concentration, whereas D3 and D4 are lower concentrations.

C. gloeosporioides isolate Cg-935, a completely randomized design displayed in a factorial arrangement with four repetitions was used. Spore suspensions were mixed at a 1:1 ratio with each one of the fungicide treatments, incubated and evaluated as described above.

Mycelium growth assays. A completely randomized design with three repetitions was used in an experiment to test the effect of culture filtrates from 14 microbial isolates (six fungi, four bacteria, two actinomycetes, and two yeasts) on mycelium growth of the *C. gloeosporioides* isolate Cg-917. Petri plates were prepared by mixing 9 mL of PDA with 1 mL of culture filtrate and left to solidify, inoculated with 5-mm disks of PDA + mycelium of Cg-917 and incubated at 27°C in the dark for 7 d. After this period, the colony diameter was measured.

Mycelium growth of the isolate Cg-935 in the presence of plant extracts (Tab. 2) was determined in a randomized block experiment with a factorial arrangement and three repetitions. Petri plates were prepared by mixing 9 mL of PDA with 1 mL of plant extract at each concentration. Once solidified, 5-mm disks of PDA with actively growing Cg-935 mycelium were placed in the center of each plate. Inoculated plates were incubated at 27°C in the dark for 7 d, at which time the colony diameter was measured.

The effect of ten fungicides at different concentrations (Tab. 2) on mycelium growth of the *C. gloeosporioides* isolate Cg-935 was tested in a completely randomized experiment with a factorial arrangement and four repetitions.

Petri plates were prepared by mixing 9 mL of PDA with 1 mL of fungicide solution. Once solidified, 5-mm disks of PDA with actively growing mycelium of either isolate of *C. gloeosporioides* were placed in the center of each plate. Inoculated plates were incubated at 27°C in the darkness for 7 d, at which time the colony diameter was measured.

Fruit protection assays. The best treatments from all previous experiments were tested for anthracnose control on detached mango fruits. A completely randomized experiment replicated three times was used to evaluate the protection ability of 14 microbial filtrates on fruits inoculated with the isolate Cg-917 of the anthracnose pathogen. Similarly, a completely randomized experiment in a factorial arrangement replicated three times was used to test the effectiveness of five fungicides on fruits inoculated with the isolate Cg-935 of the mango pathogen. Likewise, a completely randomized factorial experiment was used to test the effectiveness of two plant extracts at two concentrations each at preventing anthracnose development on fruits inoculated with the isolate Cg-935.

Mango fruits (hilacha cultivar, E2 growth stage) were washed, superficially disinfected with 1% NaOCl, and rinsed in distilled water. Treatments of fruits with culture filtrates from antagonists, plant extracts or fungicide solutions consisted of a 30-min immersion, followed by air drying at room temperature. Treated fruits were then inoculated by placing two 5-mm disks of a sporulating colony of the pathogen on the fruit surface (5 cm apart). After inoculation, the fruits were kept inside plastic boxes (two fruits per box) containing moist paper towels, at 25°C in alternating 12/12 dark/light conditions. Lesion diameter, resulting from inoculation, was assessed after 8 d.

Data analysis. All statistical analyses were performed with SAS® version 9.3. Standard ANOVA and Tukey Studentized Range Test (HSD) for mean separation ($P \leq 0.05$) were performed on data from spore germination and mycelium growth experiments. Similarly, data from fruit protection assays were analyzed by ANOVA and treatment means were separated using HSD ($P \leq 0.05$).

Results

Biological activity of microbial culture filtrates. Four exploratory tests were initially carried out in 2007 with four microbial groups (bacteria, fungi, actinomycetes, yeasts), resulting in 14 isolates being selected for further testing against the pathogen *C. gloeosporioides* isolate Cg-917 (Tab. 3). Culture filtrates from all 14 isolates reduced spore germination ($P \leq 0.0001$) in the range of 43 to 69% as compared

to the control treatment; and some of them induced germ tube malformations (data not shown). The Tukey (HSD) test indicated greater effects for the isolates Actin-38, Bacte-15, Bacte-40, Bacte-59, Paeci-85 and Trich-19. Mycelium growth of the pathogen on filtrate-amended medium was reduced by all microbial isolates ($P \leq 0.0001$), with greater reductions (90- 95% colony growth reduction) induced by the isolates Actin-35, Bacte-59, Bacte-40, Bacte-73 and Yeast-05 (Tab. 3).

TABLE 3. Effects of culture filtrates of 14 microbial isolates on germination and mycelium growth of *C. gloeosporioides* (isolate Cg-917) and lesion size on inoculated mango fruits.

Isolate	Spore germination (%) ¹	Colony diameter (cm) ²	Lesion diameter (cm) ³
Control	8.16 a	8.73 a	1.27 a
Bacte-73	4.66 b	0.63 c	0.15 c
Yeast-08	4.10 b	0.66 c	0.32 b
Yeast-05	3.76 bc	0.60 c	0.30 b
Trich-01	3.73 c	1.66 b	0.17 c
Actin-35	3.60 c	0.53 d	0.27 b
Verti-79	3.46 c	1.70 b	0.27 b
Verti-77	3.40 c	0.93 b	0.70 a
Paecil-83	3.23 cd	0.86 c	0.22 bc
Trich-19	3.13 cd	1.26 b	0.12 c
Paecil-85	2.93 d	0.90 bc	0.12 c
Bacte-59	2.86 d	0.56 c	0.25 bc
Bacte-40	2.70 d	0.56 c	0.22 bc
Bacte-15	2.60 d	0.73 c	0.20 bc
Actin-38	2.56 d	1.23 b	0.12 c
cv	14.04	18.55	60.01
F	** , $P \leq 0.0001$	** , $P \leq 0.0001$	* , $P \leq 0.05$

¹ Spore germination counts were performed after 18 h of incubation. A total of 100 spores were observed in each replicate.

² Diameter of fungal colonies was determined after incubation for 7 d.

³ Inoculated fruits were incubated for 8 d before measurements of lesion diameter were taken.

Mango fruits that were superficially sprayed with culture filtrates of five isolates (Paeci-85, Trich-19, Trich-1, Bacte-73 and Actin-38) before inoculation with Cg-917, showed lesion size reductions around 80% when compared with the untreated control ($P \leq 0.05$); however, caution should be exercised when interpreting these results due to the high variation coefficient in this experiment. This suggests the need for additional studies with a larger number of fruits to take into consideration factors like the presence of quiescent infections on the surface and the nutritional status of fruit tissues, which lead to intrinsic variability among mango fruits of the same developmental stage E-2.

Biological activity of fungicides. Ten fungicides were initially tested on spore germination and mycelium growth assays, and five were selected for fruit protection assays; the data from these experiments are presented as general

treatment means in Tab. 4. The variation coefficient for the spore germination experiment was moderate to high (37.6%), suggesting that factors related to fungicide solubility in water or conditions for optimal contact or uptake by the fungal spores were not adequately controlled in the preparation of the spore germination assays. Dose effects of three fungicides (copper hydroxide, kasugamycin, validamycin) on colony growth were statistically significant, while two fungicides (azoxystrobin and thiabendazole) had a significant dose effect on spore germination.

Eight fungicides reduced ($P \leq 0.0001$) spore germination by 80-100%, with two of them (carbendazim and iprodione) causing total inhibition (Tab. 4). Mycelium growth of the pathogen was significantly inhibited by all tested fungicides ($P \leq 0.0001$), but the strongest inhibition presented in a culture medium amended with carbendazim or difenoconazole. Based on these results, five fungicides (azoxystrobin, carbendazim, thiabendazole, difenoconazole and validamycin) were used in fruit protection assays where significant ($P \leq 0.05$) but moderate effects were observed for three fungicides (carbendazim, azoxystrobin and thiabendazole), with lesion size reductions around 50% in relation to the unprotected control.

TABLE 4. Effects of fungicides on germination and mycelium growth of *C. gloeosporioides* (isolate Cg-935) and lesion size on inoculated mango fruits.

Fungicide	Spore germination (%) ¹	Colony diameter (cm) ²	Lesion diameter (cm) ³
Control	4.75 a	7.96 a	2.14 a
Kasugamycin	1.31 b	4.95 b	-
Sulphur	1.00 b	5.74 b	-
Validamycin	0.95 b	3.88 c	2.38 a
Thiabendazole	0.68 bc	0.58 e	1.30 b
Difenoconazole	0.68 bc	0.05 f	1.68 b
Copper hydroxide	0.62 c	1.05 d	-
Azoxystrobin	0.51 c	0.99 de	1.47 b
Mancozeb	0.37 d	1.68 d	-
Iprodione	0 e	3.58 c	-
Carbendazim	0 e	0 f	1.17 b
cv	37.44	14.47	31.10
F	** , $P \leq 0.0001$	** , $P \leq 0.0001$	* , $P \leq 0.05$

¹ Spore germination counts were performed after 18 h of incubation. A total of 100 spores were observed in each replicate.

² Diameter of fungal colonies was determined after incubation for 7 d.

³ Inoculated fruits were incubated for 8 d before measurements of lesion diameter were taken.

Biological activity of plant extracts. Tab. 5 shows the effect of six plant extracts on germination, colony growth and fruit infection by the *C. gloeosporioides* isolate Cg-935. Most plant extracts tested did not produce a strong effect on the pathogen. Among the plant extracts, Desfan and

Ecoswing had significant dose effects on mycelium growth of the pathogen, with higher effects at the highest dose as detected in the colony growth tests. However, this effect was not evident on spore germination or fruit infection assays. Therefore, data from plant extract experiments are presented as general treatment means of plant extracts.

Reduction of spore germination was statistically significant ($P \leq 0.0001$) with one treatment (Desfan), inducing 98% inhibition of germination; this plant extract also caused a significant ($P \leq 0.05$) reduction (88%) in colony growth. When applied on the mango fruit surface however, none of the extracts significantly reduced lesion size (Tab. 5).

Discussion

The effects of *C. gloeosporioides* inoculum on mango include: immediate lesion development following infection of the flowers, establishment of quiescent infections on immature fruits, or causation of anthracnose lesions following new infections of physiologically mature fruits (Arauz, 2000). Since heavy losses result from fruit anthracnose, we focused on features of the pathogen important for the onset of infection and development of fruit symptoms. Various studies describing the lifecycles of anthracnose pathogens (Bailey *et al.*, 1992) suggest that spore germination needs to be blocked in order to prevent quiescent infections and reduce fruit disease. Other studies (Benito *et al.*, 2000; Osorio *et al.*, 2006; Moral *et al.*, 2009) indicate that flower infection as well as quiescent infections on foliage could be important sources of inoculum for fruit infection.

Protecting mango fruits from anthracnose development is a common practice which relies mostly

On fungicide spray programs or physical treatment following harvest (Dodd, 1991; Chowdhury *et al.*, 2007); consequently, finding alternative treatments is desirable but little progress has been made. Fruit anthracnose development in mango may be influenced by a complex array of conditions that determine the effects of the inoculum and the infection of host tissues (Wharton and Diéguez-Urbeondo, 2004). The results from this study are preliminary and represent only a small set of conditions related to the development of this disease; promising treatments identified in this study should be tested for effectiveness on anthracnose control under a wider set of conditions and a greater number of pathogen isolates, representing the diversity and complexity of the *C. gloeosporioides* population associated with mango.

TABLE 5. Effects of plant extracts on germination and mycelium growth of *C. gloeosporioides* (isolate Cg-935) and lesion size on inoculated mango fruits.

Plant extract	Spore germination (%) ¹	Colony diameter (cm) ²	Lesion diameter (cm) ³
Control	2.50 a	3.30 a	2.25
Xplode	2.18 a	1.18 b	-
Lefaria extract	1.68 b	1.09 bc	-
CapsiAlil®	1.46 b	1.30 b	-
EcoSwing®	1.18 b	0.68 c	2.01
Swinglea extract	0.62 c	1.15 b	-
Desfan-100®	0.06 d	0.38 d	2.13
CV	18.60	7.71	61.40
F	** <i>, P</i> ≤0.0001	* <i>, P</i> ≤0.05	

¹ Spore germination counts were performed after 18 h of incubation. A total of 100 spores were observed in each replicate.

² Diameter of fungal colonies was determined after incubation for 7 d.

³ Inoculated fruits were incubated for 8 d before measurements of lesion diameter were taken.

Separate experiments with microbial filtrates, fungicides and plant extracts were conducted, whereby spore germination was assessed after 18 h of incubation. Spore germination is affected by factors like age, temperature and light or chemical composition of the medium (Abril *et al.*, 2008); which could explain the variation in germination levels observed among the experiments. In this study, overall spore germination levels were low, suggesting that the fungus may have unidentified internal constraints or regulatory mechanisms for survival similar to spore dormancy, as described for several plant pathogens (De Wolf and Isard, 2007). Alternatively, *C. gloeosporioides* spores may lose viability very rapidly after removal from their source, but data were not taken to support this hypothesis. Spore germination of *C. gloeosporioides* was strongly inhibited by microbial culture filtrates, fungicides and two plant extracts, suggesting that some of these treatments may hold promise for disease control. These should be tested more thoroughly, particularly microbial isolates since they also caused a highly significant inhibition of mycelium growth of the pathogen originating from the mango tissue surfaces and therefore, could have the ability to adapt and colonize host tissues. In addition, more experiments with fungicides and plant extracts are necessary to account for factors related to product solubility in water and optimize exposure or uptake by the fungal spores.

Colony growth of *C. gloeosporioides* on culture medium was stable across the experiments and appears to be a reliable trait to study sensitivity of the pathogen to various products. Mycelium growth in natural settings could be targeted for control at two specific stages: after spore germination but before penetration or during activation

of quiescent infections on the fruit surface. Postharvest treatments for anthracnose control focus on this later stage of the interaction (Arauz, 2000; Arias and Carrizales, 2007). All microbial filtrates, most fungicides and two plant extracts (Tabs. 3, 4, 5) caused a strong inhibition in mycelium growth of the isolates Cg-917 and Cg-935 of the pathogen; therefore, it would be of interest to evaluate some of these products for their ability to specifically inactivate quiescent infections of *C. gloeosporioides* on mango fruits. Mycelium growth, and its associated enzymatic activity (Huckelhoven, 2007), may be responsible for flower damage by *C. gloeosporioides* in mango; therefore, flower protection by some of the products tested in this study is worth trying, particularly with microorganisms which may be able to colonize developing reproductive structures.

The lesion size of E-2 mango fruits inoculated with sporulating colonies of the isolates Cg-917 or Cg-935 showed high variation coefficients (31-61%) across the experiments; some intrinsic factors of the used plant material or differences in the inoculum load could explain this variability in the results. Mango fruits at the E-2 growth stage were used for inoculations based on previous studies; however there is some degree of uncertainty about the age of fruits of similar size and shape; this could explain why the reaction to the inoculation was not uniform. Results from this study indicated only small effects of fungicides or plant extracts on anthracnose development in inoculated fruits, while microbial filtrates showed significant reduction in lesion size. This is very promising since current control practices rely on fungicides or costly physical treatments (Kumar *et al.*, 2007; Sundravada *et al.*, 2007).

Anthracnose development from late infections of mature fruits and from quiescent infections of immature fruits may proceed at different rates. These differences may be due to: a) the metabolic stage of the inoculum, b) the amount of active propagules at the infection site on the fruit surface and c) the time taken for initial colonization or lesion progress (Moral *et al.*, 2009). All these factors have to be taken into consideration in future studies aiming to test potential control products with detached mango fruits, since natural disease development takes place not only from recent infection events but also from quiescent infections occurring in early stages of fruit growth. Furthermore, how much spore load migrates to the developing mango fruit and from what sources is a question that needs to be answered in order to understand some of the conditions for disease onset on the fruit and design or adjust control practices based on chemical or biological products.

Conclusions

Spore germination and mycelium growth of *C. gloeosporioides* can be inhibited or reduced *in vitro* by fungicides, plant extracts and microbial liquid culture filtrates; these types of experiments can be used to identify substances with high biological activity against the pathogen.

Development of anthracnose lesions on inoculated mango fruits was strongly reduced by filtrates from the liquid culture of various microorganisms, suggesting that some antagonists may be potential biocontrol agents of *C. gloeosporioides* and should be tested further in the field and for postharvest conditions.

Acknowledgements

The authors thank Mr. Jorge Arguelles for helping with the data analysis, Mr. Jaime Aguirre for his work in the fungicide experiments and Mr. José Dimas Segura for collecting field samples for microbial isolation.

Literature cited

- Abril, M., K.J. Curry, B.J. Smith, and D.E. Wedge. 2008. Improved microassays used to test natural product-based and conventional fungicides on plant pathogenic fungi. *Plant Dis.* 92, 106-112.
- Afanador, L., D. Minz, M. Maymon, and S. Freeman. 2003. Characterization of *Colletotrichum* isolates from tamarillo, passiflora, and mango in Colombia and identification of a unique species from the genus. *Phytopathology* 93, 579-587.
- Arauz, L.F. 2000. Mango anthracnose: economic impact and current options for integrated management. *Plant Dis.* 84, 600-611.
- Arias, B. and L. Carrizales. 2007. Control químico de la antracnosis del mango (*Mangifera indica* L.) en pre y postcosecha en el municipio de Cedeño, estado Monagas, Venezuela. *Bioagro* 19(1), 1-8.
- Bailey, J.A., R.J. O'Connell, R.J. Pring, and C. Nash. 1992. Infection strategies of *Colletotrichum* species. pp. 88-120. In: Bailey, J.A. and M.J. Jeger (eds.). *Colletotrichum: Biology, Pathology and control*. CAB, Wallingford, UK.
- Barnett, J.A., R.W. Payne, and D. Yarrow. 2000. *Yeasts: characteristics and Identification*, 3rd ed. Cambridge University Press, Cambridge, UK.
- Benítez, T., A. Rincón, M. Limón, and A.C. Codón. 2004. Biocontrol mechanisms of *Trichoderma* strains. *Intl. Microbiol.* 7, 249-260.
- Benito, E.P., M. Arranz, and A. Eslava. 2000. Factores de patogenicidad de *Botrytis cinerea*. *Rev. Iberoam. Micol.* 17, s43-s46.
- Beno-Moualem, D. and D. Prusky. 2000. Early events during quiescent infection development by *C. gloeosporioides* in unripe avocado fruits. *Phytopathology* 90(5), 553-559.
- Cadavid, M., S. Kelemu, and J.A. Osorio. 2007. Genetic diversity in the anthracnose pathogen infecting tropical fruits in Colombia. *Phytopathology* 97(7), S-16.
- Castro, A., J. Romero, J.A. Osorio, E. Martínez, and J. Hio. 2007. Distribución temporal y espacial de fuentes de inóculo de *Colletotrichum acutatum* en el follaje sano de tomate de árbol. *Fitopatol. Colomb.* 31(1)1-5.
- Cerón, L., M. Higuera, M. Sánchez, S. Bustamante, and G. Buitrago. 2005. Crecimiento y desarrollo de *Colletotrichum gloeosporioides* f. *alatae* durante su cultivo en medios líquidos. *Acta Biol. Colomb.* 11(1), 99-109.
- Cheng, W., H. Ishii, and K. Nishimura. 2006. Fungicide sensitivity and phylogenetic relationships of anthracnose fungi isolated from various fruits in Japan. *Plant Dis.* 90, 506-512.
- Chowdhury, M., M. Rahim, K. Khalequzzaman, M. Humayan, and M. Alam. 2007. Effect of plant extracts and time of application on incidence of anthracnose, yield and quality of mango. *Intl. J. Sustain. Crop Prod.* 2(5), 59-68.
- De Wolf, E.D. and S.A. Isard. 2007. Disease cycle approach to plant disease prediction. *Annu. Rev. Phytopathol.* 45, 203-220.
- Dodd, J.C. 1991. Pre and postharvest control of mango anthracnose in the Philippines. *Plant Pathol.* 40, 576-583.
- Donadio, S., P. Monciardini, R. Alduina, and P. Mazza. 2002. Microbial technologies for the discovery of novel bioactive metabolites. *J. Biotechnol.* 99, 187-198.
- Huckelhoven, R. 2007. Cell wall-associated mechanisms of disease resistance and susceptibility. *Annu. Rev. Phytopathol.* 45, 101-127.
- Korsten, L. 2006. Advances in control of postharvest diseases in tropical fresh produce. *Intl. J. Postharv. Technol. Innov.* 1, 48-61.
- Korsten, L., E. De Villiers, R.C. Wehner, and J.M. Kotzetz. 1997. Field spray of *Bacillus subtilis* and fungicides for control of pre-harvest fruit disease of avocado in South Africa. *Plant Dis.* 81, 455.
- Kumar, A., N. Reddy, K. Reddy, and M. Devi. 2007. Evaluation of fungicidal resistance among *Colletotrichum gloeosporioides* isolates causing mango anthracnose in Agri Export Zone of Andhra Pradesh, India. *Plant Pathol. Bull.* 16, 157-160.
- Liyanage, H., M. Millan, and H. Kistler. 1992. Two genetically distinct populations of *Colletotrichum gloeosporioides* from citrus. *Phytopathol.* 82(11), 1371-1376.
- MADR, Ministerio de Agricultura y Desarrollo Rural. 2009. Anuario estadístico de frutales y hortalizas 2004-2008 y sus calendarios de siembras y cosechas. Bogotá.
- Martínez, E., J. Hio, J.A. Osorio, and M.F. Torres. 2009. Identification of *Colletotrichum* species causing anthracnose on Tahiti lime, tree tomato and mango. *Agronomía Colombiana* 27 (2): 211-218.
- Moral, J., R. Oliveira, and A. Trapero. 2009. Elucidation of the disease cycle of olive anthracnose caused by *Colletotrichum acutatum*. *Phytopathology* 99, 548-556.
- Osorio, J.A., E. Viáfara, and N. Murcia. 2000. Aspectos epidemiológicos y de manejo de la antracnosis de los cítricos. pp. 41-48. In: *Memorias del Seminario Nacional sobre Enfermedades de los Cítricos*. Pereira, Colombia.
- Osorio, J.A., M.F. Torres, J.C. Hio. 2006. Detección y análisis de la distribución de infecciones latentes de *Colletotrichum*



- acutatum* J. H. Simmonds en árboles de frutales tropicales. Fitopatol. Colomb. 29(2), 45-52.
- Ospina, C.A. and J.A. Osorio. 2005. Caracterización de poblaciones del género *Colletotrichum* causante de la antracnosis de los cítricos. Fitopatol. Colomb. 29(1), 15-22.
- Páez-Redondo, A. 1995. Uso de variedades tolerantes: alternativa para el Manejo de antracnosis (*Colletotrichum gloeosporioides*) en mango (*Mangifera indica*). ASCOLFI-Informa 21, 36-39.
- Paterson, R. and P.D. Bridge. 1994. Biochemical techniques for filamentous fungi. IMI Technical Handbooks No. 1. CAB, Wallingford, UK.
- Reyes, A., J.A. Osorio, E. Martínez, and J. Hio. 2007. Estudio de la relación fenológica reproductiva y el comportamiento de las infecciones latentes de *Colletotrichum acutatum* en tomate de árbol. Fitopatol. Colomb. 31(2), 37-42.
- Rojas-Martínez, R., E. Zavaleta, D. Nieto, and M. Acosta. 2008. Virulence and genetic variation of *Colletotrichum gloeosporioides* (Penz.) Penz. and Sacc. on mango (*Mangifera indica* L.) cv. Haden. Rev. Mex. Fitopatol. 26, 21-26.
- Schaad, N.W. 2001 Laboratory guide for identification of plant pathogenic bacteria. Department of Plant Pathology, University of Georgia, St. Paul, MN.
- Stockwell, V. and J. Stack. 2007. Using *Pseudomonas* spp for integrated bio-control. Phytopathology 97, 244-249.
- Sundravadana, S., D. Alice, S. Kuttalam, and R. Samiyappan. 2006. Control of mango anthracnose by azoxystrobin. Tunis. J. Plant Protection 1, 109-114.
- Sundravadana, S., D. Alice, S. Kuttalam, and R. Samiyappan. 2007. Efficacy of azoxystrobin on *Colletotrichum gloeosporioides* Penz growth and on controlling Mango anthracnose. J. Agr. Biol. Sci. 2(3), 10-15.
- Wharton, P.S. and J. Diéguez-Urbeondo. 2004. The biology of *Colletotrichum acutatum*. Anales del Jardín Botánico de Madrid 61, 3-22.

