

Development of a new maize hybrid for the Ecuadorian lowland

Desarrollo de un nuevo híbrido de maíz para el Litoral Ecuatoriano

Ricardo Limongi-Andrade¹, Daniel Alarcón-Cobeña¹, Eddie Zambrano-Zambrano¹, Marlon Caicedo², Paúl Villavicencio-Linza², José Eguez³, Bernardo Navarrete⁴, Carlos Yanez⁵, and José L. Zambrano^{5*}

ABSTRACT

Yellow maize (*Zea mays* L. var. *indurata*) is mainly produced in the Ecuadorian lowlands (less than 1,200 m a.s.l.), primarily for feed production. Although Ecuador has recently increased maize production, new genotypes are needed for self-sufficiency and in order to avoid costly imports. Maize is sown mostly from December to January during the rainy season. However, the irregularity of rainfall has become a constraint on production. "INIAP H-603 Superior" is a new single-cross maize hybrid developed for the Ecuadorian lowlands with improved yields, which could contribute to domestic food security. The new hybrid had an average yield of 8.48 t ha⁻¹ with outstanding performance under the rainfall, sustained moisture, and irrigation conditions, outperforming the commercial hybrids INIAP H-602, DEKALB-7088 and INIAP H-553. Additionally, the new hybrid showed tolerance to the principal foliar diseases: leaf blight (*Exserohilum turicum*), rust (*Puccinia sorghi*), and leaf spot (*Curvularia lunata*), as well as good adaptability and stability, with a regression coefficient (*bi*) of 0.98 when it was evaluated in 29 locations from 2010 to 2013. The greatest yield potential of the new hybrid (10.82 t ha⁻¹) was obtained with irrigation during the dry season.

Key words: breeding, food security, climate change, stability.

RESUMEN

El maíz amarillo duro (*Zea mays* L. var. *indurata*) es producido principalmente en las zonas bajas o costeras del Ecuador (menos de 1200 m s.n.m.) y está destinado mayoritariamente para la elaboración de alimento balanceado. A pesar de que la producción de este cereal en el Ecuador se ha incrementado en los últimos años, se requieren de nuevos genotipos que permitan la autosuficiencia y eviten importaciones de maíz. El cultivo de maíz es realizado principalmente en la época de lluvia, cuando las irregulares precipitaciones se convierten en las principales limitantes para la productividad del cultivo. "INIAP H-603 Superior" es un híbrido simple de maíz desarrollado para expresar su potencial bajo condiciones de riego, lluvia o humedad remanente. El nuevo híbrido obtuvo rendimientos promedios de 8.48 t ha⁻¹, superando a los híbridos comerciales INIAP H-602, DEKALB-7088 e INIAP H-553 y mostró tolerancia a las principales enfermedades que afectaron al cultivo: lancha (*Exserohilum turicum*), roya (*Puccinia sorghi*), y mancha por curvularia (*Curvularia lunata*); así como buena adaptabilidad y estabilidad de rendimiento, con un coeficiente de regresión (*bi*) de 0.98, basado en pruebas multi-ambiente en 29 localidades, realizadas del 2010 al 2013. El mayor potencial de rendimiento del nuevo híbrido (10.82 t ha⁻¹) se obtuvo en la época seca bajo riego.

Palabras clave: fitomejoramiento, seguridad alimentaria, cambio climático, estabilidad.

Introduction

Flint yellow maize is a crop of significant importance in Ecuador. During the rainy season of 2016, farmers sowed roughly 192,000 ha of flint yellow maize, with an average yield of 5.5 t ha⁻¹ (Castro, 2016). Maize in Ecuador is a strategic crop for food security because of its multiple products and by-products, which are mainly used in the poultry, fish, and dairy feed industries.

Although this kind of maize production has increased in recent years in Ecuador, going from 540,000 t in 2002 to 1,730,000 t in 2015 (MAG, 2015), more high yielding hybrids that are adapted to the growing conditions in Ecuador are needed. Most of the maize (84% of the production area) is sown in December and January at the beginning of the rainy season (Aguilar *et al.*, 2015). The rainy season lasts three to five months, and the rainfall is irregularly distributed during this period (MAG, 2015).

Received for publication: 10 November, 2017. Accepted for publication: 28 July, 2018

Doi: 10.15446/agron.colomb.v36n2.68782

¹ National Maize Breeding Program, Instituto Nacional de Investigaciones Agropecuarias (INIAP). Portoviejo (Ecuador).

² National Maize Breeding Program, INIAP. Quevedo (Ecuador).

³ National Maize Breeding Program, INIAP. Gualaceo (Ecuador).

⁴ National Department of Plant Protection, INIAP. Portoviejo (Ecuador).

⁵ National Maize Breeding Program, INIAP. Quito (Ecuador).

* Corresponding author: jose.zambrano@iniap.gob.ec



Accordingly, the objective of the National Maize Breeding Program of INIAP is to develop and release high yielding hybrids that are adapted to the local growing conditions, with resistance to the principal foliar diseases of this crop.

Materials and methods

The single-cross maize hybrid “INIAP H-603 Superior” was obtained in 2010, crossing selected lines generated by the Maize Breeding Program (MBP) of the National Institute for Agricultural Research (INIAP) and lines introduced by the International Center for Maize and Wheat Improvement (CIMMYT), Colombia (Alarcón *et al.*, 2016).

Origin of parental lines

The female line POB.3.F4. 27-1-1-1 was developed by the MBP at the Experimental Station Portoviejo (EEP). It was derived from the Population POB.3 in 1994 using diallelic crossing between six genotypes: INIAP-541, INIAP-542, INIAP-526, Pool 26 *selección sequía* (CIMMYT), Pool 18 *selección sequía precoz* (CIMMYT), and Pacific 9205 (private company). After four cycles of mass selection, 256 inbred lines (S1) were developed with self-pollination. The lines were advanced to S4 and evaluated for adaptability and combining ability. This line was selected because of its good agronomic traits, combining ability and high yield. The male line CML-451 was introduced in 2007 from CIMMYT, Colombia. These lines are maintained at EEP by hand (plant to plant pollination), with controlled open pollination.

Experimental design and statistical analysis

This experiment was conducted in the provinces of Manabi (Santa Ana, Tosagua, Jipijapa), Guayas (Balzar, El Empalme), Los Ríos (Mocahe, Valencia) and Loja (Zapotillo, Pindal, Paltas). Ten experimental hybrids, including “INIAP H-603 Superior”, and five commercial checks (INIAP H-553, INIAP H-601, INIAP H-602, DE-KALB-1596, and DEKALB-7088) were evaluated using a randomized complete block design with three replicates in 29 locations (trials) from 2010 to 2013. To establish the differences between the hybrids, an analysis of variance and significance tests were carried out per location and combined (McIntosh, 1983; González *et al.*, 2010; Moore and Dixon, 2015) with the statistical package INFOSTAT v. 2015 (Di Rienzo *et al.*, 2015). A Tukey t-test ($P<0.05$) was used to determine the differences in the mean yield between the hybrids, growing conditions and interactions. The model used to estimate yield was: $Y_{ij} = \mu + G_i + A_j + (GA)_{ij} + B_k(j) + E_{ijk}$, where: Y_{ij} = average performance of the i-th hybrid obtained in the j-th environment; μ = effect of the

general median; G_i = effect of the i-th hybrid; A_j = effect of the j-th environment; $(GA)_{ij}$ = effect of the interaction between the i-th genotype and the j-th environment; $B_k(j)$ = effect of the k-th repetition on the j-th environment, and E_{ijk} = randomized effect of the experiment error associated with the i-th genotype on the j-th environment and k-th repetition.

The stability of the hybrids was estimated based on Eberhart and Russel’s (1966) criteria (Lin *et al.*, 1986). To evaluate the stability, the locations were grouped based on the three main maize Ecuadorian growing conditions: rainfall (20 trials), with irrigation during the dry season (four trials), and with water deficits (sustained moisture) during the dry season (five trials). The planting for the dry season occurred in June and July and, for the rainy season, in December and January. The experiment plots consisted of two rows (5 m long) with a distance between rows of 0.80 m, and 0.20 m between plants, for a population density of 62,500 plants ha^{-1} .

Agronomic management

The experiments were planted by hand. The fertilization and phytosanitary control of the crop were carried out according to Villavicencio and Zambrano (2009). The response of the hybrid to the prevalent foliar diseases: leaf blight (*Exserohilum turcicum*), rust (*Puccinia sorghi*), and leaf spot (*Curvularia lunata*) was evaluated under natural infection conditions, along with the five commercial hybrids used as controls at the EEP, from 2010 to 2013. The disease symptoms were scored after silking (reproductive stage R4), based on a 1 to 5 scale developed by CIMMYT (1995), where 1 was no symptoms and 5 was complete infection. The plant and ear (cob) heights, male and female flowering time, and grain yield were also evaluated as described by CIMMYT (1995), while the corncob type and grain texture were scored according to IBPGR (1991).

Results and discussion

Agronomic characteristics

The plant architecture and phenology of “INIAP H-603 Superior” are similar to other maize hybrids developed by INIAP (Tab. 1). The index between the ear (cob) and the plant height (AM/AP) and male and female flowering times are important traits for breeders because they contribute to lodging resistance and complete grain filling of the cob. The index AM/AP for INIAP H-603 Superior (0.5) and flowering time (55 and 57 d) were good values for lodging resistance and grain filling. These values agree with those reported by Gordon and Deras (2017) for maize

in Mesoamerica (AM/AP = 0.5, female flowering = 54 d, male flowering = 57 d), as well as the AM/AP index value reported by Vera *et al.* (2013), who evaluated eight hybrids in four locations in the central area of the Ecuadorian lowlands. Both authors observed variations caused by environmental effects, as seen in the present study.

TABLE 1. Morphological characteristics of the maize hybrid “INIAP H-603 Superior” developed for the lowlands of Ecuador.

Trait	Average and standard deviation	Range
Plant height	259 ± 26.8 cm	180-292
Ear height	127 ± 20.8 cm	75-155
Male flowering	55 ± 3.3 d	49-59
Female flowering	57 ± 2.8 d	51-62
Harvest time	120 d	118-135
Corn cob type	Conic-cylindrical	
Grain texture	Flint	

Resistance to major foliar diseases

“INIAP H-603 Superior” was rated 3 (moderate) for leaf blight (*E. turicum*), rated 1 (no symptoms) for rust (*P. sorghi*), and rated 2 (mild) for leaf spot (*C. lunata*) as shown in Table 2. These are significant foliar diseases in the provinces of Manabi, Guayas and Los Ríos (Villavicencio and Zambrano, 2009). The new hybrid showed equal or a better response (lower values) to foliar disease incidence

than the other commercial hybrids. *P. sorghi* (1 to 8%) and *E. turicum* (1 to 30%) were reported for Pergamino, Argentina by Parisi *et al.* (2014), who concluded that maize genotypes responded differently to these epiphytes. Vera *et al.* (2013) reported an average disease incidence of 3.31 and 2.44% for leaf spot and leaf blight, respectively, with an incidence not greater than 20% for maize hybrids evaluated in the Ecuadorian lowlands.

Yield performance

The combined analysis of variance for yield showed highly significant differences between the hybrids, environments and interactions (Tab. 3 and 4). A combined mean analysis showed that there were four ranges of significance for the hybrids, where “INIAP H-603 Superior”, with 8.48 t ha⁻¹ was the most stable (*bi* closest to 1) and shared the same range of significance for yield with DEKALB-1596 and INIAP H-601, with 8.20 t ha⁻¹ and 7.95 t ha⁻¹, respectively.

There were significant differences between the yield obtained with irrigation during the dry season (9.93 t ha⁻¹) and the yields obtained during the rainy season (6.68 t ha⁻¹) and the dry season with residual humidity (6.65 t ha⁻¹) (Tab. 4). This suggests that, during the rainy season, this crop may not receive the amount of water required by the plant to express its yield potential at the appropriate time. These results support the concept that high maize yields

TABLE 2. Foliar disease ratings for “INIAP H-603 Superior” and five commercial maize hybrids evaluated at the Experimental Station Portoviejo, 2011-2013.

Disease*	INIAP H-603	INIAP H-553	INIAP H-601	INIAP H-602	DEKALB1596	DEKALB 7088
Leaf blight (<i>Exserohilum turicum</i>)	3	3	3	3	4	4
Rust (<i>Puccinia sorghi</i>)	1	2	2	2	2	2
Leaf spot (<i>Curvularia lunata</i>)	2	3	3	3	4	4

* 1 to 5 scale, where 1 = no symptoms and 5 = severe infection (CIMMYT, 1995).

TABLE 3. Average yield, standard deviation, and yield stability of “INIAP H-603 Superior” and five commercial maize hybrids evaluated in 29 locations in the Ecuadorian lowlands from 2010 to 2013.

Hybrids t ha ⁻¹	Yield		
	Stability** bi	R ²	
H1	INIAP H-603	8.48 ± 1.97 a*	0.98
H6	DEKALB-1596	8.20 ± 2.29 ab	0.62
H3	INIAP H-601	7.95 ± 1.98 abc	0.84
H4	INIAP H-602	7.66 ± 2.01 bc	1.04
H5	DEKALB-7088	7.29 ± 2.08 cd	1.26
H2	INIAP H-553	6.92 ± 1.69 d	0.73
Tukey test (<i>P</i> <0.05)		0.72	

* Means and standard deviations followed by the same letter do not differ statistically according to the Tukey's range test (*P*<0.05).

** Values close to 1 are the most stable, according to Eberhart and Russel (1966).

are associated with production under irrigation (Zambrano *et al.*, 2017) and coincide with the observations of Arellano *et al.* (2011) and Caicedo *et al.* (2017), who found that irrigation with high yield hybrids resulted in a differential or yield potential of 5.0 and 7.0 t ha⁻¹, respectively.

Stability analysis

The genotype X environment interaction (GxE) is an important input in the evaluation and development of new maize hybrids based on yield (Zambrano *et al.*, 2017) and resistance to diseases (Parisi *et al.*, 2014), where climatic, soil and management variations produce a significant effect on the differentiation of hybrids. This was first proposed by Finlay and Wilkinson (1963), Eberhart and Russell (1966), and Zobel *et al.* (1988), but nowadays there are several models for estimating GxE, which are widely used in maize, such as the reliability based on differences

in yield with commercial controls (Camargo *et al.*, 2002) or the additive main effects and multiplicative interaction (AMMI) model (Caicedo *et al.*, 2017).

The Eberhart and Russel (1966) criterion for stability evaluates the relationship between environmental conditions and yield, where linear relationships are more stable. Alejos *et al.* (2006) showed how the Eberhart and Russel method is useful for evaluating genotypes with favorable or unfavorable responses to changing environments. In this study, "INIAP H-603 Superior" changed linearly with the change in the environment, with a regression coefficient (*bi*) of 0.98 and *R*² of 0.84 (Tab. 3), indicating that it is a genotype with good stability. These values are similar to those reported for new maize hybrids in Panama (Camargo *et al.*, 2002). Hybrids with *bi* values greater than 1, such as DEKALB-7088, which showed a *bi* of 1.26 and *R*² of 0.57,

TABLE 4. Average yield and standard deviation (t ha⁻¹) of the maize hybrids according to the growing conditions and hybrid X growing condition interaction evaluated in 29 locations in the Ecuadorian lowlands from 2010 to 2013.

Growing Condition		Yield (t ha ⁻¹)		
E2	Dry season (irrigated)	9.93 ± 1.37	a*	
E1	Rainy season	6.68 ± 1.87	b	
E3	Dry season (residual humidity)	6.65 ± 1.28	b	
Tukey test (<i>P</i> <0.05)		0.53		
Hybrid x Growing Condition				
H1 X E2		10.82 ± 1.15	a	
H6 X E2		10.48 ± 1.05	a	
H3 X E2		10.43 ± 1.13	a	
H4 X E2		9.69 ± 1.15	a	
H5 X E2		9.18 ± 1.69	a	b
H2 X E2		8.95 ± 1.06	a	b
H1 X E3		7.58 ± 1.44	b	c
H6 X E1		7.21 ± 2.26	c	d
H1 X E1		7.04 ± 1.58	c	d
H6 X E3		6.92 ± 1.20	d	
H4 X E3		6.79 ± 1.06	d	
H3 X E3		6.75 ± 1.12	d	
H3 X E1		6.68 ± 1.65	d	
H5 X E1		6.68 ± 2.04	d	
H4 X E1		6.50 ± 1.91	d	
H5 X E3		6.01 ± 1.15	d	
H2 X E1		5.95 ± 1.45	d	
H2 X E3		5.86 ± 1.06	d	
Tukey test (<i>P</i> <0.05)		1.95		
CV (%)		23.6		

* Means and standard deviations followed by the same letter do not differ statistically according to the Tukey's range test (*P*<0.05).

H1 = INIAP H-603 Superior, H2 = INIAP H-553, H3 = INIAP H-601, H4 = INIAP H-602, H5 = DEKALB-7088, H6 = DEKALB-1596.

are very sensitive to environmental changes. In addition, "INIAP H-603 Superior" had high yields in all tested environments. These kinds of genotypes should have a good response to high-input agriculture (Sabaghania *et al.*, 2006).

Conclusions

The crossing of domestic (INIAP lines) and CIMMYT lines has allowed the development of a high yielding maize hybrid, with an average of 8.48 t ha^{-1} , that is adapted to the varying local growing conditions. "INIAP H-603 Superior" is a hybrid well suited for the Ecuadorian lowlands, with resistance to the three principal foliar diseases: leaf blight (*Exserohilum turicum*), rust (*Puccinia sorghi*), and leaf spot (*Curvularia lunata*), and good yield stability ($bi = 0.98$). The greatest yield potential of the new hybrid (10.82 t ha^{-1}) was obtained with irrigation during the dry season.

Acknowledgments

This research was supported by the projects: "Fortalecimiento Institucional" and "Cambio de la Matriz Productiva" funded by the Ecuadorian Government. We would like to thank the farmers and technical team members that collaborated with this research on the development of the new hybrid at the Experimental Station and in farmers' fields. We would also like to thank CIMMYT, represented in Latin America by maize breeder partners Dr. Luis Narro and Dr. Felix San Vicente, for its constant technical support and germplasm, and Dr. Charles W. Barnes for his helpful comments and suggestions on the manuscript.

Literature cited

- Aguilar, D., D. Andrade, D. Álava, J. Burbano, M. Díaz, A. Garcés, W. Jiménez, D. Leiva, V. Loayza, W. Muyulema, P. Pérez, V. Ruiz, B. Simbaña, and R. Yépez. 2015. Estimación de superficie sembrada de arroz (*Oriza sativa L.*) y maíz amarillo duro (*Zea mays L.*) en las épocas de invierno y verano año 2015, en las provincias de Manabí, Los Ríos, Guayas, Santa Elena, Loja y El Oro. Quito, Ecuador. URL: http://sinagap.agricultura.gob.ec/pdf/estudios_agroeconomicos/estimacion_superficie_arroz_maiz-2015.pdf (accessed 1 November 2017).
- Alarcón, D., E. Zambrano, R. Limongi, and O. Cerón. 2016. INIAP H-603: Híbrido de maíz duro para Manabí y Los Ríos. Portoviejo, Ecuador: INIAP, Estación Experimental Portoviejo, Programa de Maíz. (Plegable no. 428). URL: <http://repositorio.iniap.gob.ec/handle/41000/4847> (accessed 1 November 2017).
- Alejos, G., P. Monasterio, and R. Rea. 2006. Análisis de la interacción genotipo ambiente para rendimiento de maíz en la región maicera del estado Yaracuy, Venezuela. Agron. Trop. 56(3), 369-384.
- Arellano, J., J. Vargas, I. Rojas, and M. Ávila. 2011. H-70: Híbrido de maíz de alto rendimiento para temporal y riego del altiplano central de México. Rev. Mex. Cienc. Agríc. 2(4), 619-626. Doi: 10.29312/remexca.v2i4.1651
- Caicedo, M., I. Ledesma, P. Villavicencio, A. Ezequiel, and D. Alarcón. 2017. Estabilidad ambiental en híbridos de maíz usando el modelo AMMI en el litoral ecuatoriano. ESPAM Ciencia 8(1), 23-32.
- Camargo, I., R. Gordón, J. Franco, A. González, E. Quirós, and A. Figueroa. 2002. Confiabilidad de nuevos híbridos de maíz, en Panamá. Agron. Mesoam. 13(1), 7-11.
- Castro, M. 2016. Rendimientos de maíz duro seco en invierno 2016. URL: http://sinagap.agricultura.gob.ec/pdf/estudios_agroeconomicos/rendimiento_maiz_duro_seco_invierno2016.pdf (accessed 1 November 2017).
- CIMMYT. 1995. Manejo de los ensayos e informe de los datos para el Programa de Ensayos Internacionales de Maíz del CIMMYT. URL: <https://repository.cimmyt.org/handle/10883/764> (accessed 1 November 2017).
- Córdova, H. 1991. Estimación de parámetros de estabilidad para determinar la respuesta de híbridos de maíz (*Zea mays L.*) a ambientes contrastantes de Centro América, Panamá y México. Agron. Mesoam. 2, 1-10.
- Di Renzo J.A., F. Casanoves, M.G. Balzarini, L. Gonzalez, M. Tablada, and C.W. Robledo. 2015. INFOSTAT versión 2015. Grupo InfoStat, FCA, Universidad Nacional de Córdoba, Argentina.
- Eberhart, S.A. and W.A. Russel. 1966. Stability parameters for comparing varieties. Crop. Sci. 6, 36-40. Doi: 10.2135/cropsci1966.0011183X000600010011x
- Finlay, K.W. and G.N. Wilkinson. 1963. The analysis of adaptation in a plant breeding programme. Aust. J. Agr. Res. 14, 742-754. Doi: 10.1071/AR9630742
- González, A., D. Pérez, J. Sahagún, O. Franco, E. Morales, M. Rubí, F. Gutiérrez, and A. Balbuena. 2010. Aplicación y comparación de métodos univariados para evaluar la estabilidad en maíces del valle de Toluca-Atlacomulco, México. Agron. Costarricense 34(2), 129-143.
- Gordon, R. and H. Deras. 2017. Evaluación de híbridos de maíz en ambientes contrastantes de Meso América, PCCMCA, 2016. LXII Reunión Anual del PCCMCA. San Salvador, El Salvador.
- Hernández, J.A. and F. Zavala. 2009. Adaptación y estabilidad del rendimiento de grano de genotipos de mijo (*Pennisetum americanum L. Leeke*) en San Luis Potosí, México. Rev. Fitotec. Mex. 32 (2), 143-152.
- IBPGR. 1991. Descriptors for maize. URL: https://www.bioversityinternational.org/fileadmin/user_upload/online_library/publications/pdfs/104.pdf (accessed 1 November 2017).
- Lin, C.S., M.R. Binns, and L.P. Lefkovitch. 1986. Stability analysis: Where do we stand? Crop Sci. 26, 894-900. Doi: 10.2135/cropscli1986.0011183X002600050012x
- MAG. 2015. Boletín situacional maíz duro seco. URL: http://sinagap.agricultura.gob.ec/phocadownloadpap/cultivo/2016/boletin_situacional_maiz_duro_seco_2015.pdf (accessed 1 November 2017).
- McIntosh, M. 1983. Analysis of combined experiments. Agron. J. 75(1), 153-155. Doi: 10.2134/agronj1983.00021962007500010041x

- Moore, K. and P. Dixon. 2015. Analysis of combined experiments revisited. *Agron. J.* 107, 763-771. Doi: 10.2134/agronj13.0485
- Parisi, L., L. Courerot, M. Hirsch, M. Fernández, and G. Magnone. 2014. Evaluación de enfermedades foliares, de raíz y tallo en un ensayo comparativo de híbridos de maíz de segunda y su relación con el rendimiento. URL: https://inta.gob.ar/sites/default/files/inta_pergamino_parisi_2014_evaluacion_de_enfermedades_foliares_de_raiz_y_tallo_en_un_ensayo_comparativo_de_hibridos_de_maiz.pdf (accessed 10 September 2017).
- Sabaghania, N., H. Dehghani, and S. Sabaghpour. 2006. Nonparametric methods for interpreting genotype x environment interaction of lentil genotypes. *Crop Sci.* 46, 1100-1106. Doi: 10.2135/cropsci2005.06-0122
- Vera, D., G. Liuba, L. Godoy, E. Díaz, F. Sabando, F. Garcés, and G. Meza. 2013. Análisis de estabilidad para el rendimiento de híbridos de maíz (*Zea mays*) en la Región Central del Litoral Ecuatoriano. *Sci. Agrop.* 4 (3), 211-218. Doi: 10.17268/sci.agropecu.2013.03.07
- Villavicencio, P. and J. Zambrano. 2009. Guía para la producción de maíz amarillo duro en la zona central del Litoral Ecuatoriano. URL: <http://repositorio.iniap.gob.ec/bitstream/41000/1551/1/Bolet%C3%ADn%20divulgativo%20N%C2%BA%20353.PDF> (accessed 9 October 2017).
- Zambrano, E., R. Limongi, D. Alarcón, P. Villavicencio, M. Caicedo, J. Eguez, and J. Zambrano. 2017. Interacción genotipo ambiente de híbridos de maíz bajo temporal en Manabí y Los Ríos, Ecuador. *ESPM Ciencia* 8(1), 7-14.
- Zobel, W.R., M.J. Wright, and H.G. Gauch. 1988. Statistical analysis of a yield trial. *Agron. J.* 80, 388-393. Doi: 10.2134/agronj1988.00021962008000030002x