

Effect of pollen doses on fruit formation and oil production in two hybrid palm genotypes (*Elaeis oleifera* H.B.K. Cortes x *Elaeis guineensis* Jacq.)

Efecto de dosis de polen en la formación de frutos y producción de aceite en dos genotipos híbridos de palma (*Elaeis oleifera* H.B.K. Cortes x *Elaeis guineensis* Jacq.)

Jéssica Socha¹, Daniel Cayón^{1*}, Gustavo Ligarreto¹, and Gabriel Chaves²

ABSTRACT

OxG oil palm hybrids (*Elaeis oleifera* x *Elaeis guineensis*) produce a small number of male inflorescences, with an irregular, slow anthesis phase and low-viability pollen that affect natural pollination, making the agronomic practice of assisted pollination necessary for plantations in order to guarantee the formation of the majority of bunch fruits. In the hybrids “Coarí x La Mé” and “(Sinú x Coarí) x La Mé”, the influence of several doses of pollen (0, 0.01, 0.05, 0.1, 0.15 g/inflorescence) on bunch weight, normal and parthenocarpic fruits, fruit set and oil/bunch potential was studied. A completely randomized experimental design was used with four replicates and nine inflorescences/replicate. The bunch weight was higher with the 0.05 g/inflorescence dose and lower with the natural pollination. The greatest fruit set was obtained with the dose of 0.1 g/inflorescence, and the oil/bunch percentage was higher with 0.05 g/inflorescence and lower with 0.01 g/inflorescence. The results confirm the need for assisted pollination in these new hybrid materials.

Key words: oil palm, physiology, pollination, fruit set, oil potential.

RESUMEN

Los híbridos OxG de palma de aceite (*Elaeis oleifera* x *Elaeis guineensis*) producen pocas inflorescencias masculinas con fase de antesis irregular lenta y polen de baja viabilidad afectando su polinización natural, lo cual hace necesaria la práctica agronómica de polinización asistida de las plantaciones para favorecer la formación de la mayoría de frutos del racimo. En los híbridos “Coarí x La Mé” y “(Sinú x Coarí) x La Mé” se estudió la influencia de varias dosis de polen (0, 0.01, 0.05, 0.1, 0.15 g/inflorescencia) sobre el peso del racimo, frutos normales y partenocárpicos, cuajado de frutos y potencial de aceite/racimo. El diseño experimental utilizado fue completamente al azar, en arreglo factorial 2x5, donde el factor 1 fueron los dos híbridos y el factor 2 las dosis de polen, con cuatro repeticiones y nueve inflorescencias/repetición. El peso del racimo fue mayor con la dosis 0.05 g/inflorescencia y menor con la polinización natural. El mayor cuajado de frutos se obtuvo con dosis de 0.1 g/inflorescencia, y el porcentaje de aceite/racimo fue superior con 0.05 g/inflorescencia y menor con 0.01 g/inflorescencia. Los resultados confirman la necesidad de la polinización asistida en los híbridos OxG.

Palabras clave: palma de aceite, fisiología, polinización, cuajado de frutos, potencial de aceite.

Introduction

OxG hybrids are crossbred between American (*Elaeis oleifera*) and African (*Elaeis guineensis*) oil palms and began to be studied in 1970 because some crosses showed resistance to the disease “Bud Rot” (BR), with production over 30 t of bunches/ha/year (Genty and Ujueta, 2013). These hybrids are a valuable alternative for oil production because of their tolerance to pests and diseases, bunch production (25 to 30 t ha⁻¹) and 18% oil extraction (Zambrano, 2004). In addition, their oil has high quality (55% oleic and 11% linoleic acids) and important contents of total vitamin E (tocotrienols 980 mg L⁻¹ and tocopherols 100 mg L⁻¹) (Mozzon *et al.*, 2013;

Mondragón and Pinilla, 2015) and carotenes (1375-1628 mg L⁻¹) (Rocha *et al.*, 2006; Rivera *et al.*, 2013; Choo and Nesaretnam, 2014).

The hybrid bunches have more parthenocarpic fruits (PF) than normal fruits (NF). The color and oil percentage of red PF are similar to those of NF, but white PF do not have any oil (González *et al.*, 2013). The advantage of hybrid red PF is that they develop and mature like NF, but contain more oil because of their higher proportion of mesocarp, which can reach 98% of the fruit (Bastidas *et al.*, 2007; Preciado *et al.*, 2011a). In addition, the contribution of red PF to the total bunch oil percentage is higher (20-50%) than that of NF (Preciado *et al.*, 2011b).

Received for publication: 2 October, 2018. Accepted for publication: 12 April, 2019

Doi: 10.15446/agron.colomb.v37n1.75313

¹ Faculty of Agricultural Sciences, Universidad Nacional de Colombia, Palmira (Colombia).

² Indupalma Farm, San Alberto (Colombia).

* Corresponding author: dgcajons@unal.edu.co



Studies on the physiological development of OxG hybrid bunches have shown that the fruits have filling and maturation problems, probably because of asynchronous openings of the flowers or insufficient pollination (Hormaza *et al.*, 2012; González *et al.*, 2013). In addition, female inflorescences are covered by fibrous spathes that hinder the entry of pollen and pollinators, which translates into a considerable percentage of bunches with aborted fruits (AF) and lower oil potential per bunch (OP) (Zambrano, 2004). For this reason, hybrid plantations employ the practice of assisted pollination, applying *E. guineensis* palm pollen to the inflorescences of hybrid palms to guarantee that female flowers in anthesis receive enough pollen for the development of fruits and reach a reasonable production of bunches (Prada and Romero, 2012; Barcelos *et al.*, 2015). The pollen quality depends on the genetic origin, viability, germination capacity and pollen tube growth (Godefroid *et al.*, 2010; Shivanna and Tandon, 2014) and is a fundamental requirement in assisted pollination in order to achieve complete fruit set of the bunch.

Haniff and Rosland (2002), in a study on assisted pollination in 12-year-old Tenera palms in Malaysia, evaluated increasing pollen doses (0.0001, 0.001, 0.01, 0.1, 1.0 and 5.0 g/inflorescence) and found that the dose of 0.01 g/inflorescence significantly increased the bunch weight (BW). Since assisted pollination is a very costly agronomic practice for plantations, it is necessary to generate specific management programs for hybrid materials that define the optimal dose of viable pollen in order to guarantee the formation of fruits and reduce costs. The objective of this study was to evaluate the influence of several doses of pollen on fruit development and potential oil production in OxG hybrids.

Materials and methods

Location and plant material

This study was conducted on the OxG hybrids Coarí x La Mé (CxLM) and (Sinú x Coarí) x La Mé (SxCxLM) in an Indupalma experimental field in San Alberto, Cesar, Colombia (10°20' N, 73°11' W, at 125 m a.s.l., maximum temperature 34°C, minimum temperature 22°C, 72% relative humidity, annual precipitation 2,497 mm/year, annual evaporation 1,208 mm and 2,130 h of sunshine/year).

Experimental design

The response of the two hybrids to four 76% germination pollen doses (0.15, 0.10, 0.05, and 0.01 g/inflorescence) and natural pollination as a control was evaluated in a completely randomized design (CRD) with a 2x5 factorial arrangement (factor 1: hybrids and factor 2: pollen doses), four replicates and nine inflorescences/replicate.

Pollination

The pollen used in this experiment was collected in August, 2013 from selected African Tenera palms, in which the male inflorescences were cut in anthesis (stage 607) according to the scale proposed by Hormaza *et al.* (2012) and taken to the Indupalma Seed Laboratory, where the finger-like spines were separated on Kraft paper and dried in a forced convection oven for 12 h at 39°C until 6% humidity; then, they were shaken carefully to release the pollen from the anthers, which was screened with No. 100 and 200 sieves to remove impurities, vacuum packed and stored at -4°C. The pollen quality used in the treatments was determined with the germination test described by Turner and Gilbanks (1974), which counts the percentage of pollen grains that emit a pollen tube; only pollen with 76% germination was used.

The female inflorescences were selected in phenological stage 602 or preanthesis II (Hormaza *et al.*, 2012) (Fig. 1A); then, the spathes were removed (Fig. 1B) and covered with insulation bags fitted to the base with an elastic band to avoid contamination from surrounding natural pollen (Fig. 1C). The pollination treatments were carried out when the inflorescences reached the anthesis stage (phenological stage 607), that is, when more than 80% of the flowers had cream-colored receptive stigmas (Hormaza *et al.*, 2011) (Fig. 1D). Before the application of the pollen, it was verified that no live insects were inside the bagged female inflorescences to avoid any contamination from foreign pollen; next, the pollen, mixed with inert talc as a transport agent (1:9 ratio), was generously sprinkled through a small hole made with a punch in the plastic window of the isolation bags (Fig. 1E).

The mature bunches were harvested 154 d after pollination, when they naturally detached one to five fruits per bunch (Fig. 1F). The physical analysis of the bunches was done with the methodology described by García and Yáñez (2000) to determine the BW, weight of NF and PF, fruit set coefficient (1) and bunch oil potential (OP) (2) with a subsample of 10 g of dry mesocarp using the Soxhlet method, with hexane as the solvent.

$$\text{Fruit set} = \frac{\text{normal fruits} + \text{parthenocarpic fruits} + \text{aborted fruits}}{\text{total fruits}} \times 100 \quad (1)$$

$$\text{Oil potential (\%)} = \frac{\% \text{ normal fruits oil}}{\text{fruits oil}} + \frac{\% \text{ parthenocarpic fruits oil}}{\text{fruits oil}} \quad (2)$$

Statistical analysis

The data were subjected to an analysis of variance (ANOVA) and, for the differences between treatments, the




FIGURE 1. Development stages of the female inflorescences (Hormaza *et al.*, 2012). A) pre-anthesis II (phenological stage 602), B) isolated inflorescence, C) enfolded inflorescence, D) anthesis (phenological stage 607), E) pollination, F) mature bunch.

Duncan test ($P < 0.05$) was used with the statistical program SAS® (SAS Institute, Cary, NC, USA).

Results and discussion

Bunch weight

Bunch weight (BW) is the sum of the combined weight of spikelets with their individual fruits and the weight of the peduncle or rachis (Mohd, 2000). Figure 2 shows that the higher doses of pollen (0.15, 0.05) produced bunches with a higher weight than the lower doses and natural pollination. Similar trials have also reported a significant increase in BW with doses higher than 0.01 g/inflorescence (Haniff and Roslan, 2002). The natural pollination of the hybrids was very low because of the low emission of male inflorescences that produced a small amount of pollen (5 to 10 g/inflorescence) with low viability (2 to 25%) (Hormaza *et al.*, 2012), which was unattractive to the pollinating insect *Elaeidobius kamerunicus* (Tan, 1985).

Fruit set

In the two hybrids, the higher pollen doses provided the largest fruit set, with significant statistical differences

when compared to the lowest dose (0.01 g) and the natural pollination control (Fig. 3). The expected result took into account the fact that fruit set depends on the proportion of NF and PF in bunches and, consequently, is directly related to BW, as effectively observed in this study with the higher BW obtained with the high doses. These results agree with those of Haniff and Roslan (2002), who observed a significant increase in BW and fruit set with a pollen dose that was increased from 0.0001 to 0.01 g when using assisted pollination on *E. guineensis* palms.

Fruits/bunch and oil potential

Table 2 shows that there were no significant statistical differences between the hybrids and pollen doses for the NF, PF and AF percentages. However, the tendency for a higher percentage of NF rather than PF was observed with the higher pollen doses. This is a logical consequence of the amount of pollen grains that reach the inflorescence in anthesis, which favors the formation of NF fruits. It has been proven that the assisted pollination of inflorescences in *E. guineensis* palms and OxG hybrids substantially increases the proportion of NF/bunch (Tam, 1981; Rosero and Santacruz, 2014, 2017).




FIGURE 2. Effect of the pollen dose on the bunch weight of two OxG oil palm hybrids. Bars with different letters in each hybrid are statistically different according to the Duncan test ($P < 0.05$).




FIGURE 3. Effect of pollen dose on the fruit set of the two oil palm OxG hybrids. Bars with different letters in each hybrid are statistically different according to the Duncan test ($P < 0.05$).

TABLE 2. Effect of pollen dose (g/inflorescence) on the normal fruits (NF), parthenocarpic fruits (PF), aborted fruits (AF), and oil potential (OP) in the bunches of the two OxG hybrids.

Hybrid	Dose	Fruits (%)			OP (%)
		NF	PF	AF	
Coarí x La Mé	0.15	44 a	30 ab	10	23.0 ab
	0.1	43 a	35 a	9	25.7 a
	0.05	46 a	29 ab	9	27.0 a
	0.01	31 ab	22 b	11	19.3 b
	Natural pollination	22 b	23 b	8	22.4 ab
(Sinú x Coarí) x La Mé	0.15	49 a	28 a	6	23.4 a
	0.1	45 a	21 b	5	22.1 ab
	0.05	44 a	24 ab	9	20.4 b
	0.01	32 b	21 b	10	21.1 b
	Natural pollination	33 b	16 c	10	18.8 c
CV (%)		30.0	26.5	60.6	24.0
F (Doses)		8.1*	4.2*	0.28 ns	2.2*
F (Hybrids)		1.3 ns	6.1*	0.42 ns	2.64 ns
F (Hybrids x Doses)		0.3 ns	0.37 ns	0.95 ns	0.89 ns

*F test significant ($P < 0.05$). ns F test not significant ($P < 0.05$)

The higher proportion of PF in hybrid bunches is attributed to the maternal inheritance of the palm *E. oleifera*, characterized by numerous PF that can form up to 90% of the total bunch (Corley and Tinker, 2009), which occurs despite a high percentage of viable pollen (Hardon and Tan, 1969). Zambrano (2004) reported a proportion of between 22 and 49% PF/bunch in OxG hybrids, while Rosero and Santacruz (2017) reported between 44 and 56% PF/bunch. PF increases in bunches have a large impact on the bunch oil potential because the oil content is similar to or even higher than that of NF since the oil in PF pulp can reach up to 98% (Bastidas *et al.*, 2011).

The low percentage of AF with all of the pollen doses indicates that the presence of these fruits was not due to the greater or lesser amount of pollen that reached the inflorescences in anthesis, but resulted from adverse factors during the initial development of the bunch fruits. Although the abortion of inflorescences and fruits usually occurs as a consequence of incorrect pollination, some extreme factors, such as water stress, fructification, pruning or shading, can increase it (Corley and Breure, 1992). The AF in the control treatment (natural pollination) were low in number because the experimental field was located between plots of *E. guineensis*, which, most likely, served as a source of

pollen for the two hybrids. Hardon and Turner (1967) used natural pollination on *E. guineensis* palm plantations to demonstrate that the majority of pollen remains inside the canopy after emission by male inflorescences, a factor that may favor its transport to the palms.

The OP was higher with the higher doses of pollen (0.15 and 0.10), which showed higher percentages of NF and PF. This coincides with Corley and Tinker (2009), who stated that the variation that exists in OP depends on the proportion of NF and PF in the bunch, which contain oil. In fact, OP indicates the amount of oil that can be extracted per unit of fresh bunch (Moreno *et al.*, 2017).

Conclusions

The pollen doses of 0.10 and 0.15 g/inflorescence increased the NF, decreased the AF in the bunches, and favored the production of a higher BW, fruit set, and OP in the two analyzed hybrids.

Literature cited

- Barcelos, E., S.A. Rios, R.N.V. Cunha, R. Lopes, S.Y. Motoike, E. Babiychuk, A. Skirycz, and S. Kushnir. 2015. Oil palm natural diversity and the potential for yield improvement. *Front. Plant Sci.* 6, 190. Doi: 10.3389/fpls.2015.00190
- Bastidas, S., E. Peña, R. Reyes, J. Pérez, and W. Tolosa. 2007. Comportamiento agronómico del cultivar híbrido RC1 de palma de aceite (*Elaeis oleifera* x *Elaeis guineensis*) x *Elaeis guineensis*. *Corpoica Cienc. Tecnol. Agropec.* 8(1), 5-11. Doi: 10.21930/rcta.vol8_num1_art:77
- Bastidas, S., C. Betancourth, C. Preciado, E. Peña, and R. Reyes. 2011. Predicción y control de la cosecha en el híbrido interespecífico *Elaeis oleifera* x *Elaeis guineensis* en la zona palmera occidental de Colombia II. Determinación del ciclo de cosecha para obtener racimos con alto contenido de aceite. *Corpoica Cienc. Tecnol. Agropec.* 12(1), 13-20. Doi: 10.21930/rcta.vol12_num1_art:211
- Corley, R.H.V. and C.J. Breure. 1992. Fruiting activity, growth and yield of oil palm. I. Effects of fruit removal. *Exp. Agric.* 28(1), 99-109. Doi: 10.1017/s0014479700023048
- Choo, Y. and K. Nesaretnam. 2014. Research advancements in palm oil nutrition. *Eur. J. Lipid Sci. Technol.* 116(10), 1301-1315. Doi: 10.1002/ejlt.201400076
- Corley, R.H.V. and P.B. Tinker. 2009. La palma de aceite. Federación Nacional de Cultivadores de Palma de aceite (Fedepalma), Bogotá.
- García, J.A. and E.E. Yáñez. 2000. Aplicación de la metodología alterna para análisis de racimos y muestreo de racimos en tolva. *Palmas* 21, 303-311.
- Genty, P. and M. Ujueta. 2013. Relatos sobre el híbrido interespecífico de palma de aceite OxG-Coarí x La Mé: esperanza para el trópico. Fedepalma, Bogotá.
- Godefroid, S., A. Van de Vyver, and T. Vanderborgh. 2010. Germination capacity and viability of threatened species collections in seed banks. *Biodiversity Conserv.* 19(5), 1365-1383. Doi: 10.1007/s10531-009-9767-3
- González, D.A., G. Cayón, J.E. López, and W.H. Alarcón. 2013. Development and maturation of fruits of two Indupalma OxG hybrids (*Elaeis oleifera* x *Elaeis guineensis*). *Agron. Col.* 31(3), 343-351.
- Haniff, H. and M.D. Rosland. 2002. Fruit set and oil palm bunch components. *J. Oil Palm Res.* 14(2), 24-33.
- Hardon, J. and Y. Tan. 1969. Interspecific hybrids in the genus *Elaeis* I. crossability, cytogenetics and fertility of F₁ hybrids of *E. guineensis* x *E. oleifera*. *Euphytica* 18(1), 372-379. Doi: 10.1007/BF00397784
- Hormaza, P., E. Mesa, and H.M. Romero. 2012. Phenology of the oil palm interspecific hybrid *Elaeis oleifera* x *Elaeis guineensis*. *Sci. Agric.* 69(4), 275-280. Doi: 10.1590/S0103-90162012000400007
- Mohd, H.H. 2000. Yield and yield components and their physiology. pp. 146-170. In: Basiron, Y., B.S. Jalani, and K.W. Chan (eds.). *Advances in oil palm research*. Volume I. MPOB, Bangi Selangor, Malaysia.
- Mondragón, A. and C. Pinilla. 2015. Aceite de palma alto oleico: propiedades fisicoquímicas y beneficios para la salud humana. *Palmas* 36(4), 57-66.
- Moreno, E., J.M. García, C.A. Díaz, and N.E. Ramírez. 2017. Optimización de la medición del potencial industrial de aceite. *Palmas* 38(4), 98-107.
- Mozzon, M., D. Pacetti, P. Lucci, M. Balzano, and N.G. Frega, 2013. Crude palm oil from interspecific hybrid *Elaeis oleifera* x *Elaeis guineensis*: fatty acid region distribution and molecular species of glycerides. *Food Chem.* 141(1), 245-252. Doi: 10.1016/j.foodchem.2013.03.016
- Preciado, C.A., S. Bastidas, C. Betancourth, E. Peña, and R. Reyes. 2011a. Predicción y control de la cosecha en el híbrido interespecífico *Elaeis oleifera* x *Elaeis guineensis* en la zona palmera occidental de Colombia. I. Determinación del período de madurez para obtener racimos de alto contenido de aceite. *Corpoica Cienc. Tecnol. Agropec.* 12(1), 5-12. Doi: 10.21930/rcta.vol12_num1_art:210
- Preciado, C.A., S. Bastidas, C. Betancourth, E. Peña, and R. Reyes. 2011b. Predicción y control de la cosecha en el híbrido interespecífico *Elaeis oleifera* x *Elaeis guineensis* en la zona palmera occidental de Colombia. II. Determinación del ciclo de cosecha para obtener racimos de alto contenido de aceite. *Corpoica Cienc. Tecnol. Agropec.* 12(1), 13-19. Doi: 10.21930/rcta.vol12_num1_art:211
- Rivera, Y.D., D.G. Cayón, and J.E. López. 2013. Physiological and morphological characterization of American oil palms (*Elaeis oleifera* HBK Cortes) and their hybrids (*Elaeis oleifera* x *Elaeis guineensis*) on the Indupalma plantation. *Agron. Colomb.* 31(3), 314-323.
- Rocha, P., F. Prada, L. Rey, and I. Ayala. 2006. Caracterización bioquímica parcial de la colección de *Elaeis oleifera* de Cenipalma proveniente de la Amazonia colombiana. *Palmas* 27(3), 35-44.
- Rosero, G. and L. Santacruz. 2014. Efecto de la polinización asistida en la conformación del racimo en material híbrido OxG en la plantación Guaicaramo S.A. *Palmas* 35(4), 11-19.

- Rosero, G., L. Santacruz, A. Ríos, and S. Carvajal. 2017. Influencia del destape de la inflorescencia en la polinización asistida del híbrido OxG. *Palmas* 38(1), 49-62.
- Shivanna, K.R. and R. Tandon. 2014. Reproductive ecology of flowering plants: A manual. Springer, New Delhi, India. Doi: 10.1007/978-81-322-2003-9
- Tam, T.K. 1981. Investigations into fruit set capacities of the *Elaeis oleifera* under controlled pollination conditions and germination requirements of the *Elaeis oleifera* x *Elaeis guineensis* (*pisifera*) hybrid [oil palms]. *The Planter* 57, 444-451.
- Tan, Y.P. 1985. Weevil pollination in the *Elaeis oleifera* x *Elaeis guineensis* hybrid. pp. 34-40. In: Proceedings of the Symposium on impact of pollinating weevil on the Malaysian oil palm industry. Institut Penyelidikan Minyak Kelapa Sawit Malaysia, Kuala Lumpur.
- Turner, P.D. and R.A. Gillbanks. 1974. Oil palm cultivation and management. The Incorporated Society of Planters. Kuala Lumpur.
- Zambrano, J. 2004. Los híbridos interespecíficos *Elaeis oleifera* x *Elaeis guineensis*. Una alternativa de renovación para la zona oriental de Colombia. *Palmas* 25 (special issue), vol. 2, 339-349.