

Interaction of triacontanol with other plant growth regulators on morphology and yield of field pea (*Pisum sativum* L.)

Interacción del triacontanol con otros reguladores de crecimiento vegetal en la morfología y el rendimiento de arveja (*Pisum sativum* L.)

Héctor Cantaro-Segura^{1*} and Amelia Huaranga-Joaquín¹

ABSTRACT

The effect of a recent plant growth regulator, triacontanol (TRIA), on plant growth and yield of *Pisum sativum* L. was investigated. The experiment was carried out under field conditions at the Instituto Nacional de Innovación Agraria (INIA), La Molina, Lima, Peru, using a completely randomized block design with eight treatments and three replicates. Treatments consisted in the foliar application of TRIA alone and in all possible combinations with three plant growth regulators based on auxins (AUX), gibberellins (GA), and cytokinins (CK), on pea plants cv. Rondo. The highest green pod yields were obtained with the application of TRIA+AUX+GA+CK, and TRIA+AUX+CK. The TRIA+AUX increased the values of the yield variables while TRIA+GA increased the values of the morphological variables. TRIA+CK showed a stimulating effect on morphological variables and number of grains per pod, while TRIA+AUX+CK acted synergistically on yield variables since their combined effect overweighed the effect of each growth regulator separately. Treatments with TRIA exceeded the control treatment in yield variables, indicating its great potential to be used in sustainable agriculture to guarantee food security in the future.

Key words: auxins, cytokinins, gibberellins, plant physiology, legumes.

RESUMEN

Se investigó el efecto de un regulador del crecimiento vegetal relativamente nuevo, el triacontanol (TRIA), sobre el crecimiento de las plantas y el rendimiento de *Pisum sativum* L. El experimento se llevó a cabo en condiciones de campo en el Instituto Nacional de Innovación Agraria (INIA), Lima, Perú, utilizando un diseño de bloques completamente al azar con ocho tratamientos y tres repeticiones. Los tratamientos consistieron en la aplicación foliar de TRIA solo y en todas las combinaciones posibles con tres reguladores del crecimiento vegetal a base de auxinas (AUX), giberelinas (GA) y citoquininas (CK), en plantas de arveja cv. Rondo. Los mayores rendimientos de vaina verde se obtuvieron con la aplicación de TRIA+AUX+GA+CK y TRIA+AUX+CK. TRIA+AUX incrementó los valores de las variables de rendimiento mientras que TRIA+GA incrementó los valores de las variables morfológicas. TRIA+CK mostró un efecto estimulante sobre las variables morfológicas y el número de granos por vaina, mientras que TRIA+AUX+CK actuó sinérgicamente en las variables de rendimiento, ya que su efecto combinado sobrepesó el efecto de cada regulador de crecimiento por separado. Los tratamientos con TRIA superaron al tratamiento control en las variables de rendimiento, lo que indica su gran potencial de uso en la agricultura sostenible para garantizar la seguridad alimentaria en el futuro.

Palabras clave: auxinas, citoquininas, giberelinas, fisiología vegetal, leguminosas.

Introduction

The field pea (*Pisum sativum* L.) belongs to the Fabaceae family and is one of the oldest domesticated species for human and livestock consumption. It is widely cultivated around the world (Wu *et al.*, 2019) and predominant in world trade, representing about 35-40% of the total trade in legumes. It is also among the most consumed vegetables worldwide (Ratnayake *et al.*, 2001). The pea is an important nutritional crop (Liu *et al.*, 2015) since its grains have high contents of protein (18-30%), vitamins and minerals. Additionally, its shoots and leaves can be used for fresh

consumption as leafy vegetables (Santos *et al.*, 2014), with a good demand in the national and international markets. Field pea is a profitable agricultural product. However, this profitability is diminished in many regions by various factors such as environmental problems, incorrect agronomic management, and increase in production costs due to the rise in the application of fertilizers and pesticides (Erisman, 2011). Agronomic management is especially critical because long-duration varieties with low yields are still sown in production areas and are susceptible to different diseases and pests (Wu *et al.*, 2019). For this crop, correct physiological and nutritional management is imperative,

Received for publication: 3 March, 2021. Accepted for publication: 17 August, 2021.

Doi: 10.15446/agron.colomb.v39n2.94096

¹ Departamento de Fitotecnia, Facultad de Agronomía, Universidad Nacional Agraria La Molina, Lima (Peru).
* Corresponding author: hcantaro@lamolina.edu.pe



with biostimulants and plant hormones playing an important role (Bertolin *et al.*, 2010; Cato *et al.*, 2013; Martínez González *et al.*, 2017).

Plant growth regulators are small molecules that can trigger different physiological processes related to the growth, development, and defense of plants (Pozo *et al.*, 2015). They are also known as phytohormones, but this term is not used frequently in agriculture (Davies, 2010). Plant hormones, growth regulators, and inhibitors have been used in practice to increase yield, improve quality, or alleviate the adverse effects induced by biotic or abiotic stresses (Csukasi *et al.*, 2009). Different classes of hormones have already been characterized, including abscisic acid, auxins, brassinosteroids, cytokinins, ethylene, gibberellins, jasmonates, strigolactones, etc. (Depuydt & Hardtke, 2011). All of them have been linked in one way or another to growth regulation (Santner *et al.*, 2009; Wolters & Jürgens, 2009). Cytokinins, auxins, gibberellins, and brassinosteroids are considered essential for growth of mutant phenotypes in which hormone biosynthesis or perception is disrupted; cytokinins regulate cell proliferation, while gibberellins promote cell elongation and auxins are involved in both processes. Furthermore, brassinosteroids are essential for cell elongation, but may also play a role in cell division (Nakaya *et al.*, 2002; Hardtke *et al.*, 2007). All these hormones can regulate a high number of processes in a unique and independent way. However, cooperation and interrelationship between signaling metabolic pathways appears to exist, as it follows from the superimposed influence on various cellular processes (Hardtke *et al.*, 2007).

New hormones and plant growth regulators are still being discovered, and their interrelationships, mechanisms of action, and relationships at the metabolic level are being exhaustively studied (Gomez-Roldan *et al.*, 2008; Kuppusamy *et al.*, 2009). Among these substances is triacontanol (TRIA), a growth regulator recently used in commercial applications despite the fact that it was discovered a few decades ago in natural waxes. It can exert stimulating effects, even at considerably low foliar concentrations (Khandaker *et al.*, 2013). TRIA is not considered a plant hormone since it is a secondary substance in plant growth (Naeem *et al.*, 2012). However, various studies show that its effects on growth and yield in plants are not shown by other plant hormones or growth regulators. The positive role of TRIA in photosynthesis, nitrogen fixation, enzymatic activities, free amino acids, reducing sugars, and soluble proteins in plants has been documented (Borowski *et al.*, 2000; Naeem *et al.*, 2009; 2010; 2011; Aftab *et al.*, 2010; Khandaker *et al.*, 2013). The application of TRIA also increases the dry

weight, chlorophyll content, protein and net photosynthetic efficiency in rice (*Oryza sativa* L.) (Chen *et al.*, 2002). In cotton, it promotes vegetative growth and increases the level of monogalactosyldiacylglycerol (MGDG), a galactolipid that appears to be involved in the synthesis of photosystem I proteins (Naeem *et al.*, 2012). The application of TRIA alone or in combination with potassium increases plant height, fresh and dry weight of the plant and leaf area in tomato (Khan *et al.*, 2009).

In this exploratory study, we evaluated the effect of the interaction of plant growth regulators with triacontanol on the field pea *Pisum sativum* L. cv. Rondo to determine the roles of growth regulators applied alone and in combination and their impact on yield components and plant growth and development.

Materials and methods

The experiment was carried out in the experimental fields of the Instituto Nacional de Innovación Agraria (INIA), La Molina, Lima, Peru (12°04'48" S, 76°56'44" W at 243 m a.s.l.) between September and December 2015. To determine the effect of the application of triacontanol (TRIA) and the growth regulators on peas, pea seeds cv. Rondo were used. Rondo is a commercial cultivar with a determinate growth habit, semi-late, with straight pod and rough grain suitable for fresh consumption, and with medium maturation appreciated by farmers for its yield and pod size. Climate characteristics of the field site are described in Table 1. Four pea seeds were sown per hole under drip irrigation at a distance between plants of 0.75 x 0.2 m, with 24 experimental plots of 9 m² and a total area of 320 m² (experimental plots + edge spaces). Average soil characteristics of the field site were: sand, 53%; silt, 28%; clay, 19%; organic matter, 1.12%; P, 17.4 mg kg⁻¹; K, 291 mg kg⁻¹; pH, 7.7, and electrical conductivity, 1.62 dS m⁻¹. The soil analysis was carried out according to the guidelines established by Burt (2014). Subsequently, manual thinning was done, leaving three plants per stroke and a final planting density of 200,000 plants ha⁻¹.

At 43 d after sowing (V4 growth stage), the treatments described in Table 2 were foliar applied. All treatments, except the control, alone or in combination with other plant growth regulators (auxins (AUX), gibberellins (GA) or cytokinins (CK)) contained TRIA, and a foliar-applied potassium fertilizer (K). Other commercial foliar fertilizers from FARMEX S.A. (Peru), such as FX Amino and Powergizer, were also applied to all treatments at the same concentration to replace nitrogen deficiencies.

TABLE 1. Average climate data during the crop cycle.

Month	Maximum temperature (°C)	Minimum temperature (°C)	Relative humidity (%)	Mean precipitation (mm/month)
September	21.52	15.34	80.21	0.12
October	22.77	16.16	79.75	0.06
November	22.9	16.64	79.33	0.09
December	25.08	17.97	77.92	0.06

To evaluate the effect of the treatments on growth and yield, the plant height, number of branches per plant, plant fresh weight, pod length, pod width, number of pods per plant, number of grains per pod and green pod yield ($t\ ha^{-1}$) were determined at harvest.

A completely randomized block design was used, with eight treatments and three replicates, for a total of 24 experimental units. The results were subjected to a normality test and analysis of variance (ANOVA) by the F test, and the means were compared with the least significant Fisher's test (LSD) at 5% probability of error, using the statistical software R. Pearson's correlation test at $P \leq 0.05$ was applied to assess the significance of correlation coefficients.

TABLE 2. Description of the treatments and doses of the plant growth regulators used in the experiment.

Name	Description
TRIA+AUX	0.5 mg L ⁻¹ triacontanol + 10 mg L ⁻¹ NAA
TRIA+GA	0.5 mg L ⁻¹ triacontanol + 10 mg L ⁻¹ GA ₃
TRIA+CK	0.5 mg L ⁻¹ triacontanol + 1 mg L ⁻¹ KIN
TRIA+AUX+CK	0.5 mg L ⁻¹ triacontanol + 0.5 mg L ⁻¹ NAA + 1 mg L ⁻¹ KIN
TRIA+AUX+GA+CK	0.5 mg L ⁻¹ triacontanol + 0.5 mg L ⁻¹ NAA + 1 mg L ⁻¹ GA ₃ + 1 mg L ⁻¹ KIN
TRIA+K	0.5 mg L ⁻¹ triacontanol + 3000 mg L ⁻¹ K ₂ O
TRIA	0.5 mg L ⁻¹ triacontanol
CONTROL	Treatment without any application of triacontanol or growth regulators

NAA - 1-Naphthaleneacetic acid; GA₃ - gibberellic acid; KIN - kinetin; K₂O - potassium oxide.

Results and discussion

Growth parameters

The plant growth regulators applied in this study caused changes in plant height. The application of TRIA + GA obtained the highest plant height (143.5 cm), and the control obtained the lowest value (79.8 cm), which was 15.6% lower than the average obtained in the experiment (Tab. 3). The height of the pea plant is a characteristic determined by its genetics (Weeden, 2007), and the interaction with growth-promoting phytohormones such as GA (Wang *et*

al., 2017). The control of this experiment showed a height greater than that reported by Anchivilca Rojas (2018) (68 cm), Santos *et al.* (2018) (60 cm), and Rodríguez Quispe (2015) (52 cm). According to Checa Coral *et al.* (2020), this could be due to environmental conditions and an efficient supply of resources since the resulting phenotype is the interaction of the genotype and the environment. The positive interaction of TRIA and GA agrees with that proposed by Shukla *et al.* (1992), who mention that triacontanol potentiates the effect of GA. The opposite is observed with AUX, since TRIA decreases its influence as a “destruction” effect is produced. The intensity of this effect is specific in each cultivar (Henry & Gordon, 1980); auxins at low concentrations inhibit stem growth in favor of root growth and development (Davies, 2010; Taiz *et al.*, 2014). No effects were observed with foliar application of potassium because the soil was rich in this element (291 mg L⁻¹) and the application of potassium would keep the plants under conditions of “luxury consumption” (Marschner, 2011).

Regarding the number of branches per plant, there were no statistical differences in the analysis of variance for the treatments. Most of the treatments obtained similar values to those of the control, except for TRIA+GA, with a value 21.8% less than the average, and TRIA+AUX+GA+CK, with a value 23.5% greater than the control and 26.3% greater than the average (Tab. 3). The trend is repeated for pod length and plant fresh weight regarding the highest and lowest values in the treatments. However, for pod width, the control showed the lowest value of 1.66 cm, TRIA+AUX showed an intermediate value of 1.76 cm, and TRIA+AUX+GA+CK obtained the highest value with 1.83 cm.

The number of branches plant (NBP) and the pod width are also genetically determined characteristics, but the former is more susceptible to variations due to environmental conditions and/or agronomic practices than the latter (presence of the *n* gene, Wehner & Gritton, (1981)). Pod width did not exhibit significant or percentage statistical variation from all variables, showing a less marked incremental effect of the negative interaction of TRIA with AUX and CK applied jointly (Fig. 1). On the other hand, for NBP, the treatments

TABLE 3. Effect of plant growth regulators on growth and yield of pea plants.

Treatment	PH (cm)	NBP	PL (cm)	PW (cm)	NPP	NGP	PFW (g)	Yield (t ha ⁻¹)
TRIA+AUX	86.5 ^b ± 8.3	2.7 ^{ab} ± 0.4	9.67 ^{ab} ± 0.7	1.77 ^{ab} ± 0.03	9.5 ^{abc} ± 1.0	6.1 ^c ± 0.7	130.2 ^{bc} ± 21.5	7.43 ^c ± 0.8
TRIA+GA	143.5 ^a ± 4.6	2.2 ^b ± 0.3	8.26 ^b ± 0.4	1.83 ^{ab} ± 0.09	3.7 ^c ± 0.9	4.6 ^d ± 0.01	93.5 ^c ± 11.5	2.15 ^d ± 0.3
TRIA+CK	95.8 ^b ± 6.4	2.8 ^{ab} ± 0.2	10.67 ^a ± 0.1	1.74 ^{ab} ± 0.03	12.2 ^{ab} ± 3.4	7.9 ^{ab} ± 0.3	143.3 ^{abc} ± 15.6	8.47 ^{bc} ± 0.9
TRIA+AUX+CK	85.0 ^b ± 6.1	2.8 ^{ab} ± 0.2	10.29 ^a ± 0.4	1.68 ^{ab} ± 0.02	15.0 ^a ± 1.5	7.7 ^{ab} ± 0.2	154.5 ^{ab} ± 7.2	10.53 ^{ab} ± 0.9
TRIA+ AUX+GA+CK	91.5 ^b ± 4.5	3.5 ^a ± 0.5	10.71 ^a ± 0.7	1.83 ^a ± 0.06	14.7 ^{ab} ± 3.2	8.7 ^a ± 0.2	198 ^a ± 34.9	12.15 ^a ± 1.4
TRIA+K	85.7 ^b ± 3.3	2.7 ^{ab} ± 0.2	10.46 ^a ± 0.4	1.81 ^{ab} ± 0.03	9.0 ^{bc} ± 1.3	7.7 ^{ab} ± 0.6	120.8 ^{bc} ± 15.4	7.30 ^c ± 1.0
TRIA	88.7 ^b ± 3.3	2.7 ^{ab} ± 0.2	10.21 ^a ± 0.3	1.73 ^{ab} ± 0.05	12.8 ^{ab} ± 0.9	7.2 ^{bc} ± 0.4	135.8 ^{bc} ± 9.3	8.88 ^{bc} ± 0.4
CONTROL	79.8 ^b ± 7.2	2.8 ^{ab} ± 0.2	9.33 ^{ab} ± 0.5	1.66 ^b ± 0.04	13.3 ^{ab} ± 1.5	5.9 ^{cd} ± 0.3	122.2 ^{bc} ± 14.6	7.98 ^{bc} ± 1.0
F-Test	**	ns	ns	ns	*	**	ns	**
Mean	94.56	2.77	9.95	1.75	11.27	6.98	137.29	8.11

TRIA - triacontanol, AUX - auxins, GA - gibberellins, CK - cytokinins, K - foliar-applied potassium, PH - plant height, NBP - number of branches per plant, PL - pod length, PW - pod width, NPP - number of pods per plant, NGP - number of grains per pod, PFW - plant fresh weight. ns - not significant at $P > 0.05$; * : $P \leq 0.05$; ** : $P \leq 0.01$.

with CK increased the branching of the shoots (Müller & Leyser, 2011) and the number of branches in the plants (Taiz *et al.*, 2014).

Pearson's correlation coefficients showed interesting data regarding the relationships between variables. Within the vegetative parameters, plant height was negatively correlated with pod length (-0.271) and the number of branches per plant was positively correlated with pod length (0.442*) and yield (0.731**). The variable pod length was positively correlated with plant fresh weight (0.626**), number of grains per pod (0.823**), and green pod yield (0.654**). A positive correlation was also observed between these last three variables. However, these variables were not positively correlated with pod width (Tab. 4). In contrast with the values obtained in the variable plant height, the average pod length in this experiment (9.9 cm) was lower than that obtained by Anchivilca Rojas (2018) (10 cm), Rondinel Ruíz (2014) (10.2 cm), and Rodríguez Quispe (2015) (11.4 cm). Triacontanol application resulted in an increase of 11.3% in yield over the control (Fig. 2). A positive interaction was also observed with treatments containing CK and, to a lesser extent, with foliar potassium. A slight negative interaction was observed with AUX and a higher negative interaction was registered with gibberellins.

The effect of CK is significant since they promote greater pod setting in legumes (Carlson *et al.*, 1987) and increase the width, length, and weight of pods (Mosjidis *et al.*, 1993). Potassium also has a slight incremental effect on fruiting and a higher pod setting in legumes (Marschner, 2011). However, this is not reflected by a luxury consumption of K. The trend of the effect of the hormonal interaction is repeated on pod length, as well as on plant fresh weight,

except that a negative interaction with foliar potassium was also recorded. Although gibberellins favor the vegetative growth of the plant when applied in excess (Taiz *et al.*, 2014), this growth is very fast, forming tall but weak stems with lower dry matter content and fresh weight of the plant. The mixture of the three hormones considerably increases the plant fresh weight due to a better regulation of physiological processes and the internal hormonal balance in the plants (Figs. 1-3).

Yield parameters

The lowest values of all yield variables (number of pods per plant, number of grains per pod, and green pod yield) were obtained with TRIA+GA. The highest values were recorded with TRIA+AUX+GA+CK, in which all the hormones were used, except in the number of pods per plant, where the best treatment was TRIA+AUX+CK (Tab. 3). The number of pods per plant showed a different behavior between treatments, although it was highly positively correlated with the number of grains per pod (0.485*) and yield (0.883**) as shown in Table 3. An increase of 308.7% over the lowest value of this variable and 12.52% over the control was recorded. Statistically significant differences between the treatments were observed, since a very strong detrimental effect of GA was seen in this variable. Figures 1A and 1B show the positive interaction of TRIA with the mixture of AUX, CK and GA, but a negative interaction with AUX and CK applied separately.

Regarding the number of grains per pod, significant statistical differences were observed between the treatments. The TRIA+AUX+GA+CK showed an increase of 48.8% over the control and 14.3% over the TRIA applied alone, which, in turn, represents a 22.6% increase over the control.




FIGURE 1. Differential effects of the application of A) triacontanol (TRIA) compared to the control and B) the plant growth regulators (AUX - Auxins, GA - Gibberellins, and CK - Cytokinins) alone or mixed compared to triacontanol. PH - plant height; NBP - number of branches per plant; PL - pod length; PW - pod width; NPP - number of pods per plant; NGP - number of grains per pod; PFW - plant fresh weight.




FIGURE 2. Differential effect of the treatments on the yield ($t\ ha^{-1}$) compared to the control. TRIA - triacontanol, AUX - auxins, GA - gibberellins, CK - cytokinins, K - foliar-applied potassium.

TRIA+AUX, TRIA+GA and the control did not exceed the average value obtained in the experiment of seven grains per pod (Tab. 3).

Within the yield components, the number of pods per plant with the application of TRIA decreased by 3.8% compared to the control (Fig. 1A). No positive interactions were observed with the mixture of AUX and CK or with the mixture of the three hormones. However, a negative interaction was recorded with AUX and CK applied separately since the AUX-CK relationship is imbalanced in physiological processes. A large decrease was observed with GA since they favor vegetative growth (Davies, 2010), and foliar potassium application obtained values below the control (32.5% less).

FIGURE 3. Differential effects of growth regulators on the yield ($t\ ha^{-1}$) compared to triacontanol. AUX - auxins, GA - gibberellins, CK - cytokinins, K - foliar-applied potassium.

The number of pods per plant is one of the most important yield components in legumes. GA decreased this yield variable because it promotes growth to the detriment of fruiting and formation of plant flowers (Davies, 2010). CK promote pod formation and show synergism with auxins (Cato *et al.*, 2013). The positive impact of CK has been observed in plants, such as *Artemisia* grown *in vitro* (Rasool *et al.*, 2013), showing a greater effect with the mixture of the three growth regulators (Cato *et al.*, 2013).

Foliar-applied potassium has a positive effect on the control and an interaction with TRIA, similar to that occurring with CK. Regarding the variable green pod yield, the values obtained ranged between 2.15 and 12.15 $t\ ha^{-1}$ with highly significant statistical differences between treatments ($P < 0.01$). TRIA+AUX+CK and TRIA+AUX+GA+CK

TABLE 4. Pearson's correlation coefficients of the variables tested.

	PH	NBP	NPP	PL	PW	PFW	NGP
NBP	-0.271	----	----	----	----	----	----
NPP	-0.573**	0.655**	----	----	----	----	----
PL	-0.443*	0.442*	0.396	----	----	----	----
PW	0.225	0.243	-0.232	0.131	----	----	----
PFW	-0.255	0.891**	0.670**	0.626**	0.267	----	----
NGP	-0.447*	0.456*	0.485*	0.823**	0.058	0.587**	----
Yield	-0.613**	0.731**	0.883**	0.654**	-0.009	0.845**	0.723**

PH - plant height; NBP - number of branches per plant; NPP - number of pods per plant; PL - pod length; PW - pod width; PFW - plant fresh weight, and NGP - number of grains per pod.

exceeded 10 t ha⁻¹, and TRIA+GA showed a decrease of 73.1% relative to the control. TRIA+AUX and TRIA+K did not exceed the control. The application of TRIA alone showed an increase of 11.3% relative to the control (Fig. 2).

Pearson's correlation coefficients of the yield variables showed that the number of pods per plant was positively correlated with number of grains per pod (0.485*), number of branches per plant (0.655**), plant fresh weight (0.670**), and yield (0.883**), and negatively correlated with plant height (-0.573**). The number of grains per pod was also positively correlated with the other variables ($P < 0.05$). The same was observed for the variable green pod yield, except for pod width, which is not correlated with any variable (Tab. 4).

Green pod yield is the agronomic trait that helps to determine the profitability of the crop. It also determines if the crop is a good alternative for the farmer, who looks for high yields with increasing emphasis on quality (Espinosa & Ligarreto, 2005; Checa Coral *et al.*, 2017). The high yield values obtained are due to the fact that the values of number of branches per plant, number of pods per plant, and number of grains per pod were also higher. These last two variables are important components of the yield and determine the productivity of plants. The lowest values were obtained in the interaction of TRIA with AUX and GA applied separately. This shows the negative effect of GA on crop yield in general since they decrease pod formation favoring vegetative growth and increasing plant height, an unfavorable issue leading to lodging (Davies, 2010; Heden, 2016; Wang *et al.*, 2017). It also shows the negative effect of AUX applied alone, since they alter the AUX-CK relationship that is key to the growth and development of plant species (Schaller *et al.*, 2015).

TRIA is a natural growth regulator widely studied in various species, with notable effects on the growth and development of rice (Jadhav *et al.*, 2017), *Capsicum annuum* L.

(Sahu *et al.*, 2017), willow (Digruber *et al.*, 2018), sweet potato (Rajak *et al.*, 2018), strawberry (Baba *et al.*, 2017), and guava (Singh *et al.*, 2017). Pearson's correlation coefficients show a highly positive correlation between yield and the number of pods per plant (0.883**), number of branches per plant (0.731**), number of grains per plant (0.723**), plant fresh weight (0.845**), and pod length (0.654**) (Tab. 4).

The differential effect for TRIA in this experiment was greater, especially in the yield components, with an increase in plant height of 11%, a decrease in number of branches per plant and number of pods per plant of 5.9 and 3.8%, respectively, and increases of 9.5% in pod length, 4.2% in pod width, 11.19% in plant fresh weight and 22.6% in number of grains per pod.

Conclusions

The results of this study showed that TRIA by itself has no remarkable effect on the growth and yield of *Pisum sativum*; only when TRIA is combined with other plant growth regulators such as AUX, GA and CK are noteworthy effects seen. AUX, GA and CK increased morphological variables with the exception of the number of branches per plant, and pod length in the case of GA. Within the yield components, AUX and CK increased the number of grains per pod and GA decreased all these variables, showing a negative interaction with TRIA. AUX and CK acted synergistically in almost all the variables evaluated, except for the number of branches for which similar values were obtained. In all these variables, the synergistic action of AUX and CK exceeds the effect of each growth regulator separately. The joint application of foliar-applied potassium fertilizer with TRIA showed an unfavorable effect on the yield and plant fresh weight. The highest green pod yields were obtained with the application of TRIA plus AUX, GA and CK (12.15 t ha⁻¹), followed by the application of TRIA plus AUX and CK (10.53 t ha⁻¹). The trihormonal application with TRIA obtained the best results in the variables

because the hormonal relationship is maintained and is not altered by the imbalance of the others.

Acknowledgments

The authors acknowledge the help of the Grain Legumes and Oilseed Research Center (Peru) for financing the research, Engineer Norma Bustamante and the company FARMEX S.A. for providing the biostimulants used in the experiment, and the National Institute of Agrarian Innovation (INIA-PERU) for facilitating the experiment site.

Conflict of interest statement

The authors declare that there is no conflict of interest regarding the publication of this article.

Author's contributions

AHJ obtained the financial support and the experiment site for the project. AHJ and HCS designed the methodology. HCS conducted the research process, performed the experiment, and analyzed the data. HCS wrote the draft and AHJ carried out the revision of the manuscript.

Literature cited

- Aftab, T., Khan, M. M. A., Idrees, M., Naeem, M., Singh, M., & Ram, M. (2010). Stimulation of crop productivity, photosynthesis and artemisinin production in *Artemisia annua* L. by triacontanol and gibberellic acid application. *Journal of Plant Interactions*, 5(4), 273–281. <https://doi.org/10.1080/17429141003647137>
- Anchivilca Rojas, G. H. (2018). *Abonamiento orgánico y fertilización NPK en arveja verde (Pisum sativum L.) cv. Rondo, bajo riego por goteo en Tupicocha, Huarochirí* [Undergraduate honors thesis, Universidad Nacional Agraria La Molina]. Repositorio Institucional Universidad Nacional Agraria La Molina. <http://repositorio.lamolina.edu.pe/handle/UNALM/3559>
- Baba, T. R., Ali, A., Kumar, A., & Husain, M. (2017). Effect of exogenous application of salicylic acid and triacontanol on growth characters and yield of strawberry. *The Pharma Innovation Journal*, 6(11), 274–279.
- Bertolin, D. C., Sá, M. E., Arf, O., Furlani Junior, E., Colombo, A. S., & Carvalho, F. L. B. M. (2010). Aumento da produtividade de soja com a aplicação de bioestimulantes. *Bragantia*, 69(2), 339–347. <https://doi.org/10.1590/S0006-87052010000200011>
- Borowski, E., Blamowski, Z. K., & Michałek, W. (2000). Effects of Tomatex/Triacontanol/ on chlorophyll fluorescence and tomato (*Lycopersicon esculentum* Mill.) yields. *Acta Physiologiae Plantarum*, 22, 271–274. <https://doi.org/10.1007/s11738-000-0030-5>
- Burt, R. (Ed.). (2014). *Soil survey field and laboratory methods manual. Soil survey investigations report no. 51, Version 2*. United States Department of Agriculture, Natural Resources Conservation Service.
- Carlson, D. R., Dyer, D. J., Cotterman, C. D., & Durley, R. C. (1987). The physiological basis for cytokinin induced increases in pod set in IX93-100 soybeans. *Plant Physiology*, 84(2), 233–239. <https://doi.org/10.1104/pp.84.2.233>
- Cato, S. C., Macedo, W. R., Peres, L. E. P., & Castro, P. R. C. (2013). Sinergism among auxins, gibberellins and cytokinins in tomato cv. Micro-Tom. *Horticultura Brasileira*, 31, 549–553. <https://doi.org/10.1590/S0102-05362013000400007>
- Checa Coral, Ó. E., Bastidas Acosta, J. E., & Narváez Taimal, O. C. (2017). Evaluación agronómica y económica de arveja arbustiva (*Pisum sativum* L.) en diferentes épocas de siembra y sistemas de tutorado. *Revista U.D.C.A Actualidad & Divulgación Científica*, 20(2), 279–288.
- Checa Coral, O. E., Getial Pantoja, J. A., & Rodríguez Rodríguez, D. M. (2020). Evaluación de ocho líneas de arveja arbustiva (*Pisum sativum* L.) en seis ambientes de la zona cerealista de Nariño. *Revista U.D.C.A Actualidad & Divulgación Científica*, 23(1), Article e1211. <https://doi.org/10.31910/rudca.v23.n1.2020.1211>
- Chen, X., Yuan, H., Chen, R., Zhu, L., Du, B., Weng, Q., & He, G. (2002). Isolation and characterization of triacontanol-regulated genes in rice (*Oryza sativa* L.): possible role of triacontanol as a plant growth stimulator. *Plant Cell Physiology*, 43(8), 869–876. <https://doi.org/10.1093/pcp/pcf100>
- Csukasi, F., Merchante, C., & Valpuesta, V. (2009). Modification of plant hormone levels and signaling as a tool in plant biotechnology. *Biotechnology Journal*, 4(9), 1293–1304. <https://doi.org/10.1002/biot.200800286>
- Davies, P. J. (2010). The plant hormones: their nature, occurrence, and functions. In P. J. Davies (Ed.), *Plant hormones: biosynthesis, signal transduction, action!* (pp. 1–15). Springer. https://doi.org/10.1007/978-1-4020-2686-7_1
- Depuydt, S., & Hardtke, C. S. (2011). Hormone signalling crosstalk in plant growth regulation. *Current Biology*, 21(9), R365–R373. <https://doi.org/10.1016/j.cub.2011.03.013>
- Digruber, T., Sass, L., Cseri, A., Paul, K., Nagy, A. V., Remenyik, J., Molnár, I., Vass, I., Toldi, O., Gyuricza, C., & Dudits, D. (2018). Stimulation of energy willow biomass with triacontanol and seaweed extract. *Industrial Crops and Products*, 120, 104–112. <https://doi.org/10.1016/j.indcrop.2018.04.047>
- Erisman, J. W. (2011). The new global nitrogen cycle. In J. C. Polacco, & C. D. Todd (Eds.), *Ecological aspects of nitrogen metabolism in plants* (1st ed., pp. 3–15). Wiley-Blackwell. <https://doi.org/10.1002/9780470959404.ch1>
- Espinosa, N., & Ligarreto, G. A. (2005). Evaluación de la habilidad combinatoria y heterosis de siete progenitores de arveja *Pisum sativum* L. *Agronomía Colombiana*, 23(2), 197–206.
- Gomez-Roldan, V., Fermas, S., Brewer, P. B., Puech-Pagès, V., Dun, E. A., Pillot, J. P., Letisse, F., Matusova, R., Danoun, S., Portais, J. C., Bouwmeester, H., Bécard, G., Beveridge, C. A., Rameau, C., & Rochange, S. F. (2008). Strigolactone inhibition of shoot branching. *Nature*, 455, 189–194. <https://doi.org/10.1038/nature07271>
- Hardtke, C. S., Dorcey, E., Osmont, K. S., & Sibout, R. (2007). Phytohormone collaboration: zooming in on auxin-brassinosteroid interactions. *Trends in Cell Biology*, 17(10), 485–492. <https://doi.org/10.1016/j.tcb.2007.08.003>
- Hedden, P. (2016). Gibberellins. In B. Thomas, B. G. Murray, & D. J. Murphy (Eds.), *Encyclopedia of applied plant sciences*

- (Vol. 1, pp. 411–420). Academic Press. <https://doi.org/10.1016/B978-0-12-394807-6.00219-7>
- Henry, E. W., & Gordon, C. J. (1980). The effect of triacontanol on peroxidase, IAA, and plant growth in *Pisum sativum* var. 'Alaska' and 'Little marvel'. *Journal of Experimental Botany*, 31(5), 1297–1303. <https://doi.org/10.1093/jxb/31.5.1297>
- Jadhav, S. M., Burondkar, M. M., Patil, M. Y., & Sawant, G. B. (2017). Effect of plant growth regulators on physiological behavior of rice (*Oryza sativa* L.) under *kharif* Konkan condition. *International Journal of Chemical Studies*, 5(6), 1365–1367.
- Khan, M. M. A., Bhardwaj, G., Naeem, M., Moinuddin, Mohammad, F., Singh, M., Nasir, S., & Idrees, M. (2009). Response of tomato (*Solanum lycopersicum* L.) to application of potassium and triacontanol. *Acta Horticulturae*, 823, 199–208. <https://doi.org/10.17660/actahortic.2009.823.29>
- Khandaker, M. M., Faruq, G., Rahman, M. M., Sofian-Azirun, M., & Boyce, A. N. (2013). The influence of 1-triacontanol on the growth, flowering, and quality of potted *Bougainvillea* plants (*Bougainvillea glabra* var. "Elizabeth Angus") under natural conditions. *The Scientific World Journal*, 2013, Article 308651. <https://doi.org/10.1155/2013/308651>
- Kuppusamy, K. T., Walcher, C. L., & Nemhauser, J. L. (2009). Cross-regulatory mechanisms in hormone signaling. *Plant Molecular Biology*, 69, Article 375. <https://doi.org/10.1007/s11103-008-9389-2>
- Liu, C., Wang, S., Copeland, L., & Wang, S. (2015). Physicochemical properties and *in vitro* digestibility of starches from field peas grown in China. *LWT - Food Science and Technology*, 64(2), 829–836. <https://doi.org/10.1016/j.lwt.2015.06.060>
- Marschner, P. (Ed.). (2011). *Mineral nutrition of higher plants* (3rd ed.). Academic Press. <https://doi.org/10.1016/C2009-0-63043-9>
- Martínez González, L., Maqueira López, L., Nápoles García, M. C., & Núñez Vázquez, M. (2017). Efecto de bioestimulantes en el rendimiento de dos cultivares de frijol (*Phaseolus vulgaris* L.) biofertilizados. *Cultivos Tropicales*, 38(2), 113–118.
- Mosjidis, C. O., Peterson, C. M., Truelove, B., & Dute, R. R. (1993). Stimulation of pod and ovule growth of soybean, *Glycine max* (L.) Merr. by 6-benzylaminopurine. *Annals of Botany*, 71(3), 193–199. <https://doi.org/10.1006/anbo.1993.1024>
- Müller, D., & Leyser, O. (2011). Auxin, cytokinin and the control of shoot branching. *Annals of Botany*, 107(7), 1203–1212. <https://doi.org/10.1093/aob/mcr069>
- Naeem, M., Idrees, M., Aftab, T., Khan, M. M. A., & Moinuddin. (2010). Changes in photosynthesis, enzyme activities and production of anthraquinone and sennoside content of coffee senna (*Senna occidentalis* L.) by triacontanol. *International Journal of Plant Developmental Biology*, 4(1), 53–59.
- Naeem, M., Khan, M. M. A., & Moinuddin. (2012). Triacontanol: a potent plant growth regulator in agriculture. *Journal of Plant Interactions*, 7(2), 129–142. <https://doi.org/10.1080/17429145.2011.619281>
- Naeem, M., Khan, M. M. A., Moinuddin, Idrees, M., & Aftab, T. (2011). Triacontanol-mediated regulation of growth and other physiological attributes, active constituents and yield of *Mentha arvensis* L. *Plant Growth Regulation*, 65, 195–206. <https://doi.org/10.1007/s10725-011-9588-8>
- Naeem, M., Khan, M. M. A., Moinuddin, & Siddiqui, M. H. (2009). Triacontanol stimulates nitrogen-fixation, enzyme activities, photosynthesis, crop productivity and quality of hyacinth bean (*Lablab purpureus* L.). *Scientia Horticulturae*, 121(4), 389–396. <https://doi.org/10.1016/j.scienta.2009.02.030>
- Nakaya, M., Tsukaya, H., Murakami, N., & Kato, M. (2002). Brassinosteroids control the proliferation of leaf cells of *Arabidopsis thaliana*. *Plant & Cell Physiology*, 43(2), 239–244. <https://doi.org/10.1093/pcp/pcf024>
- Pozo, M. J., López-Ráez, J. A., Azcón-Aguilar, C., & García-Garrido, J. M. (2015). Phytohormones as integrators of environmental signals in the regulation of mycorrhizal symbioses. *New Phytologist*, 205(4), 1431–1436. <https://doi.org/10.1111/nph.13252>
- Rajak, R., Sengupta, S., Tiwary, A. K., Verma, S., Minz, A., Bibhuti, T. O. P., & Rani, V. (2018). Effect of bio regulators on yield parameters of sweet potato (*Ipomoea batatas* L.). *Journal of Pharmacognosy and Phytochemistry*, 7(1), 2425–2427.
- Rasool, R., Ganai, B. A., Kamili, A. N., Akbar, S., & Masood, A. (2013). Synergistic effect of auxins and cytokinins on propagation of *Artemisia amygdalina* (Asteraceae), a critically endangered plant of Kashmir. *Pakistan Journal of Botany*, 45(2), 629–634.
- Ratnayake, W. S., Hoover, R., Shahidi, F., Perera, C., & Jane, J. (2001). Composition, molecular structure, and physicochemical properties of starches from four field pea (*Pisum sativum* L.) cultivars. *Food Chemistry*, 74(2), 189–202.
- Rodríguez Quispe, G. (2015). *Evaluación de 12 cultivares de arveja (Pisum sativum L.) de tipo industrial para cosecha en verde en condiciones de Tarma* [Undergraduate thesis, Universidad Nacional del Centro del Perú]. Repositorio Institucional Digital Universidad Nacional del Centro del Perú. <http://repositorio.uncp.edu.pe/bitstream/handle/20.500.12894/977/RODR%C3%8DGUEZ%20QUISPE%20GUSTAVO.pdf>
- Rondinel Ruíz, R. (2014). *Rendimiento en vaina verde de tres variedades de arveja (Pisum sativum L.) en tres modalidades de siembra bajo el sistema de agricultura de conservación. Canaán a 2750 msnm - Ayacucho* [Undergraduate honors thesis, Universidad Nacional de San Cristóbal de Huamanga]. Repositorio Institucional Universidad Nacional de San Cristóbal de Huamanga. http://repositorio.unsch.edu.pe/bitstream/handle/UNSCH/901/Tesis_Ag1139_Ron.pdf
- Sahu, G., Aslam, T., Das, S. P., Maity, T. K., & Gupta, N. K. (2017). A study on pre-flowering foliar spray of plant growth regulator on growth and yield parameters in sweet pepper (*Capsicum annuum* L.) under protected condition. *International Journal of Current Microbiology and Applied Sciences*, 6(7), 3998–4007. <https://doi.org/10.20546/ijcmas.2017.607.414>
- Santner, A., Calderon-Villalobos, L. I. A., & Estelle, M. (2009). Plant hormones are versatile chemical regulators of plant growth. *Nature Chemical Biology*, 5, 301–307. <https://doi.org/10.1038/nchembio.165>
- Santos, J., Herrero, M., Mendiola, J. A., Oliva-Teles, M. T., Ibáñez, E., Delerue-Matos, C., & Oliveira, M. B. P. P. (2014). Assessment of nutritional and metabolic profiles of pea shoots: the new ready-to-eat baby-leaf vegetable. *Food Research International*, 58, 105–111. <https://doi.org/10.1016/j.foodres.2014.01.062>

- Santos, O. F., Cunha, F. F., Taira, T. L., Souza, E. J., & Leal, A. J. F. (2018). Increase in pea productivity associated with irrigation management. *Horticultura Brasileira*, 36(2), 178–183. <https://doi.org/10.1590/s0102-053620180205>
- Schaller, G. E., Bishopp, A., & Kieber, J. J. (2015). The Yin-Yang of hormones: cytokinin and auxin interactions in plant development. *The Plant Cell*, 27(1), 44–63. <https://doi.org/10.1105/tpc.114.133595>
- Shukla, A., Abad Farooqi, A. H., Shukla, Y. N., & Sharma, S. (1992). Effect of triacontanol and chlormequat on growth, plant hormones and artemisinin yield in *Artemisia annua* L. *Plant Growth Regulation*, 11, 165–171. <https://doi.org/10.1007/BF00024071>
- Singh, K., Sharma, M., & Singh, S. K. (2017). Effect of plant growth regulators on fruit yield and quality of guava (*Psidium guajava*) cv. Allahabad Safeda. *Journal of Pure and Applied Microbiology*, 11(2), 1149–1154. <https://doi.org/10.22207/JPAM.11.2.61>
- Taiz, L., Zeiger, E., Møller, I. M., & Murphy, A. (Eds.). (2014). *Plant physiology and development* (6th ed.). Oxford University Press.
- Wang, Y., Zhao, J., Lu, W., & Deng, D. (2017). Gibberellin in plant height control: old player, new story. *Plant Cell Reports*, 36, 391–398. <https://doi.org/10.1007/s00299-017-2104-5>
- Weeden, N. F. (2007). Genetic changes accompanying the domestication of *Pisum sativum*: is there a common genetic basis to the ‘domestication syndrome’ for legumes? *Annals of Botany*, 100(5), 1017–1025. <https://doi.org/10.1093/aob/mcm122>
- Wehner, T., & Gritton, E. T. (1981). Effect of the *n* gene on pea pod characteristics. *Journal of the American Society for Horticultural Science*, 106(2), 181–183.
- Wolters, H., & Jürgens, G. (2009). Survival of the flexible: hormonal growth control and adaptation in plant development. *Nature Reviews Genetics*, 10, 305–317. <https://doi.org/10.1038/nrg2558>
- Wu, L., Chang, K. F., Hwang, S. F., Conner, R., Fredua-Agyeman, R., Feindel, D., & Strelkov, S. E. (2019). Evaluation of host resistance and fungicide application as tools for the management of root rot of field pea caused by *Aphanomyces euteiches*. *The Crop Journal*, 7(1), 38–48. <https://doi.org/10.1016/j.cj.2018.07.005>