

Concordance of estimated glomerular filtration rate according to the formulas used in Colombia for patients with chronic kidney disease not on dialysis

JORGE CORONADO-DAZA, MARÍA CAROLINA FRAGOZO-RAMOS, LINA FERNANDA REYES-FONTALVO
• CARTAGENA (COLOMBIA)

DOI: <https://doi.org/10.36104/amc.2020.1652>

Abstract

Introduction: the Colombian guidelines for chronic kidney disease (CKD) recommend estimating glomerular filtration (GF) using the Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) equation. No studies have been performed in the Colombian population to compare the accuracy of this equation to that of others used in clinical practice.

Design and methods: we evaluated the GF estimation performance of the Modification of Diet in Renal Disease (MDRD-4), Cockcroft-Gault (CG) and body surface area adjusted Cockcroft-Gault (CG-BSA) equations against the CKD-EPI equation in 757 adult patients. Performance was evaluated using bias, precision and accuracy measurements.

Results: the mean GF by CKD-EPI was 37.32 ± 12.71 mL/min/1.73m²; by MDRD-4 it was 39.8 ± 13.2 mL/min/1.73m², by CG it was 35 ± 12.6 mL/min and by CG-BSA it was 34.52 ± 11.34 mL/min/1.73m². All the equations had bias with respect to GF by CKD-EPI. The most accurate equation was GF estimated by MDRD-4 (MeGF) with 97.1 and 99.74% of measurements within 15 and 30%, respectively; and the least accurate was GF estimated by CG (CGeGF) with 59.7 and 81.77% of the measurements within 15 and 30%, respectively. The concordance correlation coefficient between GF by CKD-EPI and MDRD-4 was 0.97, with CG and CG-BSA at 0.78 and 0.85, respectively.

Conclusions: the most accurate alternative equation for estimating glomerular filtration in this Colombian population is MDRD-4, which has a high concordance with the CKD-EPI equation. Estimation of GF with the CG equation is not recommended. (Acta Med Colomb 2020; 45. DOI: <https://doi.org/10.36104/amc.2020.1652>).

Key Words: *glomerular filtration rate, estimation equations, chronic kidney disease, renal function.*

Dr. Jorge Coronado-Daza: Internista-Nefrólogo. MSc, PhD. Profesor Asociado Departamento Médico, Grupos de Investigación "Alta Tensión" Universidad de Cartagena y "Nefrología ASOCOLNEF"; Dra. María Carolina Fragozo-Ramos: Médico Internista; Dra. Lina Fernanda Reyes-Fontalvo: Médico Interno. **Grupo de Investigación "Alta Tensión" Universidad de Cartagena.** Cartagena (Colombia).
Correspondencia: Dr. Jorge Coronado-Daza, Cartagena (Colombia).
E-mail: jocodada@yahoo.es
Received: 07/II/2020 Accepted: 12/VIII/2020

Introduction

Estimated glomerular filtration (GF) is an important tool for the clinician, routinely used to identify chronic kidney disease (CKD), monitor kidney function changes and adjust pharmacological treatment in some patients (1-5). There are more precise methods which show real glomerular filtration values, such as clearance of inulin and various isotopes (6, 7). Due to the complexity and cost of these techniques, they have been replaced in clinical practice by equations for estimating the GF rate (GFR).

At this time, it has become necessary to determine which of the equations for estimating GFR have the greatest precision for classifying kidney patients. Worldwide, the concordance between the Cockcroft-Gault (CG), four-variable Modification of Diet in Renal Disease (MDRD-4) and Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) formulas has

been established, comparing them with direct measurement based on iothalamate (8). Ever since the CKD-EPI formula was published, the performance of the equations has been compared in various types of populations, finding that the CKD-EPI formula is the most accurate (9-11). Thus, in the KDIGO 2012 guidelines, this formula is used as the basis for classifying patients with CKD (12).

The Colombian 2016 CKD guidelines (13), an adoption of the KDIGO guidelines, have taken CKD-EPI as the formula for estimating glomerular filtration. However, since 2008, when the High Cost Account (CAC in Spanish) was established in Colombia, the equation that has been used to date for defining patients with CKD is CG-SC (14). The reason for using the CG formula is the fact that it is easy to run manually and does not require internet availability, which is needed for the MDRD-4 or CKD-EPI formulas.

The clinicians' feeling is that, in following the CAC guidelines, patients with CKD are being mistakenly classified, since the CG-SC formula is used for all patients without considering that it cannot be used in patients at the extremes of age and weight, those with muscle disease, amputees and/or vegetarians (15, 16). The objective of our study is to determine the concordance of GFR measured according to the CG, CG-BSA, MDRD-4 and CKD-EPI formulas in patients with Stage 3a (G3a) to Stage 5 non-dialysis (G5ND) CKD.

Materials and methods

Study population

This was an observational study of individuals seen as outpatients in the nephrology department of a healthcare institution in Cartagena, Colombia, during 2016 and 2017.

The inclusion criteria were: 1) Adult patients over the age of 18 being followed for G3a-G5ND CKD; and 2) Patients with glomerular filtration calculated using the CG (CGeGF), body surface area-adjusted CG (CG-BSAeGF), MDRD-4 and CKD-EPI formulas. 2

The following patients were excluded: pregnant women, lower-limb amputees, vegetarians, and those with active cancer, or primary or secondary muscle disease. Data was gathered on type of health insurance, age, sex, weight, height, body mass index (BMI), etiology of CKD and albuminuria and/or proteinuria.

Glomerular filtration rate estimation

The GFR was calculated in all cases using the CG (mL/min), CG-BSA, MDRD-4 and CKD-EPI (mL/min/1.73 m²) equations, as described below:

- GF estimated by Cockcroft-Gault (CGeGF) (15):
 - Creatinine clearance in men = $(140 - \text{age}) \times (\text{weight} / \text{serum creatinine} \times 72)$.
 - Creatinine clearance in women = $(140 - \text{age}) \times (\text{weight} \times 0.85 / \text{serum creatinine} \times 72)$.
- GF estimated by body surface area-adjusted Cockcroft-Gault (CG-BSAeGF) (15):
 - Men = $(140 - \text{age}) \times (\text{weight}) \times (1.73 \text{ m}^2 / \text{body surface area}) / (\text{serum creatinine} \times 72)$.
 - Women = $(140 - \text{age}) \times (\text{weight}) \times (1.73 \text{ m}^2 / \text{body surface area}) \times 0.85 / (\text{serum creatinine} \times 72)$.

Body surface area was calculated using the DuBois formula (17): $\text{body surface area (m}^2\text{)} = 0.20247 \times \text{height (1)}^{0.725} \times \text{weight (kg)}^{0.425}$
- GF estimated by four-variable MDRD (MeGF) (18):
 - $175 \times (\text{serum creatinine} / 0.95) - 1.154 \times (\text{age}) - 0.2030 \times (0.742 \text{ if female}) \times (1.210 \text{ if black race})$.
- GF estimated by CDK-EPI (CKD-EPIeGF) (19):
 - $141 \times \min \{ \text{creatinine} / k, 1 \}^\alpha \times \max \{ \text{creatinine} / k, 1 \}^{-1.209} \times 0.993 \text{ age [years]} \times 1.018 \text{ [if female]}$, where k is 0.7 for women and 0.9 for men, α is -0.329 for women and -0.411 for men.

The units used in the equations were age in years, weight

in kilograms (kg), serum creatinine (Cr) concentration in mg/dL and body surface area (BSA) in m².

Categories according to GFR

Reduced GF, defined as eGFR <60 mL/min/1.73 m² was classified in four groups according to the 2012 KDIGO guidelines (12): 45-59 mL/min/1.73 m² (G3a), 30-44 mL/min/1.73 m² (G3b), 15-29 mL/min/1.73 m² (G4) and <15 mL/min/1.73 m² (G5). Patients in category G5 under conservative management without dialysis were classified as stage G5ND. Prevalence for each category was described according to the study equations.

Statistical analysis

Descriptive statistics were applied for categorical and quantitative variables, expressing them as frequencies, percentages, mean and median according to the nature of each variable.

The concordance correlation coefficient (CCC) (20) was calculated for all patients between the GFR estimated by CKD-EPI and the GFR estimated by other formulas: MDRD-4, CGeGF and CG-BSAeGF. Concordance was classified according to the CCC as follows: 1=perfect; >0.99=almost perfect; 0.95 to 0.99=substantial; 0.900 to 0.949=moderate; and <0.900=poor.

Glomerular filtration estimated by the CKD-EPI formula was taken as the gold standard (eGFGS) for this study. The eGFGS was compared with MeGF, CGeGF and CG-BSAeGF using the Bland-Altman analysis (21) (a graphical representation of the regression line produced by using the difference in means of the formulas being compared as the dependent variable and the average of the means as the independent variable). Two standard deviations around the mean difference were taken as the limits of agreement between both measurements, as they include 95% of the observed differences. A maximum limit of agreement of ± 5 was established between the estimation formulas and the eGFGS (CKD-EPI), considering the definition of accelerated CKD progression to be a sustained decline in GF greater than 5 mL/min/year.

Bias, precision and accuracy were evaluated among the compared formulas. Bias was assessed as the mean difference in means between the eGFGS and the MeGF, CGeGF and CG-BSAeGF. Precision was defined as the range of the limits of agreement of the mean difference in means (bias) between the eGFGS and the MeGF, CGeGF and CG-BSAeGF estimates. Accuracy was measured as the percentage of results estimated with the different formulas within 15% (P15) and 30% (P30) of the measurements using the eGFGS. A P30 >90% qualifies a formula as satisfactory for clinical interpretation (22). The best formula was defined as the one with the least bias, smallest range of limits of agreement (or best precision) and highest P30. Stata 14 and XLSTAT Premium statistical software were used for analysis.

Ethical aspects

The ethical principles were in accordance with Colombia's Resolution 8430 of 1993. As it was a non-intervention study, with minimal risk, written informed consent was not requested, only verbal consent. The study was approved by the healthcare institution. Data confidentiality, integrity and transparency were respected.

Results

Data were collected from 855 patients, of whom 757 were eligible for the final analysis, having met the inclusion criteria (Figure 1).

Of the subjects analyzed, 61.96% were men and 38% were women; 50.53% had contributory health insurance and 48.68% subsidized. Arterial hypertension was the main cause of CKD (61.43%), followed by diabetes mellitus (31.04%); primary glomerulopathies and other causes made up the remaining 7.53%. The main characteristics of the included individuals are shown in Table 1.

Overweight was found in 37.12% of cases, and obesity in 11.2%. The frequency of the CKD stages was different for each of the formulas used (Figure 2). The CG-BSA formula classified a greater percentage of cases as G3b CKD than the MDRD-4 and CKD-EPI formulas. In turn, there was a greater proportion of patients in G4 CKD with CG and CG-BSA: 32.23 and 30.65%, respectively.

Serum albumin was measured in 508 patients (mean 4.0 ± 0.5 g/dL); 90.55% had normal levels and the remaining 9.48% had hypoalbuminemia. The average albuminuria was 481.58 ± 305.73 mg/gr (range: 1-5,009 mg/gr); 53.77% of the albuminuria measurements were in category A1 (<30 mg/gr), 32.43% in category A2 (30-300 mg/gr), and 13.8% in category A3 (>300 mg/gr).

The performance of the various formulas in estimating GFR is shown in Table 2. All the formulas had a small bias with respect to GF by CKD-EPI. The bias was negative for CGeGF and CG-BSAeGF (-2.23 [-2.82 to -1.65] and -2.8 [-3.24 to -2.37] mL/min/1.73m², respectively) and positive for MeGF (2.49 [2.32 to 2.65] mL/min/1.73m²). Precision was similar between CGeGF and CG-BSAeGF (-18.2 to 13.7 and -14.7 to 9.1 mL/min/1.73m², respectively); the most precise was MeGF (-2.1 to 7.1 mL/min/1.73m²). The most accurate formula was MeGF, with a proportion of measurements within 15 and 30% of 97.1 and 99.74%, respectively, and the least accurate was CGeGF with 59.7 and 81.77% of the measurements within 15 and 30%, respectively. The concordance correlation coefficient between eGFGS and MeGF was 0.97, with 0.78 for CGeGF and 0.85 for CG-BSAeGF.

Figures 3-5 show the dispersion graphs for each equation and the differences between MeGF, CGeGF, CG-BSAeGF and eGFGS. The CGeGF Bland-Altman graph has a wide limit of agreement, with predominantly underestimated measurements, but predominantly overestimated G4 and G5 CKD measurements, which is more evident in the CG-BSAeGF graph. Conversely, the MeGF Bland-Altman

graph has a narrow limit of agreement with predominantly overestimated measurements, but with measurements around zero (0) in G4 and G5 CKD.

Discussion

All GFR estimation formulas should be validated in the population in which they are to be used. The current study is the first to attempt to validate the bias, precision, accuracy and concordance between each of the formulas for measuring GFR in the Colombian population, using GFR estimated with the CKD-EPI formula as the gold standard.

In our study, we found that the MDRD-4 formula had the best performance with respect to precision, accuracy and concordance relative to the CKD-EPI formula. The precision did not exceed the established limit of a range of 10 for limits of agreement (from -5 to +5), with precision being better at the eGFGS underestimation limit. The accuracy is almost perfect, and concordance is in the substantial range. On the other hand, GFR estimates using the CG and CG-BSA equations were widely dispersed with respect to the eGFGS, both with a range of limits of agreement over the established limit, with borderline accuracy for CG-BSAeGF alone, but poor concordance for both.

In Colombia, the CKD-EPI formula has not been validated, but it has been validated internationally, including the Latin American population. In 2014, a study of healthy Mexicans with an average age of 35 took direct GFR measurements obtained by an isotope or radiopharmaceutical (Tc-DTPA: Tc99m-DTPA [diethylenetriaminepentaacetic acid]) as a reference and subsequently compared them with GFR values estimated by CKD-EPI and MDRD-4. The




Figure 1. Study patient selection flowchart.

Table 1. General characteristics of the population.

Characteristic	Mean (N=757)	Standard deviation	95% CI	
Age (years)	75.36	12.16	65.3	79.4
Weight (kg)	67.3	12.42	66.7	77.8
Height (cm)	163.53	8.2	160.5	170.2
BMI (kg/m ²)	25.11	4.14	24.3	28.5
Creatinine	1.86	0.97	1.35	3.1
GFR by CG (mL/min)	35.08	12.59	27.69	42.97
GFR by BSA-corrected CG (mL/min/1.73 m ²)	34.52	11.34	26.66	41.17
GFR by MDRD-4 (mL/min/1.73 m ²)	39.81	13.20	26.99	47.34
GFR by CKD-EPI (mL/min/1.73 m ²)	37.31	12.71	25.72	45.11
Serum albumin (g/dL)	4.03	0.52	3.29	4.39
Creatinine clearance (mL/min)	44.57	19.85	36.14	63.52

BMI: body mass index; GFR: glomerular filtration rate; CG: Cockcroft-Gault; MDRD-4: four-variable Modification of Diet in Renal Disease; CKD-EPI: Chronic Kidney Disease Epidemiology Collaboration formula.

GFR calculated by CKD-EPI was found to have a greater concordance with the direct measurement reference values (23). Similar results were obtained by Veronese FV, et al. (24) in a study including healthy, diabetic and CKD patients. Despite having different population characteristics, the two studies demonstrated that CKD-EPI is a better predictor of GFR. However, in Colombia, the CAC categorizes GFR by CGeGF (14). Nevertheless, CGeGF's results cannot be generalized, given their high heterogeneity as BMI increases (25, 26), and its performance is inferior to GFR estimated by MDRD-4 and CKD-EPI formulas, as has been reported in the previously mentioned studies and others in various population groups (27, 28). These findings are consistent with our results.

Kuo CF et al., in a study of 32,542 healthy individuals undergoing routine studies, compared GFR estimated by six-variable MDRD (MDRD-6), MDRD adjusted to the Chinese population and CG. The estimation of GFR, CKD risk factor

assignment and stratification differed significantly between the three equations, especially with the CG equation. The CG formula produced a lower GFR and, paradoxically, higher measurements in patients with hyperuricemia, hypertension, and metabolic syndrome than in those without these risk factors. In this study, as well as in ours, the use of the CG formula for CKD screening is discouraged, and the MDRD formula adjusted to the Chinese population is recommended for that population (29).

On the other hand, the previous data contrast with results obtained by Liu X et al. who reported that the CG formula was the most precise for estimating GFR at different stages of chronic kidney disease in older adults (30). Hence the importance of our study to validate the CG and MDRD-4 formulas in the Colombian population, using the globally recommended CKD-EPI formula as reference, which has also been accepted in our country, according to the guidelines adopted in 2016. In the study population, we found that the performance of MDRD-4 was statistically better in all the study variables with respect to CGeGF and CG-BSAeGF; it was also more accurate in estimating GFR. Similar to our




Figure 2. Stages of Chronic Kidney Disease according to the different formulas for calculating glomerular filtration rate. CG: Cockcroft-Gault; CG-BSA: Cockcroft-Gault corrected for body surface area; MDRD-4: four-variable Modification of Diet in Renal Disease; CKD-EPI: Chronic Kidney Disease Epidemiology Collaboration formula; ND: not on dialysis.

Table 2. Performance of the formulas in calculating glomerular filtration rate compared with CKD-EPI.

	Bias	Precision		Accuracy
	*	LL	UL	P30 (%)
CGeGF-eGFGS	-2.23	-18.21	13.74	81.77
C G - B S A e G F - eGFGS	-2.8	-14.75	9.14	90.80
MeGF-eGFGS	2.49	-2.12	7.09	99.74

CGeGF: Cockcroft-Gault-estimated glomerular filtration;
 CG-BSAeGF: glomerular filtration estimated by the body surface area-corrected Cockcroft-Gault formula;
 MeGF: glomerular filtration by the MDRD-4 formula;
 *(mL/min/1.73 m²);
 LL: lower limit (mL/min/1.73 m²);
 UL: upper limit (mL/min/1.73 m²)




Figure 3. Bland-Altman concordance between CGeGF and CKD-EPIeGF. CG: Cockcroft-Gault; CKD-EPI: Chronic Kidney Disease Epidemiology Collaboration formula; CGeGF: CG-estimated glomerular filtration; CKD-EPIeGF: glomerular filtration by CKD-EPI.




Figure 4. Bland-Altman concordance between CG-BSAeGF and CKD-EPIeGF. CG-BSA: Cockcroft-Gault corrected for body surface area; CKD-EPI: Chronic Kidney Disease Epidemiology Collaboration formula; CG-BSAeGF: glomerular filtration estimated by CG-BSA; CKD-EPIeGF: glomerular filtration estimated by CKD-EPI.




Figure 5. Bland-Altman concordance between MeGF and CKD-EPIeGF. MDRD-4: four-variable Modification of Diet in Renal Disease; CKD-EPI: Chronic Kidney Disease Epidemiology Collaboration formula; MeGF: glomerular filtration estimated by MDRD-4; CKD-EPIeGF: glomerular filtration estimated by CKD-EPI.

results, the data obtained by Schwandt A, et al. (31) concur in that the MDRD-4 formula performed better globally, and had greater precision and less bias. In this last study, the parameter used as the gold standard was GFR estimated by 24-hour urine creatinine clearance.

Chudek J, et al. (32) also compared the performance of the MDRD-4, MDRD-6 and CG equations specifically in patients 65 years old and older. The MDRD-6 formula was found to have the best concordance, precision and accuracy, compared to CG, for all CKD groups. Related results in favor of the precision of MDRD in elderly patients were demonstrated in another study performed by Aras S et al. (33).

Finally, Zaman SB (34), in a study of 4,042 type 2 diabetic patients, reported a correlation coefficient between GFRs estimated by CKD-EPI and MDRD-4 of 0.90 (95% CI 0.83 - 0.97, $p < 0.001$). However, this study found that the correlation coefficient between the CKD-EPI equation and CG was 0.56 (95% CI: 0.44 - 0.67, $p < 0.0001$), even lower than what was found in our caseload.

The current study has the strength of having been performed on a significant number of CKD patients not on dialysis, and comparing for the first time the eGFGS (CKD-EPI), globally recommended and adopted in Colombia in the 2016 CKD guidelines, with alternative estimation formulas used in Colombia: MDRD-4, and the one used by the CAC, CG-BSAeGF. Despite the fact that the data are solid and have clinical significance, reporting a global analysis of the data of patients in Cartagena, its weaknesses lie in not considering a stratified analysis according to CKD stage and patient subgroups of ages and average and extreme weights that can show similar findings.

Conclusion

Our findings suggest that the classification of CKD categories in Colombia should be based on the CKD-EPI formula, as is recommended internationally based on direct measurements with inulin and radiopharmaceuticals. Using the alternate CG-BSA formula is not advisable, despite its easy manual use, as this study has shown that it has low precision and accuracy, which could lead to a greater classification of elderly patients in this population in early stages, such as G3a CKD, with the corresponding consequences for the patient and the healthcare system.

Thus, the most precise, accurate and concordant alternate formula for estimating glomerular filtration in this Colombian population is MDRD-4, although online methods would be required for its calculation, just like CKD-EPI. The Cockcroft-Gault formula has low precision, borderline accuracy and poor concordance with the CKD-EPI formula. It is recommended that the CG-BSA formula for GFR estimation not be used to classify CKD in Colombia.

Referencias

1. Coresh J, Stevens LA. Kidney function estimating equations: where do we stand?. *Curr Opin Nephrol Hypertens*. 2006 May; **15**(3): 276-84.
2. Alaini A, Malhotra D, Rondon-Berrios H, Argyropoulos CP, Khitan ZJ,

- Raj DSC, et al. Establishing the presence or absence of chronic kidney disease: Uses and limitations of formulas estimating the glomerular filtration rate. *World J Methodol.* 2017 Sep 26; **7(3)**: 73-92.
3. Van Rijn MHC, Metzger M, Flamant M, Houllier P, Haymann J-P, Van den Brand JA, et al. Performance of creatinine-based equations for estimating glomerular filtration rate changes over time. *Nephrol Dial Transplant.* 2018;(September):1-9.
 4. Higdon EA, Kimmons LA, Jr BTD, Hudson JQ. Disagreement in Estimates of Kidney Function for Drug Dosing in Obese Inpatients. *J Pharm Pract.* 2019 Feb; **32(1)**: 41-47.
 5. Pérez Cabeza AI, Chinchurreta Capote PA, González Correa JA, Ruiz Mateas F, Rosas Cervantes G, Rivas Ruiz F, et al. Discrepancies between the use of MDRD-4 IDMS and CKD-EPI equations, instead of the Cockcroft-Gault equation, in the determination of the dosage of direct oral anticoagulants in patients with non-valvular atrial fibrillation. *Med Clin (Barc).* 2018 Feb 9; **150(3)**: 85-91.
 6. Miller WG, Jones GRD. Estimated Glomerular Filtration Rate; Laboratory Implementation and Current Global Status. *Adv Chronic Kidney Dis.* 2018 Jan; **25(1)**: 7-13.
 7. Hofman M, Binns D, Johnston V, Siva S, Thompson M, Eu P, et al. 68Ga-EDTA PET/CT imaging and plasma clearance for glomerular filtration rate quantification: comparison to conventional 51Cr-EDTA. *J Nucl Med.* 2015 Mar; **56(3)**: 405-9.
 8. Hall PM, Rolin H. Iothalamate clearance and its use in large-scale clinical trials. *Curr Opin Nephrol Hypertens.* 1995 Nov; **4(6)**: 510-3.
 9. Yamaguchi T, Higashihara E, Okegawa T, Miyazaki I, Nutahara K. Optimal equation for estimation of glomerular filtration rate in autosomal dominant polycystic kidney disease: influence of tolvaptan. *Clin Exp Nephrol.* 2018; **22(5)**: 1213-23.
 10. Cristelli MP, Cofán F, Rico N, Trullàs JC, Manzardo C, Agüero F, et al. Estimation of renal function by CKD-EPI versus MDRD in a cohort of HIV-infected patients : a cross-sectional analysis. 2017; 1-7.
 11. Dou Y, Sun X, Liu D, Zhang L, Xiao J, Cheng G, et al. Accuracy of glomerular filtration rate equations for chronic kidney disease patients at the G3a stage : a single - center cross - sectional study. *BMC Res Notes.* 2017; **2**: 1-5.
 12. KDIGO. 2012 Clinical practice guideline for the evaluation and management of chronic kidney disease. *Kidney Int* 2013; **3**: 1-150.
 13. Ministerio de protección Social. Guía de Práctica Clínica para el diagnóstico y tratamiento de la Enfermedad Renal Crónica (adopción). [Internet]. Bogotá; 2016 [citado: 15 de diciembre de 2019]. Disponible en: http://gpc.minsalud.gov.co/gpc_sites/Repositorio/Otros_conv/GPC_e_renal/GPC_enfermedad_renal_adopcion.pdf
 14. Cuenta de alto costo. Situación de la enfermedad renal crónica, la hipertensión arterial y la diabetes mellitus. [Internet] Bogotá; 2017 [citado: 15 de diciembre de 2019]. Disponible en: https://cuentadealtocosto.org/site/images/Publicaciones/2018/Libro_Situacion_ERC_en_Colombia_2017.pdf.
 15. Cockcroft DW, Gault MH. Prediction of creatinine clearance from serum creatinine. *Nephron.* 1976; **16(1)**: 31-41.
 16. Deskur-Smielecka E, Kotlinska-Lemieszek A, Chudek J, Wieczorowska-Tobis K. Assessment of renal function in geriatric palliative care patients – Comparison of creatinine-based estimation equations. *Clin Interv Aging.* 2017; **12**: 977-83.
 17. Du Bois D, Du Bois EF. A formula to estimate the approximate surface area if height and weight be known. 1916. *Nutrition.* 1989; **5(5)**: 303-11; discussion 12-3.
 18. Levey AS, Coresh J, Greene T, Stevens LA, Zhang YL, Hendriksen S, et al. Using Standardized Serum Creatinine Values in the Modification of Diet in Renal Disease Study Equation for Estimating Glomerular Filtration Rate. 2006; **145(4)**: 247-54.
 19. Levey AS, Stevens LA, Schmid CH, Zhang YL, Castro AF 3rd, Feldman HI, et al. New Equation to Estimate Glomerular Filtration Rate. 2009; **150(9)**: 604-12.
 20. Crawford SB, Kosinski AS, Lin HM, Williamson JM, Barnhart HX. Computer programs for the concordance correlation coefficient. *Comput Methods Programs Biomed.* 2007; **88(1)**: 62-74.
 21. Bland JM, Altman DG: Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet* 1: 307-310, 1986.
 22. Stevens LA, Zhang Y, Schmid CH. Evaluating the performance of GFR estimation equations. *J Nephrol.* 2008; **21(6)**: 797-807.
 23. Arreola-Guerra JM, Rincón-Pedrero R, Cruz-Rivera C, Belmont-Pérez T, Correa-Rotter R, Niño-Cruz JA. Performance of MDRD-IDMS and CKD-EPI equations in Mexican individuals with normal renal function. *Nefrologia.* 2014; **34(5)**: 591-8.
 24. Veronese FV, Gomes EC, Chanan J, Carraro MA, Camargo EG, Soares AA, et al. Performance of CKD-EPI equation to estimate glomerular filtration rate as compared to MDRD equation in South Brazilian individuals in each stage of renal function. *Clin Chem Lab Med.* 2014; **52(12)**: 1747-54.
 25. Eastwood JB, Kerry SM, Plange-rhule J, Micah FB, Antwi S, Boa FG, et al. Assessment of GFR by four methods in adults in Ashanti , Ghana : the need for an eGFR equation for lean African populations. *Nephrol Dial Transplant.* 2010 Jul; **25(7)**: 2178-87.
 26. Michels WM, Grootendorst DC, Verduijn M, Elliott EG, Dekker FW, Krediet RT. Performance of the Cockcroft-Gault, MDRD, and new CKD-EPI formulas in relation to GFR, age, and body size. *Clin J Am Soc Nephrol.* 2010 Jun; **5(6)**: 1003-9.
 27. Kamaruzaman L, Mohd R, Zaki FM, Hod R, Aziz AA. Estimating glomerular filtration rate in adult kidney transplant recipients in the Asian population. *Saudi J Kidney Dis Transpl.* 2019 May-Jun; **30(3)**: 587-596.
 28. Uche CL, Osegbe ID. Comparison of CKD-EPI versus MDRD and Cockcroft-Gault equations to estimate glomerular filtration rate among stable homozygous sickle cell patients in Southwest Nigeria. *Clin Chem Lab Med.* 2014 Dec; **52(12)**: 1747-54.
 29. Kuo CF, Yu KH, Shen YM, See LC. The Chinese Version of the Modification of Diet in Renal Disease (MDRD) Equation is a Superior Screening Tool for Chronic Kidney Disease among Middle-aged Taiwanese than the Original MDRD and Cockcroft-Gault Equations. *Biomed J.* 2014 Nov-Dec; **37(6)**: 398-405.
 30. Liu X, Cheng MH, Shi CG, Wang C, Cheng CL, Chen JX, et al. Variability of glomerular filtration rate estimation equations in elderly Chinese patients with chronic kidney disease. *Clin Interv Aging.* 2012; **7**: 409-15.
 31. Schwandt A, Denking M, Fasching P, Pfeifer M, Wagner C, Zeyfang A, et al. Comparison of MDRD, CKD-EPI, and Cockcroft-Gault equation in relation to measured glomerular filtration rate among a large cohort with diabetes. *J Diabetes Complications.* 2017 Sep; **31(9)**: 1376-1383.
 32. Chudek J, Kolonko A, Owczarek AJ, Wieczorowska K. Clinical factors increasing discrepancies of renal function assessment with MDRD and Cockcroft – Gault equations in old individuals. *Eur Geriatr Med.* 2018; **9(5)**: 713-20.
 33. Aras S, Varli M, Uzun B, Atli T, Keven K, Aras S, et al. Comparison of Different Glomerular Filtration Methods in the Elderly : Which Formula Provides Better Estimates ? Comparison of Different Glomerular Filtration Methods in the Elderly : Which Formula Provides Better Estimates ?. *Ren Fail.* 2012; **34(4)**: 435-41.
 34. Zaman SB. Detection of Chronic Kidney Disease by Using Different Equations of Glomerular Filtration Rate in Patients with Type 2 Diabetes Mellitus: A Cross-Sectional Analysis. *Cureus.* 2017 Jun 14; **9(6)**: e1352.

