





# An updated reef fish checklist of the southernmost Caribbean reef system, with comments on the lionfish invasion

Lista actualizada de los peces arrecifales del sistema arrecifal más meridional del Caribe, con notas sobre la invasión del pez león

Camilo Escobar-Sierra  , Viviana Márquez Velásquez  , Rafael Menezes  ,  
Ricardo Souza Rosa  , Alejandro Loaiza-Santana  

---

## Abstract

Chocó-Darién is an important biogeographic realm, as it is a terrestrial biodiversity hotspot and the southern limit of the Caribbean reefs. However, to date there are no compiled data on the reef fish assemblage of this region. We provide an updated checklist of marine fishes from the Chocó-Darién reef system (Colombia), with comments on their geographic distribution and conservation status. Peer-reviewed studies, unpublished data and *in situ* visual censuses were surveyed to compose this checklist. A total of 212 reef fish species across 57 families were compiled, eight of which had no previously published records, one of which (*Trachinotus falcatus*) is recorded for the first time. The most speciose families were Labridae (n = 21), Gobiidae (n = 18) and Serranidae (n = 17). Fourteen threatened species were recorded, including one critically endangered (*Epinephelus striatus*) and two endangered (*Balistes vetula* and *Scarus coelestinus*). This study contributes to fill the knowledge gaps on the reef fish diversity of the Caribbean southern limit and raises concern on the spread of the lionfish invasion into the Chocó-Darién reef system.

**Keywords.** Atlantic. Biodiversity. Chocó-Darién. Neotropic. *Pterois volitans*.

## Resumen

El Chocó-Darién es una importante área biogeográfica, pues es un *hotspot* de biodiversidad terrestre y el límite sur de los arrecifes del Caribe. Sin embargo, hasta la fecha no existen datos compilados actualizados sobre las especies de peces de arrecife de esta región. En este trabajo proporcionamos una lista actualizada de peces marinos del sistema arrecifal del Chocó-Darién (Colombia), con comentarios sobre su distribución geográfica y estado de conservación. Para la elaboración de esta lista se consultaron publicaciones, datos no publicados y censos visuales *in situ*. Se recopilieron 212 especies de peces de arrecife de 57 familias. Ocho de estas especies no tenían registros previos publicados, y una de ellas es registrada por primera vez (*Trachinotus falcatus*). Las familias con mayor número de especies fueron Labridae (n = 21), Gobiidae (n = 18) y Serranidae (n = 17). Se registraron 14 especies amenazadas, entre ellas una en peligro crítico (*Epinephelus striatus*) y dos en peligro (*Balistes vetula* y *Scarus coelestinus*). Este estudio contribuye a complementar las lagunas de conocimiento sobre la diversidad de los peces de arrecife del límite sur del Caribe y plantea la preocupación de la invasión del pez león en el sistema arrecifal del Chocó-Darién.

**Palabras clave.** Atlántico. Biodiversidad. Chocó-Darién. Neotrópico. *Pterois volitans*.

## Introduction

The continental coast of the southernmost portion of the Caribbean Sea, called Urabá Gulf, harbours particular ecological and geological features that differ markedly from other widely explored areas of the Caribbean (O'Dea, 2012). About 3.5 Myr ago, the area functioned as a deep ocean corridor connecting the fauna of the eastern Pacific and Caribbean Sea. After the closure of the Isthmus of Panama, two ecologically divergent areas were formed, with the Pacific side characterized by ocean-based environments and the Caribbean side dominated by coral reef ecosystems (Glynn, 1982).

This southernmost portion of the Caribbean Sea belongs to the Chocó-Darién biogeographic realm. It is regarded as a hotspot of global biodiversity, with high biological relevance for the Colombian Caribbean (Myers *et al.*, 2000). The region encompasses large remnants of humid forest and a mosaic of coastal habitats, including riverine, estuarine and reef ecosystems (Díaz *et al.*, 2000). Although the coastline is strongly influenced by large discharge of terrigenous sediment and freshwater from the Atrato River (Chevillot *et al.*, 1993), fringing patch reefs flourish in such harsh conditions with a typical reef fish fauna and the largest living coral cover across the region (Díaz *et al.*, 2000).

Despite its importance, the Chocó-Darién reef system remains poorly studied (but see Acero & Garzón, 1987a; Reyes-Nivia *et al.*, 2004), with a limited knowledge on its ichthyofauna when compared to other areas of the Colombian Caribbean, such as the San Andrés and Providencia archipelago (Victoria & Gómez, 1984; Mejía & Garzón-Ferreira, 2000), Santa Marta (Acero & Garzón, 1987b; Acero & Rivera, 1992; Grijalba-Bendeck *et al.*, 2004) and Islas del Rosario archipelago (Acero & Garzón, 1985; Delgadillo-Garzón & Zapata-Ramírez, 2009). Like other areas of the Colombian Caribbean, the Chocó-Darién reef has shown clear signs of degradation, evidenced by the dominance of macroalgae, pollution, unregulated tourism and, more recently, the occurrence of one of the invasive species of lionfish *Pterois volitans* (Betancur-R. *et al.*, 2011; Gómez-López *et al.*, 2018).

Lionfishes (*P. miles* and *P. volitans*) were the first non-native marine species to be established in the north-western North Atlantic to the Caribbean and Campeche Bank (Schofield, 2009). These species are considered voracious predators coupled with a high reproductive output (Côté *et al.*, 2013). Hence, declines of native fish populations in reef systems have been correlated with the increment

of their abundances (Green *et al.*, 2012). Lionfishes have sharply dispersed across the Caribbean coral reefs after the first record at the Providencia Island, Colombia in 2008 (Betancur-R. *et al.*, 2011). Over the last years, *P. volitans* has increasingly been reported at the Chocó-Darién reefs (Galvis & Galvis, 2016; García & Rueda 2018; Rojas-Vélez *et al.*, 2019), which raises concern about the potential ecological outcomes in the near future.

This study provides an updated checklist of reef fish of the Chocó-Darién region, based on compiled data of visual censuses and literature, with notes on conservation status and species distribution. In addition, the current status on lionfish (*P. volitans*) invasion is discussed here. This compilation lays foundation for supporting ongoing and future studies for conservation of reef fishes and management of marine resources and services of the region.


## Material and methods

**Study area.** The study region is located in the western tip of the coast of the Urabá Gulf, Colombia, the southernmost portion of the Caribbean Sea. This area includes three sampling localities: Capurganá Bay (8°38'13.45"N; 77°20'39.29"W), El Aguacate (8°37'5.86"N; 77°19'28.53"W) and Sapzurro (8°39'37.02"N; 77°21'40.37"W) (Figure 1).


The waters of the Urabá Gulf are affected by an overload of freshwater and sediments from the Atrato River, which plays a critical role for local biogeography (Restrepo & Kjerfve, 2004). The Chocó-Darién reef system is characterized by a mosaic of fringing reefs, mainly composed by large colonies of *Siderastrea siderea* (Figure 2a) and a less extensive coral cover of *Porites porites*, *Pseudodiploria strigosa*, *Agaricia agaricites*, *Agaricia tenuifolia* and *Millepora complanata* (Gómez-López *et al.*, 2018).

**Checklist compilation.** The checklist was compiled from three sources: visual censuses in field expeditions, literature survey of peer-reviewed articles, and unpublished data (dissertations/theses).

Six visual censuses were performed, using roving diving technique (snorkeling) to quantify reef fish species richness in the Capurganá coastal reefs on April 2011 (Hill & Wilkinson, 2004). Three fringing coastal reefs were chosen, based on a previous survey and information on reef spatial arrangement (Díaz *et al.*, 2000).



**Figure 1.** Western Caribbean coast of Colombia where the samplings of reef fish were carried out.



**Figure 2.** A small representation of the reef fish diversity recorded for the Chocó-Darién reef system, Colombia. A, *Sparisoma viride*, *Thalassoma bifasciatum* and *Cephalopholis cruentata* over a colony of *Siderastrea siderea* coral at El Aguacate; B, *P. volitans* at the Capurganá bay, ca. 7 m depth; C, *Aetobatus narinari* at the Capurganá bay.

In the sampling sites, two divers freely swam for 30 minutes, identifying the fishes and, when possible, recording images and videos. The distance from the observer to the fish was a maximum of three meters, depending on the transparency, to avoid possible taxonomic errors. New records were only considered when two divers recorded the same species, or when a high-resolution image was available. Identifications of species in photos and videos were confirmed by comparisons with those provided by Reef Fish Identification Guide (Humann & Deloach, 2003).

The list of bony fishes was organized based on Eschmeyer's Catalog of Fishes (Van der Laan *et al.*, 2021), except for Labridae, in which Scarinae was included (Westneat & Alfaro, 2005), whereas Weigmann (2016) was followed for elasmobranchs. Genera and species are listed in alphabetical order. For each species

included in the list, we confirm the geographic distribution based on Fishbase (Froese & Pauly, 2019). We also included the conservation status according to the Colombian National Red Book of marine fishes (Chasqui *et al.*, 2017).

## Results

**Species list.** A total of 212 reef fish species were compiled (Table 1), with 68 species recorded by the visual census. Eight out of the 212 species (3.7 %) had no previous published records: *Coryphopterus dicrus*, *Dactylopterus volitans*, *Diodon holocanthus*, *Epinephelus adscensionis*, *Epinephelus guttatus*, *Ginglymostoma cirratum*, *Hypanus americanus*, and *Trachinotus falcatus*. The latter species is a new record for the area and was sighted in our visual censuses. Additionally, we report a species of the *Mugil curema* complex (Nirchio *et al.*, 2017) (Table 1).

**Table 1.** Species of reef fishes recorded for the Chocó-Darién reef ecosystem, Colombia. References: 1, Acero & Garzón (1987a); 2, Reyes-Nivia *et al.* (2004); 3, Peláez de la Torre (2010); 4, Guzmán & Leal (2011); 5, Ramírez & Gaviria (2013); 6, this study. Geographic range: CG, circumglobal; CT, Circumtropical; PAC, Pacific and Transatlantic (Appearing in both Western Atlantic and Eastern Atlantic); TA+MED, Transatlantic and Mediterranean; WA, Western Atlantic [Bahamas, Florida (USA), and northern Gulf of Mexico to Brazil]; WCA, Western Central Atlantic (Greater Caribbean); WI, Western Atlantic and Oceanic islands (Islands of St. Helena and Ascension). Conservation categories (IUCN 2019): CR, Critically Endangered; DD, Data deficient; EN, Endangered; LC, Least concern; NE, Not evaluated; NT, Near Threatened; VU, Vulnerable.

Species	Reference	Geographic range	Conservation Status
<b>ORDER ORECTOLOBIFORMES</b>			
<b>Family Ginglymostomatidae</b>			
<i>Ginglymostoma cirratum</i> (Bonnaterre 1788)	3, 4	TA	VU
<b>ORDER MYLIOBATIFORMES</b>			
<b>Family Dasyatidae</b>			
<i>Hypanus americanus</i> (Hildebrand & Schroeder 1928)	3, 6	WA	NT
<i>Styracura schmardae</i> (Werner 1904)	1	WA	NE
<b>Family Myliobatidae</b>			
<i>Aetobatus narinari</i> (Euphrasen 1790)	1, 6	TA	NE
<b>ORDER ANGUILLIFORMES</b>			
<b>Family Muraenidae</b>			
<i>Echidna catenata</i> (Bloch 1795)	1, 2, 3, 6	WI	NE
<i>Enchelycore carychroa</i> Böhlke & Böhlke 1976	1	WI	NE
<i>Enchelycore nigricans</i> (Bonnaterre 1788)	1	TA	NE

Species	Reference	Geographic range	Conservation Status
<i>Gymnothorax funebris</i> Ranzani 1839	1, 4, 5	WA	NE
<i>Gymnothorax miliaris</i> (Kaup 1856)	1, 2, 3	TA	NE
<i>Gymnothorax moringa</i> (Cuvier 1829)	1, 4, 5	WI	NE
<i>Gymnothorax vicinus</i> (Castelnau 1855)	1	TA	NE
<b>Family Chlopsidae</b>			
<i>Kaupichthys hyoproroides</i> (Strömman 1896)	1	CT	NE
<b>Family Ophichthidae</b>			
<i>Ahlia egmontis</i> (Jordan 1884)	1	WA	NE
<i>Myrichthys ocellatus</i> (Lesueur 1825)	1	TA	NE
<b>Family Moringuidae</b>			
<i>Moringua edwardsi</i> (Jordan & Bollman 1889)	1	WCA	NE
<b>ORDER CLUPEIFORMES</b>			
<b>Family Engraulidae</b>			
<i>Anchoa lyolepis</i> (Evermann & Marsh 1900)	1	WA	NE
<i>Anchoviella perfasciata</i> (Poey 1860)	1	WCA	NE
<b>Family Clupeidae</b>			
<i>Harengula clupeola</i> (Cuvier 1829)	1	WA	NE
<i>Jenkinsia lamprotaenia</i> (Gosse 1851)	1	WCA	NE
<i>Opisthonema oglinum</i> (Lesueur 1818)	1	WA	NE
<b>ORDER AULOPIIFORMES</b>			
<b>Family Synodontidae</b>			
<i>Synodus intermedius</i> (Spix & Agassiz 1829)	1, 2, 3	WA	NE
<i>Synodus synodus</i> (Linnaeus 1758)	1	TA	NE
<b>ORDER HOLOCENTRIFORMES</b>			
<b>Family Holocentridae</b>			
<i>Holocentrus adscensionis</i> (Osbeck 1765)	1, 2, 3, 4, 5	TA	NE
<i>Holocentrus rufus</i> (Walbaum 1792)	2, 3, 4, 5	WA	NE
<i>Myripristis jacobus</i> Cuvier 1829	1, 2, 3, 4, 6	TA	NE
<i>Neoniphon marianus</i> (Cuvier 1829)	2	WA	NE
<i>Neoniphon vexillarium</i> (Poey 1860)	1, 2, 3	WA	NE
<i>Plectrypops retrospinis</i> (Guichenot 1853)	1	WCA	NE
<b>ORDER OPHIDIIFORMES</b>			
<b>Family Ophidiidae</b>			
<i>Parophidion schmidti</i> (Woods & Kanazawa 1951)	1	WA	NE

Species	Reference	Geographic range	Conservation Status
<b>Family Bythitidae</b>			
<i>Ogilbia cayorum</i> Evermann & Kendall 1898	1	WCA	NE
<b>ORDER SCOMBRIFORMES</b>			
<b>Family Scombridae</b>			
<i>Scomberomorus regalis</i> (Bloch 1793)	2	TA	NE
<b>ORDER SYNGNATHIFORMES</b>			
<b>Family Aulostomidae</b>			
<i>Aulostomus maculatus</i> Valenciennes 1841	1, 2, 3, 4, 6	WA	NE
<b>Family Fistulariidae</b>			
<i>Fistularia tabacaria</i> Linnaeus 1758	4, 5	TA	NE
<b>Family Syngnathidae</b>			
<i>Cosmocampus brachycephalus</i> (Poey 1868)	1	WA	NE
<i>Halicampus crinitus</i> (Jenyns 1842)	1	WCA	NE
<i>Hippocampus reidi</i> Ginsburg 1933	1	WA	VU
<i>Microphis brachyurus</i> (Bleeker 1854)	1	CT	NE
<b>Family Dactylopteridae</b>			
<i>Dactylopterus volitans</i> (Linnaeus 1758)	5	TA+MED	NE
<b>ORDER GOBIIFORMES</b>			
<b>Family Gobiidae</b>			
<i>Barbulifer ceuthoecus</i> (Jordan & Gilbert 1884)	1	WA	NE
<i>Bathygobius soporator</i> (Valenciennes 1837)	1	TA	NE
<i>Coryphopterus dicrus</i> Böhlke & Robins 1960	3, 6	WA	NE
<i>Coryphopterus eidolon</i> Böhlke & Robins 1960	2	WCA	NE
<i>Coryphopterus glaucofraenum</i> Gill 1863	1, 2, 3	WA	NE
<i>Coryphopterus lipernes</i> Böhlke & Robins 1962	2	WCA	NE
<i>Coryphopterus personatus</i> (Jordan & Thompson 1905)	2, 3, 6	WCA	NE
<i>Coryphopterus thrix</i> Böhlke & Robins 1960	2, 3, 6	WCA	NE
<i>Elacatinus illecebrosus</i> (Böhlke & Robins 1968)	1, 2, 3, 6	WCA	NE
<i>Elacatinus</i> sp.	4, 5		
<i>Ginsburgellus novemlineatus</i> (Fowler 1950)	1	WCA	NE
<i>Gnatholepis thompsoni</i> Jordan 1904	1, 2, 3, 6	TA	NE
<i>Lythrypnus nesiotus</i> Böhlke & Robins 1960	1	WCA	NE
<i>Lythrypnus</i> sp.	1		
<i>Lythrypnus spilus</i> Böhlke & Robins 1960	1	WCA	NE

Species	Reference	Geographic range	Conservation Status
<i>Priolepis hipoliti</i> (Metzelaar 1922)	1	WA	NE
<i>Tigrigobius multifasciatus</i> (Steindachner 1876)	1	WCA	NE
<i>Tigrigobius saucrus</i> (Robins 1960)	2, 3	WCA	NE
<b>ORDER CARANGIFORMES</b>			
<b>Family Sphyraenidae</b>			
<i>Sphyraena barracuda</i> (Edwards 1771)	1, 3	CG	NT
<b>Family Bothidae</b>			
<i>Bothus lunatus</i> (Linnaeus 1758)	1, 2, 3	TA	NE
<b>Family Carangidae</b>			
<i>Caranx bartholomaei</i> Cuvier 1833	1, 2, 3	WI	NE
<i>Caranx hippos</i> (Linnaeus 1766)	1, 2, 4	TA+MED	VU
<i>Caranx ruber</i> (Bloch 1793)	1, 3, 4, 5, 6	WA	NE
<i>Decapterus</i> sp.	2		
<i>Trachinotus falcatus</i> (Linnaeus 1758)	6	WA	NE
<b>ORDER BELONIFORMES</b>			
<b>Family Belonidae</b>			
<i>Strongylura</i> sp.	1, 6		
<b>ORDER MUGILIFORMES</b>			
<b>Family Mugilidae</b>			
<i>Mugil gr. curema</i>	6	CG	NE
<b>ORDER GOBIESOCIFORMES</b>			
<b>Family Gobiesocidae</b>			
<i>Acyrtops beryllinus</i> (Hildebrand & Ginsburg 1927)	1	WA	NE
<i>Acyrtus rubiginosus</i> (Poey 1868)	1	WCA	NE
<i>Gobiesox punctulatus</i> (Poey 1876)	1	WA	NE
<b>ORDER BLENNIIFORMES</b>			
<b>Family Tripterygiidae</b>			
<i>Enneanectes altivelis</i> Rosenblatt 1960	1	WA	NE
<i>Enneanectes boehlkei</i> Rosenblatt 1960	1	WCA	NE
<b>Family Labrisomidae</b>			
<i>Gobioclinus kalisherae</i> (Jordan 1904)	1	WA	NE
<i>Labrisomus nuchipinnis</i> (Quoy & Gaimard 1824)	1	TA	NE
<i>Malacoctenus macropus</i> (Poey 1868)	1	WCA	NE
<i>Malacoctenus triangulatus</i> Springer 1959	1, 2	WI	NE

Species	Reference	Geographic range	Conservation Status
<i>Paraclinus nigripinnis</i> (Steindachner 1867)	1	WA	NE
<i>Starksia variabilis</i> Greenfield 1979	1	WCA	NE
<i>Stathmonotus gymnodermis</i> Springer 1955	1	WA	NE
<b>Family Chaenopsidae</b>			
<i>Acanthemblemaria rivasi</i> Stephens 1970	2, 3	WCA	NE
<i>Coralliozetus</i> sp.	1		
<i>Ekemblemaria nigra</i> (Meek & Hildebrand 1928)	1	WCA	NE
<i>Lucayablennius zingaro</i> (Böhlke 1957)	2, 3	WCA	NE
<b>Family Dactyloscopidae</b>			
<i>Dactyloscopus tridigitatus</i> Gill 1859	1	WA	NE
<i>Platygillemus rubrocinctus</i> (Longley 1934)	1	WCA	NE
<b>Family Blenniidae</b>			
<i>Entomacrodus nigricans</i> Gill 1859	1	WCA	NE
<i>Hypsoblennius invemar</i> Smith-Vaniz & Acero P. 1980	1	WA	NE
<i>Ophioblennius macclurei</i> (Silvester 1915)	1, 3, 4, 6	WA	NE
<b>ORDER ACANTHURIFORMES</b>			
<b>Family Pomacanthidae</b>			
<i>Holacanthus ciliaris</i> (Linnaeus 1758)	1, 2, 4	WI	NE
<i>Holacanthus tricolor</i> (Bloch 1795)	2, 4	WA	NE
<i>Pomacanthus arcuatus</i> (Linnaeus 1758)	1, 2, 3, 4, 6	WA	NE
<i>Pomacanthus paru</i> (Bloch 1787)	1, 2, 3, 4, 6	WI	NE
<b>Family Chaetodontidae</b>			
<i>Chaetodon capistratus</i> Linnaeus 1758	1, 2, 3, 4, 5, 6	WA	NE
<i>Chaetodon ocellatus</i> Bloch 1787	1, 2, 3, 4, 5, 6	WA	NE
<i>Chaetodon sedentarius</i> Poey 1860	2, 6	WA	NE
<i>Chaetodon striatus</i> Linnaeus 1758	2, 3, 4, 5, 6	WI	NE
<b>Family Acanthuridae</b>			
<i>Acanthurus chirurgus</i> (Bloch 1787)	1, 2, 3, 4, 5, 6	TA	NE
<i>Acanthurus coeruleus</i> Bloch & Schneider 1801	1, 2, 3, 4, 5, 6	TA	NE
<i>Acanthurus tractus</i> Poey 1860	1, 2, 3, 4, 5, 6	WCA	NE
<b>ORDER LOPHIIFORMES</b>			
<b>Family Antennariidae</b>			



Species	Reference	Geographic range	Conservation Status
<i>Antennarius multicephalus</i> (Valenciennes 1837)	1	WI	NE
<b>ORDER TETRADONTIFORMES</b>			
<b>Family Diodontidae</b>			
<i>Diodon holocanthus</i> Linnaeus 1758	4	CT	NE
<i>Diodon hystrix</i> Linnaeus 1758	1, 2, 3, 6	CT	NE
<b>Family Tetraodontidae</b>			
<i>Canthigaster rostrata</i> (Bloch 1786)	1, 2, 3, 4, 6	WCA	NE
<i>Sphoeroides spengleri</i> (Bloch 1785)	1	WA	NE
<i>Sphoeroides testudineus</i> (Linnaeus 1758)	1	WA	NE
<b>Family Ostraciidae</b>			
<i>Acanthostracion polygonius</i> Poey 1876	2, 3	WA	NE
<i>Acanthostracion quadricornis</i> (Linnaeus 1758)	2, 3	TA	NE
<i>Lactophrys bicaudalis</i> (Linnaeus 1758)	2, 3, 6	WI	NE
<i>Lactophrys trigonus</i> (Linnaeus 1758)	1	WA	NE
<i>Lactophrys triqueter</i> (Linnaeus 1758)	2, 3	WA	NE
<b>Family Monacanthidae</b>			
<i>Aluterus scriptus</i> (Osbeck 1765)	2, 3	CT	NE
<i>Cantherhines macrocerus</i> (Hollard 1853)	2, 3, 4	WI	NE
<i>Cantherhines pullus</i> (Ranzani 1842)	1, 2, 3, 4, 6	TA	NE
<i>Monacanthus tuckeri</i> Bean 1906	2	WCA	NE
<b>Family Balistidae</b>			
<i>Balistes vetula</i> Linnaeus 1758	2	TA	EM
<i>Canthidermis sufflamen</i> (Mitchill 1815)	1, 2, 3, 6	TA	NE
<i>Melichthys niger</i> (Bloch 1786)	2	CT	NE
<b>ORDER CENTRARCHIFORMES</b>			
<b>Family Kyphosidae</b>			
<i>Kyphosus</i> sp.	2, 3, 6		
<i>Kyphosus sectatrix</i> (Linnaeus 1758)	1, 4, 5	TA+MED	NE
<b>Family Cirrhitidae</b>			
<i>Amblycirrhitus pinos</i> (Mowbray 1927)	2, 3	WA	NE
<b>ORDER ACROPOMATIFORMES</b>			
<b>Family Pempheridae</b>			

Species	Reference	Geographic range	Conservation Status
<i>Pempheris schomburgkii</i> Müller & Troschel 1848	1, 2, 3, 4, 5, 6	WA	NE
<b>ORDER PERCIFORMES INCERTAE SEDIS</b>			
<b>Family Serranidae</b>			
<i>Cephalopholis cruentata</i> (Lacepède 1802)	1, 3, 4, 5, 6	WCA	NE
<i>Cephalopholis fulva</i> (Linnaeus 1758)	1, 3	WA	NE
<i>Epinephelus adscensionis</i> (Osbeck 1765)	4, 5	TA	NE
<i>Epinephelus guttatus</i> (Linnaeus 1758)	4, 5	WA	NT
<i>Epinephelus striatus</i> (Bloch 1792)	1, 2, 3, 6	WA	CR
<i>Hypoplectrus nigricans</i> (Poey 1852)	2, 3	WCA	NE
<i>Hypoplectrus puella</i> (Cuvier 1828)	1, 2, 3, 4, 5, 6	WCA	NE
<i>Hypoplectrus unicolor</i> (Walbaum 1792)	2, 3	WCA	NE
<i>Liopropoma rubre</i> Poey 1861	2	WA	NE
<i>Mycteroperca bonaci</i> (Poey 1860)	1, 2	WA	VU
<i>Mycteroperca tigris</i> (Valenciennes 1833)	2	WA	NT
<i>Mycteroperca venenosa</i> (Linnaeus 1758)	2	WA	VU
<i>Pseudogramma gregoryi</i> (Breder 1927)	1	WA	NE
<i>Rypticus saponaceus</i> (Bloch & Schneider 1801)	1, 2, 4, 5	TA	NE
<i>Rypticus subbifrenatus</i> Gill 1861	1	TA	NE
<i>Serranus baldwini</i> (Evermann & Marsh 1899)	1	WA	NE
<i>Serranus tigrinus</i> (Bloch 1790)	1, 2, 3, 4, 5, 6	WCA	NE
<b>Family Grammatidae</b>			
<i>Gramma loreto</i> Poey 1868	1, 2, 3, 4, 5, 6	WA	NE
<i>Gramma melacara</i> Böhlke & Randall 1963	2	WCA	NE
<b>Family Opistognathidae</b>			
<i>Opistognathus whitehursti</i> (Longley 1927)	1	WA	NE
<b>Family Priacanthidae</b>			
<i>Heteropriacanthus cruentatus</i> (Lacepède 1801)	2, 3	CT	NE
<b>Family Apogonidae</b>			
<i>Apogon maculatus</i> (Poey 1860)	1	WA	NE
<i>Astrapogon puncticulatus</i> (Poey 1867)	1	WA	NE
<i>Phaeoptyx conklini</i> (Silvester 1915)	1	WA	NE
<i>Phaeoptyx pigmentaria</i> (Poey 1860)	1	TA	NE
<b>Family Malacanthidae</b>			
<i>Malacanthus plumieri</i> (Bloch 1786)	1	WI	NE

Species	Reference	Geographic range	Conservation Status
<b>Family Lutjanidae</b>			
<i>Lutjanus analis</i> (Cuvier 1828)	1, 2, 3, 6	WA	VU
<i>Lutjanus apodus</i> (Walbaum 1792)	2, 3, 4	WA	NE
<i>Lutjanus cyanopterus</i> (Cuvier 1828)	1, 2	WA	VU
<i>Lutjanus griseus</i> (Linnaeus 1758)	1	WA	NE
<i>Lutjanus jocu</i> (Bloch & Schneider 1801)	1, 2, 3	TA	DD
<i>Lutjanus mahogoni</i> (Cuvier 1828)	1, 2, 3, 4	WCA	NE
<i>Lutjanus synagris</i> (Linnaeus 1758)	2, 3	WA	NE
<i>Ocyurus chrysurus</i> (Bloch 1791)	1, 2, 3, 4, 5, 6	TA	NT
<b>Family Gerreidae</b>			
<i>Eucinostomus</i> sp.	1		
<i>Gerres cinereus</i> (Walbaum 1792)	1	WA	NE
<b>Family Haemulidae</b>			
<i>Anisotremus surinamensis</i> (Bloch 1791)	2, 3	WA	NE
<i>Anisotremus virginicus</i> (Linnaeus 1758)	1, 2, 3, 4, 6	WA	NE
<i>Brachygenys chrysargyreum</i> (Günther 1859)	2, 3	WA	NE
<i>Haemulon album</i> Cuvier 1830	2, 3	WA	LC
<i>Haemulon aurolineatum</i> Cuvier 1830	1, 2, 3, 4, 6	WA	NE
<i>Haemulon bonariense</i> Cuvier 1830	2, 3	WCA	NE
<i>Haemulon carbonarium</i> Poey 1860	1, 2, 3, 4, 6	WCA	NE
<i>Haemulon flavolineatum</i> (Desmarest 1823)	1, 2, 3, 4, 6	WA	NE
<i>Haemulon macrostoma</i> Günther 1859	1, 2, 3, 4, 5, 6	WA	NE
<i>Haemulon parra</i> (Desmarest 1823)	2, 3	WA	NE
<i>Haemulon plumierii</i> (Lacepède 1801)	1, 2, 3, 6	WA	NE
<i>Haemulon sciurus</i> (Shaw 1803)	2, 3, 4, 5	WA	NE
<b>Family Sciaenidae</b>			
<i>Equetus lanceolatus</i> (Linnaeus 1758)	2, 3	WA	NE
<i>Equetus punctatus</i> (Bloch & Schneider 1801)	2, 3, 4, 6	WA	NE
<i>Odontoscion dentex</i> (Cuvier 1830)	2, 3	WA	NE
<b>Family Mullidae</b>			
<i>Mulloidichthys martinicus</i> (Cuvier 1829)	1, 2, 3, 4, 5, 6	TA	NE
<i>Pseudupeneus maculatus</i> (Bloch 1793)	1, 2, 3, 4, 5, 6	WA	NE

Species	Reference	Geographic range	Conservation Status
<b>ORDER PERCIFORMES</b>			
<b>Family Pomacentridae</b>			
<i>Abudefduf saxatilis</i> (Linnaeus 1758)	1, 2, 3, 4, 5, 6	TA+MED	NE
<i>Abudefduf taurus</i> (Müller & Troschel 1848)	1, 6	TA	NE
<i>Chromis cyanea</i> (Poey 1860)	2	WCA	NE
<i>Chromis insolata</i> (Cuvier 1830)	2	WCA	NE
<i>Chromis multilineata</i> (Guichenot 1853)	1, 2, 3, 4, 6	TA	NE
<i>Microspathodon chrysurus</i> (Cuvier 1830)	1, 2, 3, 4, 6	WA	NE
<i>Stegastes adustus</i> (Troschel 1865)	1, 2, 3, 4, 5, 6	WCA	NE
<i>Stegastes diencaeus</i> (Jordan & Rutter 1897)	2, 3	WCA	NE
<i>Stegastes leucostictus</i> (Müller & Troschel 1848)	1, 2, 3, 6	WA	NE
<i>Stegastes partitus</i> (Poey 1868)	1, 2, 3, 4, 6	WCA	NE
<i>Stegastes planifrons</i> (Cuvier 1830)	1, 2, 3, 4, 6	WCA	NE
<i>Stegastes xanthurus</i> (Poey 1860)	1, 2, 3	WCA	NE
<b>Family Labridae</b>			
<i>Bodianus rufus</i> (Linnaeus 1758)	1, 2, 3, 4, 6	WA	NE
<i>Clepticus parrae</i> (Bloch & Schneider 1801)	2, 3, 4	WCA	NE
<i>Doratonotus megalepis</i> Günther 1862	1	TA	NE
<i>Halichoeres bivittatus</i> (Bloch 1791)	1, 2, 3, 4, 5, 6	WA	NE
<i>Halichoeres cyanocephalus</i> (Bloch 1791)	2	WA	NE
<i>Halichoeres garnoti</i> (Valenciennes 1839)	1, 2, 3, 4	WA	NE
<i>Halichoeres maculipinna</i> (Müller & Troschel 1848)	1, 2, 3, 4, 5, 6	WA	NE
<i>Halichoeres pictus</i> (Poey 1860)	2, 3	WCA	NE
<i>Halichoeres poeyi</i> (Steindachner 1867)	1, 2, 3, 6	WA	NE
<i>Halichoeres radiatus</i> (Linnaeus 1758)	1, 2, 3, 4, 5, 6	WI	NE
<i>Scarus coelestinus</i> Valenciennes 1840	2	WA	EM
<i>Scarus iseri</i> (Bloch 1789)	1, 2, 3, 6	WCA	NE
<i>Scarus taeniopterus</i> Lesson 1829	2, 3	WA	NE
<i>Scarus vetula</i> Bloch & Schneider 1801	2, 3	WCA	NT
<i>Sparisoma atomarium</i> (Poey 1861)	2, 6	WA	NE
<i>Sparisoma aurofrenatum</i> (Valenciennes 1840)	1, 2, 3, 4, 5, 6	WCA	NE
<i>Sparisoma chrysopterus</i> (Bloch & Schneider 1801)	2, 3, 6	WCA	NE


Species	Reference	Geographic range	Conservation Status
<i>Sparisoma radians</i> (Valenciennes 1840)	1	WA	NE
<i>Sparisoma rubripinne</i> (Valenciennes 1840)	1, 2, 3, 4, 5, 6	WA	NE
<i>Sparisoma viride</i> (Bonnaterre 1788)	1, 2, 3, 4, 5, 6	WA	NT
<i>Thalassoma bifasciatum</i> (Bloch 1791)	1, 2, 3, 4, 5, 6	WA	NE
<b>Family Scorpaenidae</b>			
<i>Pterois volitans</i> (Linnaeus 1758)	4, 6	PAC	NE
<i>Scorpaena bergii</i> Evermann & Marsh 1900	1	WA	NE
<i>Scorpaena grandicornis</i> Cuvier 1829	1	WA	NE
<i>Scorpaena isthmensis</i> Meek & Hildebrand 1928	1	WA	NE
<i>Scorpaena plumieri</i> Bloch 1789	1, 4	WI	NE
<i>Scorpaenodes caribbaeus</i> Meek & Hildebrand 1928	1	WA	NE
<i>Scorpaenodes tredecimspinosus</i> (Metzelaar 1919)	1	WA	NE

Chocó-Darién reef fishes are distributed in 21 orders and 57 families, the most representative orders being Perciformes plus Perciformes *incertae sedis* (43.86 %), followed by Gobiiformes (8.49%), Blenniiformes (8.49 %), and Tetraodontiformes (8.02 %). The most speciose families were Labridae (9.91 %), followed by Gobiidae (8.49 %), Serranidae (8.02 %), Haemulidae and Pomacentridae (5.66 %). Likewise, the most speciose genera were *Haemulon* ( $n = 9$ ), followed by *Halichoeres* and *Lutjanus* ( $n = 7$ ), *Sparisoma*, *Stegastes* and *Coryphopterus* ( $n = 6$ ).

From the recorded fishes, 96 are widespread species of the Western Atlantic, 50 of the Greater Caribbean, 33 Trans-Atlantic, seven Circumtropical, four Transatlantic and Mediterranean, two Circumglobal and one restricted to the Pacific Ocean (Figure 3). The latter species is the introduced lionfish *P. volitans* (Figure 2b).

Of the 212 species, one is listed as Data Deficient (DD), one species as Least Concern (LC), seven as Near-threatened (NT), seven as Vulnerable (VU), two as Endangered (EN) and one as Critically Endangered (CR). A total of 10 species are in a threat category for the Chocó-Darién coral reef, based in the National Red Book (Chasqui et al., 2017). The Nassau grouper (*Epinephelus striatus*) is

Critically Endangered. In addition, the Queen Triggerfish *Balistes vetula* and the Midnight Parrotfish *Scarus coelestinus* are categorized as Endangered. However, most species (186) have not yet been evaluated in Colombia.



**Figure 3.** Relative frequency of the geographic range categories of reef fish recorded for the Chocó-Darién reef system, Colombia. CG, circumglobal; CT, circum-tropical; PAC, Pacific; TA, Transatlantic (Western Atlantic and Eastern Atlantic); TA+MED, Transatlantic and Mediterranean; WA, Western Atlantic; WCA, Western Central Atlantic (Caribbean); WI, Western Atlantic and Oceanic islands.

## Discussion

This work represents an extensive compilation on the ichthyofauna diversity of the Chocó-Darién reef system, Colombian Caribbean. Previous studies have documented contrasting number of fish species, such as 146 (Acero & Garzón, 1987a) and 119 (Reyes-Nivia *et al.*, 2004), which is possibly due to the use of different sampling methodologies.

The most speciose families recorded in this work (e.g., Labridae, Gobiidae, Serranidae, Haemulidae and Pomacentridae) are commonly found in the continental margins of the tropical Atlantic (Floeter, 2008). Nevertheless, the fish richness recorded for the study area (212 species) represents only 30 % of the total species accounted across the Caribbean region, indicating its relatively low richness. For example, Acero & Garzón (1987b) recorded 372 species at the Santa Marta reef systems, Colombian Caribbean, and Starck (1968) recorded 389 species at the Alligator reef, Florida Keys.

Recently, a biogeographic analysis for fish data of both reef and soft bottom divided the Caribbean region into three major provinces: (1) a central, tropical province comprising the West Indies, Bermuda and Central America; (2) a southern, upwelling-affected province spanning the entire continental shelf of northern South America; and (3) a northern, subtropical province that includes all of the Gulf of Mexico, Florida and south-eastern USA (Robertson & Cramer, 2014). The Chocó-Darién reef system is located at the southern province covering the entire continental shelf of northern South America, holding the lowest number of fish species and percentage of local endemics (3.4 %). Likely, the particular environmental conditions of this region, such as the high loads of nutrients, low pH, temperature and salinity variations caused by the Atrato River (McLaughlin *et al.*, 2003, Manzello, 2010) explain, in part, the low species richness of the region.

Most of the listed species present distributions across the Western Atlantic. Biogeographically, the Western Atlantic comprises the Greater Caribbean and Brazil, with their faunas considered until recently to be partially separated by the freshwater discharge from the Amazon and Orinoco river mouths (Floeter *et al.*, 2008). However, an extensive and diverse reef system was recently mapped for the Amazon region (called Great Amazon Reef), which represents the northern limit of

the Brazilian Province and may function as an ecological corridor connecting the fauna of the Brazil and Caribbean (Francini-Filho *et al.*, 2018). Importantly, the Caribbean shelters a higher species richness (774 species) and endemism levels (57 %) compared to Brazil (Kulbicki *et al.*, 2013, Pinheiro *et al.*, 2018), which, in turn, explains the high percentage of Caribbean or Western Central Atlantic endemics species in our study (23 %).

The presence of the species *Gnatholepis thompsoni* and *Pterois volitans* in the Chocó-Darién region is related to different events. *G. thompsoni*, native from the Indian Ocean, possibly dispersed to the north Atlantic during the last interglacial period, and its expansion range has been spreading in face of climate change (Rocha *et al.*, 2005). On the other side, *P. volitans* was introduced in the Florida Keys coastal waters in the 1980's, as a consequence of escapes of the aquarium trade (Morris *et al.*, 2009). Its range has increased over the years, reaching a broad extension of the tropical and subtropical Western Atlantic and Caribbean (Schofield, 2009; Betancur-R *et al.*, 2011). In Colombia, this lionfish was firstly reported in the oceanic islands, and subsequently in the coastal areas of the Caribbean (Betancur-R *et al.*, 2011). Our study raises concerns of the species' invasion at the Chocó-Darién reef system (Galvis & Galvis, 2016; Rojas-Vélez *et al.*, 2019), and supports the previous reports of the expansion of *P. volitans* toward southernmost portion of the Caribbean Sea. This species feeds on a wide variety of juveniles of large-bodied fish (Green & Côté, 2014), and crustaceans, as already reported for Colombian Caribbean regions (Muñoz-Escobar & Gil-Agudelo, 2012; Acero *et al.*, 2019). Such feeding behavior may trigger impacts on local fish populations, and consequently, in the food web dynamics (Valdez-Moreno *et al.*, 2012).

The present study contributes to fill up the knowledge gap on reef fish of the Chocó-Darién reef system. Further research including new technologies such as ROVs (Auster, 1997) and baited remote underwater video systems (BRUVs) (White *et al.*, 2013) are recommended, which would enable exploring remote reef areas, such as mesophotic and rariphotic ecosystems (cf. Francini-Filho *et al.*, 2019). Finally, in face of the fast invasion of the *P. volitans* and the high occurrence of threatened fishes, we recommend strengthening fish monitoring programs to subsidize management and conservation measures at the Chocó-Darién reef system.

## Acknowledgments

The authors thank Lizette Irene Quan Young for assistance and valuable comments on the first drafts of this paper. To the Universidad de Antioquia for supporting the field expedition. Thanks to Eco Hotel Playas de Capurganá and to the Echavarría family, especially Nora Ramírez and Norman Jr. Echavarría, for hosting us and facilitating the samplings. Moreover, to the many generations of marine researchers that have dedicated their time to study the most beautiful corner of the Colombian Caribbean, Capurganá.

## References

- Acero, A. & Garzón, J. (1985). Peces de las islas Rosario y de San Bernardo (Colombia). I. características del área y lista de peces. *Actualidades Biológicas*, 14 (54), 137-148.
- Acero, A. & Garzón, J. (1987a). Los peces marinos hallados durante la expedición Urabá II al Caribe chocoano (Colombia). *Anales del Instituto de Investigaciones Marinas de Punta Betín*, (11), 3-1.  
<https://doi.org/10.25268/bimc.invemar.1987.17.0.459>
- Acero, A. & Garzón, J. (1987b). Peces arrecifales de la región de Santa Marta (Caribe colombiano). I. Lista de especies y comentarios generales. *Acta Biológica Colombiana*, 1 (3), 83-104.
- Acero, A. & Rivera, M. (1992). Peces de las familias Chaetodontidae y Pomacanthidae en la región de Santa Marta (Colombia): densidad y relación con la calidad del arrecife. *Caribbean Journal of Science*, 28 (3-4), 184-190.
- Acero, A., Bustos-Montes, D., Quintero, P. P., Polo-Silva, C. J. & Muñoz, A. S. (2019). Feeding habits of *Pterois volitans*: a real threat to Caribbean coral reef biodiversity. In Makowski, C. & Finkl, C. W. (Eds.). *Impacts of Invasive Species on Coastal Environments*. Pp. 269-314. Springer, Cham.  
[https://doi.org/10.1007/978-3-319-91382-7\\_8](https://doi.org/10.1007/978-3-319-91382-7_8)
- Auster, P. J. (1997). ROV technologies and utilization by the science community. *Marine Technology Society Journal*, 31 (3), 72.
- Betancur-R., R., Hines, A., Acero P., A., Ortí, G., Wilbur, A. E. & Freshwater, D. W. (2011). Reconstructing the lionfish invasion: insights into Greater Caribbean biogeography. *Journal of Biogeography*, 38 (7), 1281-1293.  
<https://doi.org/10.1111/j.1365-2699.2011.02496.x>
- Chasqui, V., Polanco, L. A., Acero, F. A., Mejía-Falla, P. A., Navia, A., Zapata, L. A. & Caldas, J. P. (2017). *Libro rojo de peces marinos de Colombia*. Santa Marta, Colombia: Invemar. 93 pp.
- Chevillot, P., Molina, A., Giraldo, L. & Molina C. (1993). Estudio geológico e hidrológico del Golfo de Urabá. *Boletín Científico CIOH*, 14, 79-89.  
<https://doi.org/10.26640/22159045.62>
- Côté, I. M., Green, S. J. & Hixon, M. A. (2013). Predatory fish invaders: insights from Indo-Pacific lionfish in the western Atlantic and Caribbean. *Biological Conservation*, 164, 50-61.  
<https://doi.org/10.1016/j.biocon.2013.04.014>
- Delgadillo-Garzón, O. & Zapata-Ramírez, P. (2009). Evaluación rápida de peces arrecifales y su relación con la estructura del sustrato en las Islas del Rosario, área marina protegida del Caribe colombiano. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y naturales*, 33(127), 273-283.
- Díaz, J. M., Díaz-Pulido, G. & Sánchez, J. (2000). Distribution and structure of the southernmost Caribbean coral reefs: Golfo de Urabá, Colombia. *Scientia Marina*, 64, 327-336.  
<https://doi.org/10.3989/scimar.2000.64n3327>
- Floeter, S. R., Rocha, L. A., Robertson, D. R., Joyeux, J. C., Smith-Vaniz, W. F., Wirtz, P. & Brito, A. (2008). Atlantic reef fish biogeography and evolution. *Journal of Biogeography*, 35 (1), 22-47.
- Francini-Filho, R. B., Asp, N. E., Siegle, E., Hocevar, J., Lowyck, K., D'Avila, N. & Thompson, C. C. (2018). Perspectives on the Great Amazon Reef: Extension, biodiversity, and threats. *Frontiers in Marine Science*, 5, 142.  
<https://doi.org/10.3389/fmars.2018.00142>
- Francini-Filho, R. B., Márquez Velázquez V., Barbosa da Silva M., Rogerio Rosa M., Gomes Sumida P. Y., Tercio Pinheiro H., Alves Rocha L., Leite Ferreira C. E., Bezerra Francini C. L. & de Souza Rosa, R. (2019). Brazil. In: Loya Y., Puglise K., Bridge T. (Eds) *Mesophotic Coral Ecosystems. Coral Reefs of the World, vol 12*. Springer, Cham, pp. 163-198.  
[https://doi.org/10.1007/978-3-319-92735-0\\_10](https://doi.org/10.1007/978-3-319-92735-0_10)
- Froese, R. & Pauly D. Editors. 2019. FishBase. World Wide Web electronic publication. Available from: <https://www.fishbase.org>, (accessed 01 February 2019)
- Galvis, N. H. & Galvis, R. H. (2016). Colombian citizen science to improve coral reef conservation. In: Birke-land C., Coles S. I. & Spies N. P. (Eds.) *Proceedings of 13th International Coral Reef Symposium*. Pp: 606-619. Honolulu, Hawaii.
- García, C. B. & Rueda, M. (2018). The lionfish *Pterois volitans* (Scorpaeniformes: Scorpaenidae) invades soft-bottoms: evidence from survey data. *Pan-American Journal of Aquatic Sciences*, 13(3), 211-215.

- Glynn, P. W. (1982). Coral communities and their modifications relative to past and prospective Central American seaways. *Advances in Marine Biology*, 19, 91-132.  
[https://doi.org/10.1016/S0065-2881\(08\)60087-5](https://doi.org/10.1016/S0065-2881(08)60087-5)
- Gómez-López, D. I., Acosta-Chaparro A., Gonzalez J. D., Sánchez L., Navas-Camacho, R. & Alonso, D. (2018). *Reporte del estado de los arrecifes coralinos y pastos marinos en Colombia (2016-2017)*. Santa Marta, Colombia: Invemar. 100 pp.  
<https://doi.org/10.3391/ai.2009.4.3.12>
- Green, S. J. & Côté, I. M. (2014). Trait-based diet selection: prey behaviour and morphology predict vulnerability to predation in reef fish communities. *Journal of Animal Ecology*, 83 (6), 1451-1460.  
<https://doi.org/10.1111/1365-2656.12250>
- Green, S. J., Akins, J. L., Maljković, A. & Côté, I. M. (2012). Invasive lionfish drive Atlantic coral reef fish declines. *PloS one*, 7(3), e32596.  
<https://doi.org/10.1371/journal.pone.0032596>
- Grijalba-Bendeck, M., Castañeda-Moya, E. & Acero, A. (2004). Estructura de un ensamblaje íctico asociado a fondos duros en el Caribe colombiano empleando la técnica del censo visual estacionario (CVE). *Actualidades Biológicas*, 26(81), 197-211.
- Guzmán, G. R. & Leal, J. (2011). *Lista de peces asociados a formaciones coralinas en la bahía Capurganá (Chocó) y algunos descriptores de su estructura comunitaria*. (Trabajo de grado). Turbo, Antioquia: Programa de Ecología de Zonas Costeras, Corporación Académica Ambiental, Universidad de Antioquia. 26 pp.
- Hill, J. & Wilkinson, C. (2004). *Methods for ecological monitoring of coral reefs*. Townsville: Australian Institute of Marine Science. 117 pp.
- Humann, P. & Deloach, N. (2003). *Reef fish identification: Florida, Caribbean, Bahamas*. Florida: New World Publications. 481 pp.
- Kulbicki, M., Parravicini, V., Bellwood, D. R., Arias-González, E., Chabanet, P., Floeter, S. R. & Mouillot, D. (2013). Global biogeography of reef fishes: a hierarchical quantitative delineation of regions. *PLoS One*, 8(12), e81847.  
<https://doi.org/10.1371/journal.pone.0081847>
- Manzello, D. P. (2010). Coral growth with thermal stress and ocean acidification: lessons from the eastern tropical Pacific. *Coral reefs*, 29(3), 749-758.  
<https://doi.org/10.1007/s00338-010-0623-4>
- McLaughlin, C. J., Smith, C. A., Buddemeier, R. W., Bartley, J. D. & Maxwell, B. A. (2003). Rivers, runoff, and reefs. *Global and Planetary Change*, 39(1-2), 191-199.  
[https://doi.org/10.1016/S0921-8181\(03\)00024-9](https://doi.org/10.1016/S0921-8181(03)00024-9)
- Mejía, L. S. & Garzón-Ferreira, J. (2000). Estructura de comunidades de peces arrecifales en cuatro atolones del Archipiélago de San Andrés y Providencia (Caribe suroccidental). *Revista de Biología Tropical*, 48 (4), 883-896.
- Morris, J. A., Akins, J. L., Barse, A., Cerino, D., Freshwater, D. W., Green, S. J. & Whitfield, P. E. (2009). *Biology and ecology of the invasive lionfishes, Pterois miles and Pterois volitans*. Marathon F.L. Gulf and Caribbean Fisheries Institute. 61 pp.
- Muñoz-Escobar, L. & Gil-Agudelo, D. L. (2012). Dietary composition of the lionfish, *Pterois volitans* (pisces: scorpaenidae), in Santa Marta and Tayrona national park. *Boletín de Investigaciones Marinas y Costeras-INVEMAR*, 41(2), 471-477.
- Myers, N., Mittermeier, R. A., Mittermeier, C. G., da Fonseca A. B. & Kent, J. (2000). Biodiversity hotspots for conservation priorities. *Nature*, 403, 853-858.  
<https://doi.org/10.1038/35002501>
- Nirchio, M., Oliveira, C., Siccha-Ramírez, Z. R., de Sene, V. F., Sola, L., Milana, V. & Rossi, A. R. (2017). The *Mugil curema* species complex (Pisces, Mugilidae): a new karyotype for the Pacific white mullet mitochondrial lineage. *Comparative cytogenetics*, 11(2), 225-237.
- O'Dea, A., Hoyos, N., Rodríguez, F., Degracia, B. & De Gracia, C. (2012). History of upwelling in the tropical Eastern Pacific and the paleogeography of the Isthmus of Panama. *Palaeogeography Palaeoclimatology Palaeoecology*, 348-349, 59-66.  
<https://doi.org/10.1016/j.palaeo.2012.06.007>
- Peláez de la Torre, S. (2010). *Comparación temporal de la estructura de dos comunidades ícticas de arrecifes someros en las bahías de Capurganá y en la ensenada del aguacate Caribe-Chocó*. (Trabajo de grado). Bogotá D.C.: Pontificia Universidad Javeriana, Facultad de Ciencias, Departamento de Biología. 65 pp.
- Pinheiro, H. T., Rocha, L. A., Macieira, R. M., ..., Floeter, S. R. (2018). South-western Atlantic reef fishes: Zoogeographical patterns and ecological drivers reveal a secondary biodiversity centre in the Atlantic Ocean. *Diversity and Distributions*, 24(7), 951-965.
- Ramírez J. J. & Gaviria G. (2013). *Catálogo de peces asociados a los arrecifes de coral de bahía Capurganá-Choco (Colombia), una aproximación a su conocimiento*. (Trabajo de grado). Turbo, Antioquia: Programa de Ecología de Zonas Costeras, Corporación Académica Ambiental, Universidad de Antioquia. 123 pp.
- Restrepo, J. D. & Kjerfve, B. (2004). The Pacific and Caribbean rivers of Colombia: water discharge, sediment transport and dissolved loads. In de Lacerda,



- L. D., Santelli, R. E., Duursma, E. K. & Abrao, J. J. (Eds.), *Environmental geochemistry in tropical and subtropical environments*. Pp. 169-187. Berlin, Heidelberg: Springer.
- [https://doi.org/10.1007/978-3-662-07060-4\\_14](https://doi.org/10.1007/978-3-662-07060-4_14)
- Reyes-Nivia, M. C., Rodríguez-Ramírez, A. & Garzón-Ferreira, J. (2004). Peces asociados a formaciones coralinas de cinco áreas del Caribe colombiano: listado de especies y primeros registros para las áreas. *Boletín de Investigaciones Marinas y Costeras-INVEMAR*, 33(1), 101-115.
- <https://doi.org/10.25268/bimc.invemar.2004.33.0.250>
- Robertson, D. R. & Cramer, K. L. (2014). Defining and dividing the greater Caribbean: insights from the biogeography of shore fishes. *PLoS One*, 9 (7), e102918.
- <https://doi.org/10.1371/journal.pone.0102918>
- Rocha, L. A., Robertson, D. R., Rocha, C. R., Van Tassell, J. L., Craig, M. T. & Bowen, B. W. (2005). Recent invasion of the tropical Atlantic by an Indo-Pacific coral reef fish. *Molecular Ecology*, 14 (13), 3921-3928.
- <https://doi.org/10.1111/j.1365-294X.2005.02698.x>
- Rojas-Vélez, S., Tavera, J. & Acero, A. (2019). Unraveling lionfish invasion: Is *Pterois volitans* truly a morphologically novel predator in the Caribbean? *Biological Invasions*, 21(6), 1921-1931.
- <https://doi.org/10.1007/s10530-019-01946-6>
- Schofield, P. J. (2009). Geographic extent and chronology of the invasion of non-native lionfish (*Pterois volitans* [Linnaeus 1758] and *P. miles* [Bennett 1828]) in the Western North Atlantic and Caribbean Sea. *Aquatic Invasions*, 4, 473-479.
- <https://doi.org/10.3391/ai.2009.4.3.5>
- Starck, W. A. (1968). A list of fishes of Alligator Reef, Florida with comments on the nature of the Florida reef fish fauna. *Undersea Biology*, 1(1), 4-40.
- Valdez-Moreno, M., Quintal-Lizama, C., Gómez-Lozano, R. & del Carmen García-Rivas, M. (2012). Monitoring an alien invasion: DNA barcoding and the identification of lionfish and their prey on coral reefs of the Mexican Caribbean. *PloS one*, 7(6), e36636.
- <https://doi.org/10.1371/journal.pone.0036636>
- Van der Laan, R., Fricke, R. & Eschmeyer, W. N. (eds). (2021). Catalogue of fishes: classification. California Academy of Sciences. Available from: <http://www.calacademy.org/scientists/catalog-of-fishes-classification/> (Accessed 10 Feb 2021).
- Victoria, P. & Gómez, D. P. (1984). Nuevos registros de peces para la isla de San Andrés (mar Caribe de Colombia). *Anales del Instituto de Investigaciones Marinas de Punta Betín*, 14, 115-132.
- Weigmann, S. (2016). Annotated checklist of the living sharks, batoids and chimaeras (Chondrichthyes) of the world, with a focus on biogeographical diversity *Journal of Fish Biology*, 88(3), 837-1037.
- <https://doi.org/10.1111/jfb.12874>
- Westneat, M. W. & Alfaro, M. E. (2005) Phylogenetic relationships and evolutionary history of the reef fish family Labridae. *Molecular Phylogenetics and Evolution*, 36, 370-390.
- White, J., Simpfendorfer, C. A., Tobin, A. J. & Heupel, M. R. (2013). Application of baited remote underwater video surveys to quantify spatial distribution of elasmobranchs at an ecosystem scale. *Journal of Experimental Marine Biology and Ecology*, 448, 281-288.
- <https://doi.org/10.1016/j.jembe.2013.08.004>



**Camilo Escobar-Sierra**

Universidad de Antioquia.

Medellín, Colombia.

<https://orcid.org/0000-0001-9105-4378>

[camilo.escobar@udea.edu.co](mailto:camilo.escobar@udea.edu.co)

Autor para correspondencia

**Viviana Márquez Velásquez**

Universidade Federal da Paraíba.

João Pessoa, Brasil.

<https://orcid.org/0000-0002-1205-7720>

[vmarquez@squalus.org](mailto:vmarquez@squalus.org)

**Rafael Menezes**

Universidade Federal da Paraíba.

João Pessoa, Brasil.

<https://orcid.org/0000-0003-2378-3805>

[rafaelmenez@gmail.com](mailto:rafaelmenez@gmail.com)

**Ricardo Souza Rosa**

Universidade Federal da Paraíba.

João Pessoa, Brasil.

<https://orcid.org/0000-0002-4289-2241>

[rsrosa@dse.ufpb.br](mailto:rsrosa@dse.ufpb.br)

**Alejandro Loaiza-Santana**

Universidad de Antioquia.

Medellín, Colombia.

<https://orcid.org/0000-0003-2608-8218>

[alejandro.loaiza.san@gmail.com](mailto:alejandro.loaiza.san@gmail.com)

---

**An updated reef fish checklist of the southernmost Caribbean reef system, with comments on the lionfish invasion**

**Citación del artículo:** Escobar-Sierra, C., Márquez-Velásquez, V., Menezes, R., Souza-Rosa, R. Loaiza-Santana, A. (2021). An updated reef fish checklist of the southernmost Caribbean reef system, with comments on the lionfish invasion. *Biota Colombiana*, 22(2), 70-87.

<https://doi.org/10.21068/c2021.v22n02a04>

**Recibido:** 27 de septiembre 2020

**Aceptado:** 2 de marzo 2021