

OCCURRENCE OF *SELLITHYRIS SELLA* (BRACHIOPODA) IN THE ROSABLANCA FORMATION, BOYACÁ-COLOMBIA

Alexis Rojas-Briceño, Pedro Patarroyo

ABSTRACT

The brachiopod shells discovered in the Rosablanca Formation from the Santa Sofia locality, NW of Villa de Leyva, Boyacá province, can be identified as *Sellithyris sella* (J. de C. Sowerby, 1823). The studied material shows characteristic features of this taxon including the loop wide, triangular with high-arched transverse band. This material was collected during a paleontological excursion with geology students. *Sellithyris* was previously reported from Zapatoca, Santander province by Dietrich (1938) and Karsten (1958). This new record documents its wide Late Valanginian-?Hauterivian distribution in Colombia (Uppermost Rosablanca Formation) and confirms the direct transatlantic connections with Western Europe.

Key words: *Sellithyris sella*, Rosablanca Formation, Late Valanginian - ? Hauterivian, Boyacá-Colombia.

REGISTRO DE *SELLITHYRIS SELLA* (BRACHIOPODA) EN LA FORMACIÓN ROSABLANCA, BOYACÁ-COLOMBIA

RESUMEN

Se describe la ocurrencia de *Sellithyris sella* (J. de C. Sowerby, 1823) en la Formación Rosablanca, proveniente del área de Santa Sofia, Departamento de Boyacá. El material estudiado comprende dos ejemplares completos y seis moldes internos colectados en una excursión con estudiantes de geología. El *brachidium* amplio, triangular y provisto de una banda transversa alta, preservado en uno de los moldes internos, es característico del taxón. *Sellithyris* ha sido registrado previamente en el área de Zapatoca, Departamento de Santander. Este nuevo registro indica su amplia distribución durante el Valanginiano Tardío-? Hauteriviano de Colombia y confirma las conexiones con Europa Occidental.

Palabras clave *Sellithyris sella*, Formación Rosablanca, Valanginiano Tardío -? Hauteriviano, Boyacá- Colombia.

INTRODUCCIÓN

The Colombian Cretaceous brachiopods include the terebratulids: *Arenaciarcula beaumonti* (d'Archaic) and *Gemmarcula* cf. *menardi* (Lamarck) deposited in the d'Orbigny Collection, and *Sellithyris sella* (J. de C. Sowerby, 1823), reported as *Terebratula Haueri* Karsten (1958) and *Terebratula sella* Dietrich (1938) from Zapatoaca, Santander Province. This study documents the occurrence of *Sellithyris sella* (J. de C. Sowerby,

1823) in a stratigraphic section with good outcrop exposure of the Upper Rosablanca Formation from Santa Sofia (FIGURE 1), near of the "la Fábrica" cave, Boyacá province. This genus is a low latitude Tethyan form (Sandy, 1997), and is present in Western Europe, North Africa, North America and Mexico (Sandy, 1997; Gaspard, 2005). Manceñido (2002) proposed that the occurrence of *Sellithyris* in Colombia and *Musculina* from northern? South America may suggest direct connections with Western Europe.




FIGURE 1. Localization of the studied area to the east of Santa Sofia, Boyacá-Colombia. Illustration from <http://maps.google.com/maps?client=firefox-a&rls=org.mozilla:es>

MATERIAL AND METHODS

The brachiopod shells were examined under a camera lucida and transverse serial sections were obtained by gridding one internal mold with 600 powder on glass plate. The dimensions (in mm): length dorsal valve (ldv), length (l), maximum width (mw), position of maximum width (pmw) and thickness (t) were taken with a caliper. All material used for this report kept in the palaeontological collections in the Departamento de Geociencias, Universidad Nacional de Colombia, Bogotá.

SYSTEMATIC PALEONTOLOGY

Order *Terebratulida* WAAGEN, 1883
Suborder *Terebratulidina* WAAGEN, 1883
Superfamily *Terebratuloidea* GRAY, 1840
Family *Sellithyrididae* MUIR-WOOD, 1965
Subfamily *Sellithyridinae* MUIR-WOOD, 1965
Genus *Sellithyris* MIDDLEMISS, 1959

Sellithyris sella (J. de C. Sowerby, 1823)
(FIGURE 2)




FIGURE 2. *Sellithyris sella* Rosablanca Formation, Santa Sofia, Boyacá Province. a. Dorsal view; b. Frontal view. UN-DG-BR-1002.

Description: Pentagonal shell, smooth, maximum wide slightly anterior mid line, anterior commissure episulcate, median sinus smaller and narrower than the lateral, umbo sub-erect. Loop wide and triangular with

high-arched transverse band. Other internal features are not observed. Dimensions in the TABLE 1.

Material examined: Two complete specimen and six internal molds.

TABLE 1. Dimensions (in mm). Length Dorsal valve (LDV), length (L), Maximum width (MW), Position of maximum width (PMW) and Thicknes (T).

Specimen	LDV	L	MW	PMW	T
UN-DG-BR-1002	12,4	16,0	14,1	9,3	8,0




FIGURE 3. Selected transverse serial sections illustrating the development of the loop of *Sellithyris sella*, a-c Rosablanca Formation, Santa Sofia, Boyacá Province. The sections were obtained by gridding the internal mold with 600 powder on glass plate. The distances were measured in mm from the posterior end of the internal mold. a, high-arched transverse band; b and c, descending lamellae; d-e loop reconstruction of *S. sella* modified from Middlemiss (1959). .

IMPLICATIONS

The occurrence of *Sellithyris sella* in the Rosablanca Formation from Zapatoca, Santander Province, and Santa Sofia, Boyacá province suggest that it was widely distributed during Late Valanginian–?Hauterivian in Colombia and confirms the direct transatlantic connections with Western Europe.

Recently, Gaspard (2005) observed shape variations in *Sellithyris cenomanensis* from Sarthe, Charente-maritime, Indre et Loire, Maine et Loire, Vendée and S.E. France, associated to the position during the Cenomanian boreal and mesogean provinces. Gaspard (2005) suggested that the brachiopods react quickly to the environmental modification to explain the observed variations. Then, the extensive morphological study of *Sellithyris* could improve the understanding of the relationship between Andean and Tethyan realms.

Geological Society of America and Kansas University Press, Lawrence, Kansas, pp. 762-816.

Sandy, M. R. 1990. Biogeographic affinities of some Jurassic-Cretaceous brachiopod faunas from the Americas and their relation to tectonic and paleoceanographic event. In: MacKinnon, D.I., Lee, D.E., Campbell, J. D. (Eds.), *Brachiopods through Time*. A.A. Balkema, Rotterdam, pp. 415-422.

Sandy, M. R. 1997. Mesozoic brachiopods of Mexico- A review and some comments on their paleobiogeographic affinities and paleoecology. *Revista Mexicana de Ciencias Geológicas*, 14 (2): 219-224.

REFERENCIAS

Dietrich, W. 1938. Lamelibranquios cretácicos de la Cordillera Oriental. En. Scheibe, R.: *Estudios geológicos y paleontológicos sobre la Cordillera Oriental de Colombia*. Minist. Ind. y Petrol., (2), 7-80 Parte tercera, pp. 81-108, pl. 2. Bogotá.

Gaspard, D. 2005. Shape variation in Cretaceous biplicate terebratulid brachiopods. Impact environnemental sur la dynamique des populations au Crétacé. Réunions thématiques du Groupe Français du Crétacé. Paris. 24. p.

Grădinaru, E.; Bárbulescu, A.; Owen, E. 2006. Latest Albian (Vraconian) brachiopod fauna North Dobrogea (Romania): Taxonomy, palaeoecology and palaeobiogeography. *Acta Geologica Polonica*, 56 (1): 67-88.

Karsten, H. 1858. Über die geognostischen Verhältnisse des westlichen Columbien, der heutigen Republiken Neu-Granada und Ecuador. Amtl. Ber. über die 32 Versammlg. d. Deutsch. Naturforsch. in Wien 1856, Viena.

Manceñido, M. O. 2002. Paleobiogeography of Mesozoic brachiopod faunas from Andean-Patagonian areas in a global context. *Geobios*, 35 (17):176-192.

Middlemiss, F. A. 1959. English Aptian Terebratulidae. *Paleontology*, 2(1): 94-142.

Muir-Wood, H. M. 1965. Mesozoic and Cenozoic Terebratulidina. In Moore, R. C. (ed.): *Treatise on Invertebrate Paleontology*, part H (Brachiopoda).

Trabajo recibido: Junio 23 de 2009

Trabajo aceptado: Octubre 16 de 2009