

Molt strategy and aging of Rufous-collared Sparrow (*Zonotrichia capensis*) in Bogotá, Colombia

Estrategia de muda y determinación de edad del copetón
(*Zonotrichia capensis*) en Bogotá, Colombia

DIEGO CUEVA

Universidad Nacional de Colombia (sede Bogotá). Facultad de Ciencias. Instituto de Ciencias Naturales, Grupo de Ornitología y Colección de ornitología, Cra. 30 # 45 – 03, edificio 425. Bogotá, Colombia. Current Address: Seção de aves, Museu de Zoologia da Universidade de São Paulo, Pós-Graduação, CEP 04218-970, São Paulo, SP, Brasil. dacuevac@usp.br

ABSTRACT

Molting is an important process of birds' natural history and learning about molt and plumage allows a better understanding of various aspects of birds' life. In temperate zones, knowledge about molt strategy, type of plumage characteristics and their association with age are well documented. In contrast, knowledge about molt in Neotropical birds is not well developed. One species with numerous studies regarding molt in several localities is the widely-distributed Rufous-collared Sparrow (*Zonotrichia capensis*). These studies covered various localities along the species range, facilitating comparisons among populations, and providing an excellent model for the study of plumage and molt variation. In this study, I described both molt strategy and extent of molt *Z. capensis costaricensis*, through sampling data for one year at two localities in Bogotá city. I captured and banded *Z. capensis* individuals and collected information about characteristics such as molt limits, skull ossification and iris coloration, which allow for the identification of molt/plumage type and its relation to age categories of a cycle-based age classification system. I found that *Z. capensis* has a complex basic strategy, where all prebasic molts, including prejuvenal molt, are complete and the preformative molt is a partial molt. The results are congruent with previous findings for this species in some areas. Nevertheless, the molt varies largely in *Z. capensis* across its distribution and individuals out of Neotropics have an additional partial-incomplete prealternate molt and it is still unknown why.

Key Words. Iris color, plumage, age classification, Neotropics, W–R–P system.

RESUMEN

La muda es un proceso importante de la historia natural de las aves y aprender sobre este proceso permite entender varios aspectos de su vida. En zonas templadas, el conocimiento sobre las estrategias de muda, las características de cada tipo de plumaje y su relación con la edad del ave, son bien conocidas. Por el contrario, en el neotrópico, este conocimiento no está bien desarrollado. Una especie con bastantes estudios sobre muda en varias localidades es el ampliamente distribuido Copetón (*Zonotrichia capensis*). Esos estudios cubren localidades de su ámbito geográfico facilitando comparaciones entre poblaciones y proveyendo un excelente modelo para estudiar la variación del plumaje y la muda. En este estudio, determiné el alcance y la estrategia de muda de *Z. capensis costaricensis* mediante muestreos durante un año en dos localidades de Bogotá. Capturé y anillé individuos de *Z. capensis* y tomé información de características como los límites de muda, osificación del cráneo y color del iris que me permitieron identificar el tipo de plumaje/muda y relacionarlo con una categoría de edad del sistema de clasificación de edad basado en los ciclos de muda. Encontré que *Z. capensis* tiene una estrategia de muda básica compleja, donde todas las mudas prebásicas, incluida la prejuvenil, son completas y la muda

preformativa es parcial. Estos resultados son congruentes con los de otras regiones. Sin embargo, la muda de *Z. capensis* varía mucho, e individuos lejos de los trópicos tienen una muda alterna parcial-incompleta adicional y aún desconocemos por qué.

Palabras Clave. Color del iris, plumaje, clasificación de edad, neotrópico, sistema W–R–P.

INTRODUCTION

Molting, an energy demanding activity, is one of the most important processes in birds' life cycle (Howell *et al.* 2003, Echeverry-Galvis and Córdoba-Córdoba 2008). Learning about plumage variation, molting, sexing and aging in birds, permit us to understand their ecology and plumage evolution. Furthermore, with long-term monitoring, they will allow a greater understanding of species demography, contributing to more effective decisions on bird conservation (Gómez *et al.* 2012).

Molt strategies, especially for passerines, are well documented in temperate zones (see Jenni and Winkler 1994, Pyle 1997, 2008). However, patterns present in temperate zones do not completely coincide with those of Neotropical birds, varying in duration and seasonality (Wolf 1969, Ryder and Wolfe 2009). This could be partly due to differences in resource availability throughout the year between these regions (e.g. Miller 1961, Poulin *et al.* 1992, Echeverry-Galvis and Hau 2013). The new W-R-P system permits classifying tropical birds into age classes (Wolfe *et al.* 2010, Johnson *et al.* 2011) and eases the understanding of plumage and molts in tropical birds and leaves an open door for ornithological research (Hernandez 2012, Johnson and Wolfe 2014). Specifically, in Colombia, research about molt and age criteria is a growing research area (e.g. Botero-Delgadillo *et al.* 2012, Hernández 2012, Gómez *et al.* 2012, Moreno-Palacios *et al.* 2013).

The molt of *Z. capensis* (Müller, 1776) has been fairly well investigated with multiple studies in Colombia and Neotropics (Miller

1961, Wolf 1969, Davis 1971, King 1972, 1973, Zerda *et al.* 1986, Pyle *et al.* 2015). Because of this, comparisons can be made among locations, latitudes and populations. In Colombia, the most detailed research about plumage, molt and breeding of *Z. capensis* was carried out near Cali, Valle del Cauca (Miller 1961). This study established the molt strategy, a brief description of different plumages such as juvenal, formative, basic, and estimates the duration of molt and reproduction seasons for the species.

There exist various studies about plumage, molt and breeding in *Z. capensis*. The existing information and its broad distribution in Central and South America make of *Z. capensis* an excellent model to understand the variation of duration and timing in this process in the tropical region. *Z. capensis* is a widely distributed species and its molt strategy varies across its distribution, even between populations of the same subspecies. In this study, I documented both, strategy and extent of molt in *Z. capensis* in Bogotá, describing new external characteristics useful for age determination in Bogotá, as iris color. Also, I compared my results with previous investigations and propose the hypothesis of molt variation across latitudinal gradient.

MATERIALS AND METHODS

The study area is located in the Eastern Cordillera of the Colombian Andes, in the city of Bogotá D.C. (2560 m a.s.l.). I established a total of five sampling points, two in the Bogotá Botanical Garden-José Celestino Mutis (4°40'North, 74°6'West) and three in the campus of the Universidad Nacional

de Colombia (4°38'North, 74°5'West). I conducted 65 mist net samplings using 12 m nets for a total of thirteen samplings per site, between 06:00 and 11:00 h, totaling 1883.5 net hours. Samplings occurred from the 1st Apr 2013 to the 5th Apr 2014. Each site was sampled every four weeks in zones with low vegetation and grasslands.

Molt strategy and pattern

Birds were banded, allowing for identification and tracking of individuals over time. I took detailed notes about plumage, body molt, flight feathers molt, molt limits, shape of external retrices, iris color categorized following Smithe's color guide (1975) and skull ossification (Ralph *et al.* 1996, Pyle 1997). Ossification was determined following the methodology of Pyle (1997) assigning a percentage of the total area of the skull. In this methodology, 0 % means a skull with a single layer of bone and 100 % means a skull with two layers and complete pneumatization. The terminology used for plumage and molt follows Humphrey and Parkes (1959) modified by Howell *et al.* (2003).

Ageing, molt and plumage description

The bird's age was determined with criteria such as: the presence/absence of brown streaks on the chest of juveniles (Miller 1961), molt limits, shape of flight feathers, sharp in juvenal feathers, broad in individuals with basic feathers, growth bars, aligned in juvenal flight feathers and not aligned in basic feathers (Pyle 1997). Additionally, I took notes describing other useful age characteristics of *Z. capensis* like iris color and skull ossification. I used the mentioned criteria to assign every bird to an age category, following the molt cycle-based age-classification system – WRP (Wolfe *et al.* 2010, Johnson *et al.* 2011).

RESULTS

Molt strategy and molt pattern

I banded 102 Rufous-collared Sparrows and recaptured 42 during the sampling year. I found that besides the basic plumage, *Z. capensis* counts with a formative plumage. Also, I did not find traces of alternate plumage, as it will be discussed later. Thus, the molt strategy of *Z. capensis* in Bogotá is a Complex Basic Strategy (CBS; *sensu* Howell *et al.* 2003). The prebasic molts present a complete molt pattern, i.e. molting all the feathers in the same molting process, and the formative plumage is the result of a partial molt in which flight feathers, primaries, secondaries, tertials, retrices and primary coverts, are not molted. Nevertheless, some captured individuals with a formative plumage had molted tertial feathers (30 %) or central retrices (17 %).

Ageing, molt and plumage description

In Bogotá, I categorized *Z. capensis* in eight age classes within the WRP system (Wolfe *et al.* 2010, Johnson *et al.* 2011). The first category, first-cycle prejuvenal molt (FPJ), occurs in the nest when the bird substitutes the natal down plumage for the juvenal. I did not sample nests and I do not have information about the FPJ. The first-cycle juvenal (FCJ) birds are the easiest to identify. They have a juvenal plumage which highly contrasts with posterior plumages. Juvenal plumage presents cryptic colors, brown streaks on stomach and breast and the absence of the specie's characteristic rufous collar (Fig. 1a). Moreover, 90 % of FCJ birds showed a skull ossification between 0 and 20 % and a raw amber iris color (Table 1; Fig. 2a).

After FCJ, *Z. capensis* reaches a first-cycle preformative molt (FPF). The molt in FPF

Table 1. Distribution of the captured birds in the categories of iris color and skull pneumatization. Bold numbers highlight the higher percentage values at each age class. Some percentages may not reach 100 % because I could not conduct a proper inspection of some captured birds (n). i.e. molting shades may difficult the observation of the ossification.

W-R-P System	n	Iris color (Figure 2)				Skull pneumatization (%)					
		a	b	c	d	0-20	20-40	40-60	60-80	80-95	95-100
FPJ	0										
FCJ	10	100 %				90 %	10 %				
FPF	6	50 %	50 %			50 %	33 %				
FCF	26		77 %	23 %		8 %	54 %	23 %	8 %	4 %	
SPB	22		18 %	68 %	14 %			5 %		59 %	9 %
DCB (SCB)	12		17 %	67 %	17 %				17 %	50 %	33 %
DPB	25			4 %	88 %						88 %
DCB	18			11 %	83 %						94 %




Figure 1. Plumages of Rufous-collared Sparrow (*Z. capensis costaricensis*) in Bogotá, Colombia. **a.** Juvenile plumage, **b.** Formative plumage, and **c.** Basic plumage. Pictures by Diego Cueva.

age does not include flight feathers. FPF birds presented molting body feathers, brown juvenal feathers on head and remaining streaks on chest, skull ossification under 40 % with a majority under 20 % (Table 1) and a raw amber/chestnut iris color (Fig. 2b). The acquired formative plumage named the next age class, first-cycle formative

(FCF; Fig. 1b). As expected (Miller 1961, 1962), formative plumage is superficially similar to later basic plumages (Fig. 1c). Thus, body feathers became useless to classify individuals into an age class. FCF Rufous-collared Sparrow molted all secondary coverts but not primary coverts, they exhibited molt-limits between greater

coverts and primaries coverts. Juvenal primary coverts had an external buffy (Fig. 3a) color that almost disappears in posterior moltings (Fig. 3b). Also, external retrices (juvenal) were narrower shaped, and flight feathers presented aligned growth bars. The preformative molt extent is similar to the one described for *Z. capensis* in other regions (Pyle *et al.* 2015) and for other *Zonotrichia* species (Pyle 1997). Furthermore, 77 % of captured FCF individuals had ossification between 20 and 60 %, and 54 % of the total FCF birds did not get over 40 % ossification. Finally, 77 % FCF individuals presented a chestnut colored iris, and 23 % presented an intermediate iris color, between chestnut and definitive Pratt's Rufous color (Table 1; Fig. 2b).

After FCF, *Z. capensis* turned into a second-cycle prebasic molt (SPB) individual. At this stage, 64 % of the individuals still have skull windows, most of them with skull ossification between 80 and 95 %. Also, 68 % of the birds presented an intermediate iris color between chestnut and Pratt's Rufous (Fig. 2c). Finally, juvenal primaries, secondaries, retrices and primary coverts can be identified before they are completely molted, according to the descriptions of the FCF birds. After second prebasic molt, *Z. capensis* enters into the second-cycle basic (SCB) age class. Birds in this stage did not show differences with subsequent plumages. Nevertheless, some individuals, presumably at the beginning of the SCB, presented skull windows and iris that did not reach a Pratt's Rufous coloration yet (Fig. 2c) permitting age identification. However, at this stage, *Z. capensis* develops the final iris color, Pratt's Rufous, and completes skull ossification, making birds indistinguishable from birds in definitive-cycle basic (DCB). This forces categorization as after first cycle (FAJ), following Wolfe *et al.* (2010) and Johnson *et al.* (2011).

Finally, definitive prebasic molt (DPB) birds differ from previous molt stages because the primaries, secondaries and primaries coverts are not juvenal and do not present the characteristics described in the FCF section. Furthermore, the individuals have a Pratt's Rufous iris (Fig. 2d) and a completely ossified skull. The definitive-cycle basic (DCB) birds have a plumage that will not change in posterior cycles. Moreover, they present a Pratt's Rufous iris color and 100 % skull ossification.

Additionally, I captured several times an individual 7.4 years old, first banded by the Ornithology Group of Universidad Nacional de Colombia (GOUN) on the 20th Sep 2006 and last time captured on the 16th Feb 2014. This bird allowed me to confirm the characteristics of DCB. Also, outstanding captures were two FCF females with high feather wear, category three over four, and developed brood patch. Moreover, seven males with suspended molt (P2-3), all with a cloacal protuberance. These captures highlight that in Bogotá *Z. capensis* reach sexual maturity and can reproduce at an early age class, during the FCF. Finally, the seven males are evidence of molt suspension due to overlapping with reproduction. Similar cases were found in Tucumán, Argentina (King 1972, 1973).

DISCUSSION

In Bogotá *Z. capensis* has a Complex Basic Strategy. All prebasic molts, including pre juvenal molt, are complete molts. Only the preformative molt is a partial molt. The strategy and extent of the molts in Bogotá do not differ from the populations around Cali, Colombia (~3° North; Miller 1961). Furthermore, both populations did not show evidence of alternate plumage, differently from several studies occurred at higher




Figure 2. Iris color variation throughout different age stages. Colors according to [Smithe \(1975\)](#). **a.** Color 223 Raw umber, **b.** Color 32 Chestnut, **c.** Intermediate between 32 and 140 colors, and **d.** Color 140 Pratt's Rufous. Pictures by Diego Cueva (a, b) and Julian Avila-Campos (c, d).

latitudes such as Costa Rica ($\sim 10^\circ$ North, [Wolf 1969](#)), Peru ($\sim 12^\circ$ South, [Davis 1971](#)) and Chile ($\sim 30^\circ$ South, [Pyle et al. 2015](#)), where an alternate plumage was present. However, studied populations of *Z. capensis* in Tucumán, Argentina ([King 1972, 1973](#)) did not show prealternate plumage either.

A second important fact about molt of *Z. capensis* is that reported by [Miller \(1961\)](#) indicating individuals going through two

complete prebasic molts in a single calendar year. Is it possible that the incomplete alternate molt reported in Peru by [Davis \(1971\)](#), which included 2-6 primaries, became a complete molt in tropical latitudes? If it is a complete prealternate molt, there is no way to differentiate it from a complete prebasic molt. Even if two molting peaks occur in Cali and Bogotá during the same year ([Miller 1961, Zerda et al. 1986](#)), neither of them may be associated with prebasic



Figure 3. Color variation in primary coverts of *Z. capensis* in Bogotá. **a.** A FPF individual with buffy color on the external border (arrow) of the retained primary coverts and, **b.** a DCB individual without the buffy border after molting the same group of feathers.

or prealternate molt, due to the existence of complete molts all year around (Miller 1961, Zerda *et al.* 1986). Differently to temperate areas, where most birds respond to an annual climate cycle (Pyle 1997), in the Neotropics *Z. capensis* is subjected to a two-cycle climate year (Miller 1961, Zerda *et al.* 1986). Here, energy consuming activities such as reproduction have two seasons per year, and a single individual can be part of both (Miller 1961, 1962). I think, there are reasons to believe that *Z. capensis* has two complete prebasic molts, instead of one prebasic and one prealternate molts. Nevertheless, the discussion remains open.

Finally, I want to underline that females are sexually mature at the FCF age. In temperate zones, for most birds, there is only one favorable season for reproduction, when several resources become available. Thus, hatching individuals wait for a complete year to the next reproduction season. Contrastingly, the Neotropics have two climate cycles per year and some species, as *Z. capensis*, have two reproductive seasons (Miller 1962). This way, hatched birds concur with a reproductive peak that occurs approximately six months later, and if they reach sexual maturity, they will start to reproduce. However, subsequent molts will

be delayed due to the high energy demand during reproduction and this will lead to a trade-off between good quality plumage and early reproduction.

There is not clarity regarding which environmental factors in the Neotropics trigger the physiological process that leads to seasonal reproduction. However, [Wikelski et al. \(2000\)](#) presented evidence that tropical birds can respond to photoperiod variation and used it to synchronize reproduction.

CONFLICT OF INTEREST

The author declares that there is no conflict of interests.

ACKNOWLEDGEMENTS

I am grateful to F. Gary Stiles for his orientation and support in all aspects of this research. I am thankful to Idea Wild for their funding support. I thank Juan Pablo López, Alejandro Pinto, Rafael Gutiérrez and Raúl Sedano for supplying materials. I thank the Ornithology Group of the Universidad Nacional - GOUN (<https://gounaves.wordpress.com/integrantes/>), Sebastian Mantilla and Darwin Morales for helping me with the morning samplings. I am grateful to Bogotá's Jose Celestino Mutis botanical garden for permission to sample and Julián Ávila for helping with it. Finally, I thank Orlando Acevedo, Diego Carantón, Courtney Dwyer, Anna Ferraroni and Gustavo Bravo for the comments on this document.

LITERATURE CITED

Botero-Delgado E, Bayly N, Gómez-Montes C. 2012. Ciclos de muda en *Diglossa albilatera* (Thraupidae) y evidencia de un patrón en la adquisición del dicromatismo sexual. *Boletín SAO* 21:1–7.

Davis J. 1971. Breeding and molt schedules of the Rufous-collared Sparrow in coastal Peru. *Condor* 73:127–146.

Echeverry-Galvis MÁ, Córdoba-Córdoba S. 2008. Una visión general de la reproducción y muda en aves en el Neotrópico. *Ornitol. Neotrop.* 19:197–205.

Echeverry-Galvis MÁ, Hau M. 2013. Flight performance and feather quality: paying the price of overlapping moult and breeding in a tropical highland bird. *Plos One* 8(5):1–9. doi: 10.1371/journal.pone.0061106.

Gómez C, Botero-Delgado E, Bayly NJ, Moreno MI, Páez CA. 2012. Documentando estrategias de muda en aves Neotropicales: ejemplos de la Sierra Nevada de Santa Marta, Colombia. *Ornitol. Neotrop.* 23:129–138.

Hernández A. 2012. Molt patterns and sex and age criteria for selected landbirds of southwest Colombia. *Ornitol. Neotrop.* 23:215–223.

Howell SNG, Corben C, Pyle P, Rogers DI. 2003. The first basic problem: a review of molt and plumage homologies. *Condor* 105(4):635–653. doi: 10.1650/7225.

Humphrey PS, Parkes KC. 1959. An approach to the study of molts and plumages. *Auk* 76(1):1–31.

Jenni L, Winkler R. 1994. *Moult and aging of European passerines*. Academic Press. New York. 224 pp.

Johnson EI, Wolfe JD. 2014. Thamnophilidae (antbird) molt strategies in a central Amazonian rainforest. *Wilson J. Ornithol.* 126(3):451–462. Doi: 10.1676/13-163.1.

Johnson EI, Wolfe JD, Ryder TB, Pyle P. 2011. Modifications to a molt-based aging system proposed by Wolfe *et al.* (2010). *J. Field Ornithol.* 82(4):422–424. doi: 10.1111/j.1557-9263.2011.00345.x.

King JR. 1972. Postnuptial and postjuvenile molt in Rufous-collared Sparrows in northwestern Argentina. *Condor* 74(1):5–16. doi: 10.2307/1366444.

King JR. 1973. The annual cycle of the Rufous-collared Sparrow (*Zonotrichia capensis*) in three biotopes in north-western Argentina. *J. Zool.* 170(2):163–188. doi: 10.1111/j.1469-7998.1973.tb01373.x.

Miller AH. 1961. Molt cycles in equatorial andean sparrows. *Condor* 63(2):143–161. doi: 10.2307/1365527.

Miller AH. 1962. Bimodal Occurrence of Breeding in an Equatorial Sparrow. *P. Natl. Acad. Sci. USA.* 48(3):396–400. doi: 10.1073/pnas.48.3.396.

- Moreno-Palacios M, Losada-Prado S, Echeverry-Galvis MÁ. 2013. Ciclos de reproducción y muda del Volatinero Negro (*Volatinia jacarina*) y el Semillero Gris (*Sporophila intermedia*) en un matorral secundario al norte del Tolima, Colombia. *Ornitol. Neotrop.* 24:421–431.
- Poulin B, Lefebvre G, Mcneil R. 1992. Tropical avian phenology in relation to abundance and exploitation of food resources. *Ecology* 73(6):2295–2309. doi: 10.2307/1941476.
- Pyle P. 1997. Identification guide to North America birds: Part I. Bolinas: Slate Creek Press.
- Pyle P. 2008. Identification guide to North America birds: Part II. Bolinas: Slate Creek Press.
- Pyle P, Engilis jr. A, Kelt DA. 2015. Manual para la estimar edad y sexo en aves del Parque Nacional Bosque Fray Jorge y Chile central, con notas sobre rangos de distribución y estación reproductiva. Baton Rouge: Special Publication of the Occasional Papers of the Museum of Natural Science.
- Ralph CJ, Geupel GR, Pyle P, Martin TE, Desante DF, Milá B. 1996. Manual de métodos de campo para el monitoreo de aves terrestres. Albany: Pacific Southwest Research Station, Forest Service, U.S. Department of Agriculture, USA.
- Ryder TB, Wolfe JD. 2009. The current state of knowledge on molt and plumage sequences in selected Neotropical bird families: a review. *Ornitología Neotropical* 20(1):1–18.
- Smithe FB. 1975. Naturalist's color guide. The American Museum of Natural History, New York.
- Wikelski M, Hau M, Wingfield JC. 2000. Seasonality of reproduction in a neotropical rain forest bird. *Ecology* 81(9):2458–2472.
- Wolfe JD, Ryder TB, Pyle P. 2010. Using molt cycles to categorize the age of tropical birds: an integrative new system. *J. Field Ornithol.* 81(2):186–194. doi: 10.1111/j.1557-9263.2010.00276.x.
- Wolf LL. 1969. Breeding and molting periods in a Costa Rican populations of the Andean sparrow. *Condor* 71(2):212–219. doi: 10.2307/1366080.
- Zerda E, Rodríguez P, Snow DW. 1986. Periodos de muda del copetón (*Zonotrichia capensis*) en el Jardín Botánico de Bogotá, Bogotá, Colombia. *Pérez-Arbelaezia* 1(3):371–376.

Received: 17/04/2017

Accepted: 21/05/2018