

Original Article

Remifentanil at induction of general anesthesia for cesarean section: Double blind, randomized clinical trial

Remifentanilo en la inducción de la anestesia general para la cesárea: estudio clínico doble ciego, ensayo clínico aleatorizado

Behdad, Shekoufeh^a; Ayatollahi, Vida^a; Harrazi, Hamid^b; Nazemian, Naderali^a; Heiranizadeh, Najmeh^a; Behnam, Baghianimoghadam^a.

^a Shahid Sadoughi Hospital, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

^b Shohaday-e-kargar Hospital, Yazd, Iran

Shekoufeh B, Ayatollahi V, Harrazi H, Nasemian N, Heiranizadeh N, Baghianimoghadam B. Remifentanil at induction of general anesthesia for cesarean section: Double blind. *Colomb.Med.*2013;44(2): 87-91

Article history:

Received Sept 20 2012
 Received in revised form 5
 Oct 2012
 Accepted Apr 2013

Keywords:

Remifentanil, Placebo,
 caesarean.

Palabras clave:

El remifentanilo, Placebo,
 cesárea.

Abstract

Introduction: Remifentanil, with its rapid activity onset and short duration of action, may be more effective than other opioids for providing hemodynamic stability during obstetric anesthesia. However, there is some evidence of adverse effects on neonatal respiratory function. We investigated maternal and fetal effects of remifentanil during cesarean section surgery.

Methods: Eighteen women with singleton term pregnancies, and physical class status of I or II as defined by the American Society of Anesthesia (ASA), who were undergoing general anesthesia for semi-elective cesarean section were randomized into two groups (40 in each group) that received either an intravenous bolus of 0.5 µg/kg remifentanil or the same dose of saline as a placebo. Maternal hemodynamic variables and neonatal umbilical artery pH and Apgar score at first and fifth minutes were evaluated in both groups.

Results: Systolic and diastolic blood pressure were significantly lower after tracheal intubation and skin incision in the remifentanil group as compared with the control group ($p < 0.05$). There were no significant differences regarding heart rate between groups at any time ($p > 0.05$). Apgar scores at first and fifth minutes were not significantly different among groups ($p > 0.05$). No neonate required assisted ventilation or naloxan administration.

Conclusion: Remifentanil may be a safe and effective drug for the induction of general anesthesia and surgical stimulation without subsequent neonatal depression.

Resumen

Introducción: El remifentanilo, con su rápido comienzo de actividad y corta duración de acción, puede ser más eficaz que otros opioides para proporcionar la estabilidad hemodinámica durante la anestesia obstétrica. Sin embargo, hay alguna evidencia de efectos adversos sobre la función respiratoria neonatal. Se investigaron los efectos maternos y fetales de remifentanilo durante la cirugía cesárea.

Métodos: Dieciocho mujeres con embarazo de feto uno y con clasificación Asa I o II (sociedad americana de anestesiología y que fueron sometidas a anestesia general para cesárea semi electiva, se dividieron aleatoriamente en dos grupos (40 mujeres en cada grupo) que recibieron un bolo intravenoso de 0.5 mm/Kg de remifentanil o en el grupo control la misma dosis de solución salina como placebo. Se evaluaron las variables hemodinámicas maternas y pH de la arteria umbilical neonatal y la puntuación de Apgar al primer y quinto minuto en ambos grupos.

Resultados: La presión arterial sistólica y diastólica fueron significativamente más bajas después de la intubación traqueal y la incisión de la piel en el grupo remifentanilo en comparación con el grupo control ($p < 0,05$). No hubo diferencias significativas en cuanto a la frecuencia cardíaca entre los grupos en cualquier tiempo ($p > 0,05$). Las puntuaciones de Apgar a los minutos primero y quinto no fueron significativamente diferentes entre los grupos ($p > 0,05$). Ningún neonato requirió ventilación asistida o administración naloxan.

Conclusión: El remifentanilo es un fármaco seguro y eficaz para la inducción de la anestesia general y la estimulación quirúrgica sin depresión neonatal posterior.

*Corresponding Author:

E-mail Address : behdadsopyusd@yahoo.com (Shekoufeh B), ayatollahivida-tryuior@yahoo.com (Ayatollahi V), harrazihamidtryui@yahoo.com (Harrazi H), naderalidrnazemiknhgtyi@yahoo.com (Nasemian N), heyranajmeshoddarkhane@yahoo.com (Heiranizadeh N), behnam.baghian@gmail.com (Baghianimoghadam B)

Introduction

Cesarean section is a common obstetric surgery and the anesthetic technique selected by anesthesiologists for this kind of operation is very important. While regional anesthesia is the preferred method for cesarean section, in high risk obstetric conditions, such as maternal coagulopathy, cardiac and neurologic diseases and

cases of urgent delivery, general anesthesia is indicated¹. Opioid drugs are not usually given for induction of general anesthesia for cesarean sections because of neonatal respiratory depression²⁻³; however, without opioid administration, patients are at risk of complications, such as increased heart rate and hypertension following laryngoscopy, tracheal intubation and skin incision. These neuroendocrine stressors can be life-threatening in mothers with cardiovascular diseases. In such cases it is necessary to consider a safe and effective analgesic agent for such situations.

The characteristics of Remifentanyl make it a good choice for usage before delivery instead of other opioids. Remifentanyl is a synthetic μ -specific agonist opioid with a rapid onset of action, a short duration to reach peak effect, a rapid clearance and quick fetal metabolism². Also, it can be administered for prevention of unwanted maternal hemodynamic instability during induction of general anesthesia and provides cardiovascular stability in high-risk patients^{4, 5}. Remifentanyl has been successfully used in cases of maternal cardiac diseases⁶⁻⁹ intracranial cysts¹⁰, and preeclampsia¹¹.

Although previous studies reported newborn chest wall muscle rigidity for short periods¹², as well as neonatal depression^{13,14} following remifentanyl administration, other studies suggest that low Apgar and neonatal depression may be dose-dependent and the use of a low dose of remifentanyl may be safe for newborns^{15,16}.

In this study we investigated the maternal and neonatal effects of remifentanyl as compared with the administration of a saline solution during the induction of general anesthesia for cesarean delivery.

Material and Methods

A double blind, randomized, placebo-controlled clinical trial was conducted from October 2009 to November 2010 at the Shahid Sadooghi Hospital, Yazd, Iran. After a review by the university institutional ethics committee, the study was registered in the Iranian registry of clinical trials (<http://irct.ir>) as IRCT201102252963N2. Forty patients were chosen for each group on the basis of similar studies with a power of 85% and α confidence level of 0.05. A minimal difference to be detected in heart rates was set at 10 beats.

Women with singleton, term pregnancy with a physical status of I or II as defined by the American Society of Anesthesia (ASA) were included in the study. Patients were selected from a pool of approximately 1000 C/S candidates. All patients were candidates for elective cesarean section by general anesthesia because of contraindications for regional anesthesia. The indicators for induction of general anesthesia included: refusal of spinal anesthesia by patients, anticoagulation therapy, and previous spinal surgery.

Patients with preeclampsia, cardiovascular and neurologic diseases, hypertension, hepatic insufficiency, renal failure, thyroid dysfunction, drug or alcohol abuse history, known fetal abnormalities, and predicted difficulty with airway management were all excluded from the study. The patients were randomly divided into two groups: an experimental group with remifentanyl administration (n= 40) and a control group with normal saline injection (n= 40). (Fig 1) Random selection to each group was performed by drawing a piece of paper from a bag containing equal numbers for each method option. The randomization process was conducted by a biostatistician who was blinded to patient groups during the randomization and analysis processes. A written informed consent was obtained from patients before starting the procedure.




Figure 1: Flow of the study

Anesthetic management:

For premedication, 10 mg metoclopramide was given one hour before induction of the anesthesia. In the operating room, two IV lines were inserted and 1000 mL lactating ringer solution infused before the induction of anesthesia.

Routine monitoring included non-invasive measurements of blood pressure, electrocardiography, pulse oximetry and capnography. Systolic blood pressure (SBP), diastolic blood pressure (DBP), mean arterial pressure (MAP) and heart rate (HR) were recorded before induction of anesthesia, immediately after laryngoscopy and tracheal intubation, immediately after skin incision and after delivery by using a BCI Advisor Vital Signs Monitor. Hemodynamic variables were measured by anesthetic technicians who were unaware of patient groups.

The patients were pre-oxygenated for three minutes before the induction of anesthesia while in a supine position. General anesthesia was induced by using a rapid sequence administration of sodium thiopental 5 mg/kg and succinylcholine 1.5 mg/kg. Following direct laryngoscopy and endotracheal intubation within 15 seconds, anesthesia was maintained with 0.7% isoflurane in a 50% nitrous oxide with 50% oxygen mixture. Isoflurane concentration was decreased to 0.5% after delivery and a dose of 0.5 mg/kg atracurium was administered as a muscle relaxant.

In the experimental group, a 0.5 µg/kg remifentanyl bolus was injected over 30 seconds immediately before induction of anesthesia. For the control group; the same volume of saline was infused as a placebo.

After delivery and clamping the umbilical cord, 1.5 µg/kg fentanyl was given to patients of both groups. Following extubation, post-operative analgesia was managed using a patient-controlled morphine injection pump.

After delivery, neonates were evaluated by a pediatrician who was blinded to the mother's group. The Apgar scores at the first and fifth minutes were also determined. Additionally, the umbilical artery pH of the newborns was assessed by the use of the Gem 3000 Blood Gas Analyzer.

All registered data were digitalized for the SPSS-(r)15 software and were analyzed by T-test. P-value < 0.05 was considered as the level for statistical significance.

Results

Demographic, maternal, and fetal characteristics of the two groups are indicated and compared in Table 1. There was no significant difference found between the two groups studied, according to the demographic characteristics of participants (Table 1). Maternal age, maternal weight, the duration of pregnancy, as well as the neonatal weight were not statistically different between the groups studied. Elective (non-emergent) cesarean section was performed for all women. The duration between uterine incision and delivery was found to be 51.53 ± 9.50 and 49.34 ± 9.20 in remifentanyl and control groups, respectively ($P > 0.05$).

No differences were found between the groups regarding systolic and diastolic blood pressure before induction of anesthesia and

after delivery. Systolic and diastolic blood pressures were reduced significantly after tracheal intubation and skin incision in the remifentanyl group when compared with controls ($P < 0.05$) (Table 2). There was no significant difference between the two groups regarding heart rate at any time. All neonates had Apgar scores above seven after the first minute. Assisted ventilation, tracheal intubation and naloxane administration were not necessary.

The mean first-minute Apgar scores were 8.83 ± 0.79 and 8.87 ± 0.40 for the remifentanyl and control groups, respectively, with no statistical differences ($P > 0.05$). Mean Apgar scores at the fifth minute were 9.90 ± 0.48 and 9.97 ± 0.16 for the experimental and control groups, respectively, with no significant differences ($P > 0.05$). Umbilical artery blood PH was measured as 7.37 ± 0.07 for the remifentanyl group and 7.35 ± 0.02 for the control group. Differences were not found to be significant ($p > 0.05$).

Discussion

In cesarean section under general anesthesia, it is ideal to minimize maternal surgical stress and newborn depression. Based on the present study, significant reduction in systolic and diastolic blood pressures after laryngoscopy, intubation and skin incision was found among women in the remifentanyl group when compared with the control group. The results indicated that a single bolus of 0.5 µg/kg remifentanyl may prevent the hypertensive responses following laryngoscopy, tracheal intubation and skin incision.

The positive effect of remifentanyl on hemodynamic stability was observed in previous studies using different doses of remifentanyl. Van de Velde et al administered 0.5 µg/kg bolus of remifentanyl followed by continuous infusion of 0.2 µg/kg/min combined with propofol. A constant mean arterial pressure during anesthesia was reported¹⁵. Similarly, Ngan Kee et al used a bolus of 1 µg/kg remifentanyl before induction of anesthesia and found that patients in the placebo group had higher SBP and MAP when compared with the remifentanyl group³. In contrast, Draisci et al who used 0.5 µg/kg bolus of remifentanyl followed by a continuous infusion of 0.15 µg/kg/min did not find any significant differences in blood pressure between the remifentanyl and control groups¹³.

No significant differences between the two groups regarding SBP and DBP after delivery were found in our study. This may be explained by the prompt action and rapid metabolism of remifentanyl. The mean induction-to-delivery interval found in this study was 7.5 min. It is assumed that remifentanyl with its short half life¹⁷ did not decrease the blood pressures after delivery in the experimental group. Similarly, in the Draisci et al study¹³ no significant differences were found regarding the blood pressure at the end of surgery between the groups studied. However, Van de Velde et al found a significant difference in mean arterial pressure at the time of delivery between the groups studied¹⁵.

Our study was similar to that of the Draisci et al survey in that no significant difference was found in the heart rates between the studied groups¹³. In contrast, Ngan Kee et al³ reported that the heart rate for the control group was higher than that found for the remifentanyl group. This difference may be due to the lower dose of remifentanyl used in our study when compared with the dose used in the Ngan Kee et al study. Management of the increased

	Remifentanyl group* (n=40)	Control group* (n=40)	P value
Age (years)	27.78±4.57	27.46±4.54	0.750
Weight (kg)	80.50±13.61	76.00±12.32	0.125
Gestational age (weeks)	39.21±1.47	39.07±1.59	0.680
Birth weight (kg)	3.16±0.33	3.19±0.37	0.736
Uterine incision - delivery time (sec)	51.53±9.50	49.34±9.20	0.303

Table 1. Maternal and neonatal characteristics in both groups.

* Values are presented as the mean±SD.

heart rate following laryngoscopy and intubation seems to require higher doses of anesthetic and analgesic drugs¹⁸.

In the current study, no differences were found between groups according to the mean Apgar scores at the first and fifth minutes. All neonates had Apgar scores higher than seven at the first minute and no neonate required naloxan administration or tracheal intubation. Given the remifentanyl dose used in our study, it appears safe for use with newborns. However, Ngan Kee claimed that even low doses of remifentanyl can cause neonatal respiratory depression. Despite giving naloxan to 10% of the neonates in the remifentanyl group, our study found that no differences were present between groups regarding the Apgar score³. There are, however, other reports concerning naloxan administration¹⁰ where apnea and generalized chest wall rigidity^{12,19} in neonates was found when remifentanyl was given during cesarean section.

Kan et al reported a high transplacental passage of remifentanyl, but only transient neonatal depression². Van de Velde's study reported that approximately 50% of infants showed a brief respiratory depression with the need for naloxan administration or assisted ventilation and that two of 13 neonates studied had Apgar scores less than seven at the fifth-minute evaluation¹⁵. This disparity may be due to the use of remifentanyl in combination with propofol as was done in the Van de Velde study. Draisci *et al* found a significant reduction in Apgar scores at the first and fifth minutes in the remifentanyl group; however, all Apgar scores were greater than eight at the fifth minute without requiring naloxan. Further, Draisci indicated that a high bolus dose and a lower infusion of remifentanyl may be useful for surgical stress management with minimal neonatal sedation¹³.

According to our study, by avoiding the use of remifentanyl infusion, neonatal depression may be decreased to minimum level because of the rapid metabolism of the single dose of remifentanyl. However, in the Van de Velde study, a single bolus dose of remifentanyl caused respiratory depression in two neonates that required naloxan administration¹⁵. It should be noticed that the single bolus dose of remifentanyl which was used in Van de Velde's study was two times greater than the single bolus dose we used. By using a single bolus of 0.5 µg/kg remifentanyl in our study, no neonatal depression occurred and all neonates had Apgar scores greater than seven without naloxan being required.

Mean umbilical artery pH was greater than 7.35 in both groups of neonates. Since low umbilical artery pH reflects both respiratory and metabolic acidosis, it can be concluded that remifentanyl administration in the dose given in our study will possibly not cause respiratory and metabolic acidosis. Similar results were reported in other studies^{13,15}. However, some neonates in these studies experienced fetal distress following remifentanyl administration to their mothers. In a cohort study, only 19% of the infants with low Apgar scores had severe acidosis at birth and 27% of the infants with acidosis had a low Apgar score at the fifth minute²⁰.

Conclusion:

According to our observations, the administration of a single bolus dose of remifentanyl in women undergoing caesarean under general anesthesia, may possibly reduce systolic and diastolic blood pressures, but not the heart rate. Remifentanyl with its rapid onset of action and rapid metabolism can be recommended as a safe opioid with minimal newborn depression, especially when it is administered in a single bolus dose. It seems that further stud-

	Remifentanyl group* (n=40)	Control group* (n=40)	P value
Before induction			
Systolic blood pressure	129.80±11.54	125.81±10.22	0.066
Diastolic blood pressure	81.07±11.22	79.71±11.66	0.596
After intubation			
Systolic blood pressure	129.64±16.52	137.97±19.59	0.041
Diastolic blood pressure	81.97±16.70	90.12±19.32	0.045
Skin incision			
Systolic blood pressure	124.65±15.88	137.23±16.73	0.001
Diastolic blood pressure	79.75±13.56	88.00±15.91	0.015
After delivery			
Systolic blood pressure	129.97±12.33	127.31±15.61	0.915
Diastolic blood pressure	74.41±13.24	75.42±13.77	0.744

Table 2. Blood pressure changes between induction and delivery in both groups

*Values are presented as the mean±SD.

ies with larger sample sizes and different remifentanyl doses are required to evaluate the risk/benefit ratio for both mothers and neonates.

Conflict of interest:

This study was financially supported by the Faculty of Medicine, Shahid Sadoughi University of Medical Sciences, Yazd, Iran as part of Dr. Hamid Harrazi's dissertation in Anesthesiology. The authors declare that no conflict of interest was present.

Acknowledgment

The authors wish to thank the Shahid Sadoughi Hospital's operating room staff for their skillful technical assistance during the course of this study.

References

- 1- Jenkins JG, Khan MM. Anaesthesia for Caesarean section: a survey in a UK region from 1992 to 2002. *Anaesthesia* 2003; 58 (11): 1114-8.
- 2- Kan RE, Hughes SC, Rosen MA, Kessin C, Preston PG, Lobo EP. Intravenous remifentanyl: placental transfer, maternal and neonatal effects. *Anesthesiol.* 1998; 88(6):1467-74.
- 3- Ngan Kee WD, Khaw KS, Ma KC, Wong AS, Lee BB, Ng FF. Maternal and neonatal effects of remifentanyl at induction of general anesthesia for cesarean delivery: a randomized, double-blind, controlled trial. *Anesthesiol* 2006;104(1):14-20.
- 4- Prakash N, McLeod T, Gao Smith F. The effects of remifentanyl on haemodynamic stability during rigid bronchoscopy. *Anaesthesia* 2001;56 (6): 576-80.
- 5- Orme RM, Grange CS, Ainsworth QP, Grebenik CR. General anaesthesia using remifentanyl for caesarean section in parturients with critical aortic stenosis: a series of four cases. *Int J Obstet Anesth* 2004;13(3):183-7.
- 6- Macfarlane AJ, Moise S, Smith D. Caesarean section using total intravenous anaesthesia in a patient with Ebstein's anomaly complicated by supraventricular tachycardia. *Int J Obstet Anesth* 2007;16(2): 155-9.
- 7- Amini S, Yaghmaei M. The use of remifentanyl in general anesthesia for cesarean section in a parturient with severe mitral stenosis and pulmonary edema. *Middle East J Anesthesiol* 2010;20(4): 585-8.
- 8- Bilehjani E, Kianfar AA, Toofan M, Fakhari S. Anesthesia with etomidate and remifentanyl for cesarean section in a patient with severe peripartum cardiomyopathy--a case report. *Middle East J Anesthesiol* 2008; 19 (5):1141-9.
- 9- Miyawaki Y, Yumiba T, Asaga T, Taie S, Shirakami G. Remifentanyl is useful for cardiovascular stability during cesarean delivery in a parturient with Marfan's syndrome. *Masui* 2010;59(10):1305-7.
- 10- Imarengiaye C, Littleford J, Davies S, Thapar K, Kingdom J. Goal oriented general anesthesia for Cesarean section in a parturient with a large intracranial epidermoid cyst. *Can J Anaesth* 2001;48(9): 884-9.
- 11- Richa F, Yazigi A, Nasser E, Dagher C, Antakly MC. General anesthesia with remifentanyl for Cesarean section in a patient with HELLP syndrome. *Acta Anaesthesiol Scand* 2005;49(5): 418-20.
- 12- Carvalho B, Mirikitani EJ, Lyell D, Evans DA, Druzin M, Riley ET. Neonatal chest wall rigidity following the use of remifentanyl for cesarean delivery in a patient with autoimmune hepatitis and thrombocytopenia. *Int J Obstet Anesth* 2004;13(1): 53-6.
- 13- Draisci G, Valente A, Suppa E, Frassanito L, Pinto R, Meo F, et al. Remifentanyl for cesarean section under general anesthesia: effects on maternal stress hormone secretion and neonatal well-being: a randomized trial. *Int J Obstet Anesth* 2008;17(2):130-6.
- 14- Yoo KY, Jeong CW, Park BY, Kim SJ, Jeong ST, Shin MH, et al. Effects of remifentanyl on cardiovascular and bispectral index responses to endotracheal intubation in severe pre-eclamptic patients undergoing Caesarean delivery under general anaesthesia. *Br J Anaesth* 2009;102(6):812-9.
- 15- Van de Velde M, Teunkens A, Kuypers M, Dewinter T, Vandermeersch E. General anaesthesia with target controlled infusion of propofol for planned caesarean section: maternal and neonatal effects of a remifentanyl-based technique. *Int J Obstet Anesth* 2004;13(3):153-8.
- 16- Volmanen P, Akural EI, Raudaskoski T, Alahuhta S. Remifentanyl in obstetric analgesia: a dose-finding study. *Anesth Analg* 2002; 94(4): 913-7.
- 17- Glass PSA, Hardman D, kamiyama Y, Quil TJ, Marton G, Donn KH, et al. Preliminary pharmacokinetics and pharmacodynamics of an ultra-short-acting opioid: Remifentanyl. *Anesth Analg* 1993;77(5): 1031-40.
- 18- Minto CF, Schnider TW, Short TG, Gregg KM, Gentilini A, Shafer SL. Response surface model for anesthetic drug interactions. *Anesthesiol.* 2000; 92 (6): 1603-16.
- 19- McCarroll CP, Paxton LD, Elliott P, Wilson DB. Use of remifentanyl in a patient with peripartum cardiomyopathy requiring Caesarean section. *Br J Anaesth* 2001;86(1):135-8.
- 20-Sykes GS, Johnson P, Ashworth F, Molloy PM, Gu W, Stirrat GM, et al. Do Apgar scores indicate asphyxia? *Lancet* 1982; 1(8270): 494-6.