



Original Article

Elders Health Empowerment Scale. Spanish adaptation and psychometric analysis

Escala de Empoderamiento sobre la Salud para Adultos Mayores. Adaptación al español y análisis psicométrico

Daniel Jorge Luis Serrani Azcurra

Professor at Department of Biological Sciences, Faculty of Psychology, National University of Rosario, Rosario, Argentina. Maimónides University, Buenos Aires, Argentina.

Serrani Azcurra DJL. Elders Health Empowerment Scale. Spanish adaptation and psychometric analysis. *Colomb Med.* 2014; 45(4): 179-85.

© 2014 Universidad del Valle. This is an Open Access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Article history

Received: 11 February 2014
Revised: 20 September 2014
Accepted: 20 November 2014

Keywords

Patient empowerment, factor construct rating scale, chronic diseases, elderly, health care utilization

Palabras clave:

Empoderamiento del paciente, Escala de calificación de el factor del constructo, enfermedades crónicas, anciano, uso del servicio de salud.

Abstract

Introduction: Empowerment refers to patient skills that allow them to become primary decision-makers in control of daily self-management of health problems. As important the concept as it is, particularly for elders with chronic diseases, few available instruments have been validated for use with Spanish speaking people.

Objective: Translate and adapt the Health Empowerment Scale (HES) for a Spanish-speaking older adults sample and perform its psychometric validation.

Methods: The HES was adapted based on the Diabetes Empowerment Scale-Short Form. Where “diabetes” was mentioned in the original tool, it was replaced with “health” terms to cover all kinds of conditions that could affect health empowerment. Statistical and Psychometric Analyses were conducted on 648 urban-dwelling seniors.

Results: The HES had an acceptable internal consistency with a Cronbach’s α of 0.89. The convergent validity was supported by significant Pearson’s Coefficient correlations between the HES total and item scores and the General Self Efficacy Scale ($r=0.77$), Swedish Rheumatic Disease Empowerment Scale ($r=0.69$) and Making Decisions Empowerment Scale ($r=0.70$). Construct validity was evaluated using item analysis, half-split test and corrected item to total correlation coefficients; with good internal consistency ($\alpha > 0.8$). The content validity was supported by Scale and Item Content Validity Index of 0.98 and 1.0, respectively.

Conclusions: HES had acceptable face validity and reliability coefficients; which added to its ease administration and users’ unbiased comprehension, could set it as a suitable tool in evaluating elder’s outpatient empowerment-based medical education programs.

Resumen

Introducción: Empoderamiento se refiere a las habilidades que le permiten al paciente convertirse en responsable de tomar las decisiones para el control diario de sus problemas de salud. A pesar de ser un concepto tan importante, particularmente para adultos mayores con problemas crónicos de salud, hay pocos instrumentos accesibles que hayan sido validados para su uso en hispano-hablantes.

Objetivo: Traducir y adaptar la Escala de Empoderamiento sobre la Salud (EES) para una muestra de adultos mayores hispano-hablantes y llevar a cabo su validación psicométrica.

Métodos: La EES se adaptó basándose en la Escala de Empoderamiento de la Diabetes versión corta. Donde se mencionaba “diabetes” en el instrumento original, se reemplazó con el término “salud” para cubrir todos los tipos de condiciones que podrían afectar el Empoderamiento sobre la salud. Se realizaron análisis estadísticos y psicométricos sobre 648 adultos mayores residentes urbanos.

Resultados: La EES tuvo una consistencia interna aceptable con un α de Cronbach de 0.89. la validez convergente se apoyó en un coeficiente de correlación de Pearson significativo entre la EES total y por ítems y la Escala General de Auto-eficacia ($r=0.77$), la Escala de Empoderamiento para la Enfermedad Reumática versión Sueca ($r=0.69$) y la Escala de Empoderamiento Tomando Decisiones ($r=0.70$). La validez de Constructo se evaluó mediante análisis de ítem, test de las dos mitades y coeficiente de correlación ítem corregido total ($\alpha > 0.8$). La validez de contenido se apoyó por los Índices de Validez de Contenido para la Escala y para los ítems de 0.98 y 1.0, respectivamente.

Conclusiones: La EES tuvo una validez y confiabilidad aceptables, que sumados a su facilidad de administración y comprensión simple y sin sesgos podría constituirse en una herramienta confiable para evaluar programas educativos médicos basados en el Empoderamiento de pacientes mayores ambulatorios.

Corresponding author:

Daniel Jorge Luis Serrani Azcurra. Professor at Department of Biological Sciences, Faculty of Psychology, National University of Rosario, Rosario, Argentina. Maimónides University, Buenos Aires, Argentina. e-mail: danielserrani@argentina.com

Introduction

The medical health literature on empowerment has increased exponentially since the early 1990s, particularly in relation to chronic conditions¹. Empowerment is a workable and patient-centered approach leading to effective interventions for addressing the psychosocial components of living with chronic diseases. In order to accomplish those goals, it must be recognized that patients are the primary decision-makers in control of the daily self-management of their health problems, and emphasizing patient autonomy and expansion of freedom of choice is mandatory². The concept of empowerment is deeply rooted in social sciences³ and has been defined as the complex and multifaceted process of recognizing self needs, skills and resources, improve owns abilities to solve problems, reach a sense of power or control and enable people gain mastery over their lives⁴. This concept is being included as part of the usual health practices in modern medicine occurring within the context of a nurturing physician-client relationship⁵. It is expected that through this process of empowerment, client's perceptions of competence regarding the ability to maintain good health and manage interactions with the health care system would improve, as a result of the internalization of current health ideas and goals at the individual and societal level. In the case of older adults, empowerment should promote well-being, healthy lifestyles and social connectedness⁶. According to those trends, interventions have been actively developed based on empowerment theory⁷ and various instruments have been in

use to acknowledge different chronic medical conditions ranging from rheumatic diseases⁸ to cardiovascular illness or diabetes⁹. The Diabetes Empowerment Scale-short form (DES-SF)¹⁰ measures empowerment in patients with diabetes, has been tested with elder patients and has been adapted to evaluate health-related empowerment. However, a Spanish translation or adaptation is lacking, so the goals of this study were to translate and adapt the DES-SF as a Health Empowerment Scale for Spanish-speaking elders, and assess its psychometric properties.

Materials and Methods

Instruments

The Health Empowerment Scale (HES) was adapted from The Diabetes Empowerment Scale Short-form (DES-SF), which was selected for its brevity, reducing the chance of non-response due to poor concentration, and its excellent validity and reliability criteria, while reflecting the attributes of empowerment. The mean age of subjects in the original data set and follow-up studies was 60 years, which further supports its use for elderly subjects. After substituting the word "diabetes" with "health" in each item from the DES-SF to assess health-related empowerment, the instrument retained 8 items, scored on a 5 points Likert scale ranging from 5 (strongly agree) to 1 (strongly disagree). Higher scores indicate stronger level of health-related empowerment (Table 1).

Table 1. Relationship between Health Empowerment Scale and properties of empowerment.

Subscales of HES	Questions of HES (spanish/english)	
Satisfaction and dissatisfaction related to health	Sé muy bien con que parte(s) del cuidado de mi salud no estoy satisfecho	Self-Control
	I know what part(s) of taking care of my health that I am dissatisfied with.	
Identification and achievement of personally meaningful goals	Soy capaz de alcanzar mis metas de salud mediante planes concretos de acción	Self-efficacy
	I am able to turn my health goals into a workable plan.	
Application of a systematic problem-solving process	Tengo diferentes maneras de superar los obstáculos para lograr mis objetivos de salud	Problem solving
	I can try out different ways of overcoming barriers to my health goals	
Coping with the emotional aspects of living with health	Tener salud me hace sentir mejor	Psychosocial coping
	I can find ways to feel better about having health.	
Stress management	Puedo afrontar el estrés por mis problemas de salud de manera positiva	Psychosocial coping
	I Know the positive ways I cope with health-related stress.	
Appropriate social support	Puedo solicitar ayuda para cuidar y mantener mi salud cuando lo necesito	Support
	I can ask for support for having and caring for my health when I need it.	
Self-motivation	Reconozco lo que me motiva para cuidar mi salud	Motivation
	I know what helps me stay motivated to care for my health.	
Making cost/benefit decisions about making behavior changes	Me conozco lo suficiente para escoger lo que más conviene a mi salud	Decision making
	I know enough about myself as a person to make health care choices that are right for me.	

Other related instruments used for concurrent validity were:

- Making Decisions Empowerment Scale¹¹ is a 28-item self report questionnaire designed to measure empowerment in patients with psychological disorders; with five subscales (self-efficacy, power, community activism, righteous anger and optimism over the future) responded to on a four-point scale, exhibiting good internal consistency (Cronbach's alpha= 0.81).
- Swedish Rheumatic Disease Empowerment Scale (SWE-RES-23)⁸ adapted for use in patients with rheumatic diseases, has 23 items responded on a 4-points Likert scale, containing five factors (empowerment subscales), and a Cronbach's $\alpha = 0.59$ to 0.92.
- The General Self Efficacy Scale (Spanish Adaptation)¹² assess people's believe about their ability to appropriately manage life stressors. It has 10 items rated with a 10 point Likert scale and has high internal consistency (between 0.87) and predictive value with other correlated scales.

Procedures

To assess the validity and reliability of the Health Empowerment Scale (HES), the DES-SF was translated into Spanish by a certified translator and two physicians specialized in gerontology. Each time the word "diabetes" appeared, it was replaced by the word "health" to accomplish the general purpose of the instrument. Another certified translator, blind to the first translation, translated the HES back into English. After that, a content analysis was performed by two gerontology professors and one licensed nurse with training in gerontology. They were asked to make comments on individual questions in relation to the accuracy, clarity, cultural relevance semantic, conceptual and operational equivalence of the translation¹³. Questions number 4 and 8 were evaluated as difficult to understand directly (regarding fluency) and were rephrased, and question 6 mixed the terms "health care" and "health". All questions were edited in light of the comments. A pilot study (n= 32) was performed with the panel-modified version at a community senior center on August 2013. The researcher read the scale in a Spanish consistent manner and recorded the responses. The HES required an average of 18 min to complete. The panel-modified version was readily accepted. Furthermore, to assess its readability properties the scale was processed with the INFLESZ 1.0 software which issued a legibility index of 69.87 (acceptable when ≥ 55) meaning that the scale was considered simple to understand according to an elementary literacy level¹⁴. Therefore, in September 2013, the main study was conducted for 700 urban-dwellers senior citizens assisting to community day-centers located in Rosario (Argentina). The data were collected by three nursing majors who had been trained to perform the survey in a similar and constant manner. They administered the questionnaire one-on-one. Of a total of 700 copies distributed for the survey, 648 were completed and returned.

Ethical considerations

The study was carried out after receiving approval from the institutional review board at National University of Rosario. Written informed consents were obtained when the participants agreed to participate in the study.

Sample

Subjects were randomly selected from a large population of senior citizens regularly assisting to community day-centers located in Rosario. A sample size of 103 subjects was required in order to get an effect size of 0.3, a significance level of 0.01 for a type I error and a power of 0.8 at correlation analysis¹⁵. Below is the formula used to calculate the sample size, according to Altman's¹⁶.

However, taking into account number of participants to figure out sample size in factor analysis, some authors¹⁷ have suggested the following guide samples sizes: 50 as very poor; 100 as poor, 200 as fair, 300 as good, 500 as very good and 1,000 as excellent. According to that, we decided to include in the final sample 648 elderly individuals.

Statistical analysis

Descriptive statistics were used to establish the frequency, range, mean, and SD of demographic and clinical characteristics of the main sample. The HES reliability was assessed through internal consistency with total Cronbach's alpha, half-split analysis, Item-to-total correlations and Cronbach's alpha without item. Validity was evaluated through Principal and Confirmatory Factor Analysis (maximum likelihood). The SPSS[®] statistical package for the Social Sciences (SPSS[®]-19) was used to compute descriptive statistics, correlation and internal consistency together with t-test analyses. Principal component analysis allowed performing statistical significance testing of factor loadings, correlations among factors and computation of confidence intervals used for factor extraction. After extraction, factors were retained for rotation according to eigenvalues greater than 1.0, items loadings above 0.30, no or few item cross-loadings and no factors with fewer than three items. Varimax was the selected orthogonal rotation method as it assumes no correlation exists between factors, rendering a more accurate solution. Factorability of the correlation matrix was based on Bartlett's test of sphericity to estimate the probability that correlations in a matrix were 0, and the Kaiser-Meyer-Olkin (KMO) measure of sampling adequacy which accounts for the relationship of partial correlations to the sum of squared correlations, thus indicating the extent to which a correlation matrix actually contains factors or simply chance correlations between a small subset of variables. Values ≥ 0.60 are required for good factor analysis. Confirmatory factor analysis (CFA) was carried out using the maximum likelihood (ML) method. Since departures from multivariate normality can have a significant impact on maximum-likelihood estimation, we calculated descriptive analytical measures prior to conducting CFA analysis. As kurtosis statistics were found to indicate normality, no other correction was used to adjust the model chi-square¹⁸. According to Schweizer's recommendations¹⁹ additional measures of model fit were used: root mean square error of approximation (RMSEA) with values below 0.08 considered acceptable fit; comparative fit index (CFI) ≤ 0.06 as good fit values; GFI and NNFI with values ≥ 0.90 as acceptable fit. Three putative models were compared with one, two and three factors. The last (three factor) was a less parsimonious model, in which the factor inter-correlation was freely estimated; while the first model (one factor) was a more parsimonious model, the independence model, in which the factor inter-correlation was set to 0 to represent the factor structure of the original scale. Concurrent validity was tested with Pearson correlation between HES and GSES, SRDES and MDES. Content

Validity was evaluated with Scale and Item Construct Validity Index. Statistical significance was established at $p= 0.01$ (two-tails).

Results

Participants' socio-demographic data are displayed in Table 2. Most of them were married females, with a mean age of 75.5 for men and 74.1 for women, ranging from 64 to 93 years; most of them female (71.0%), and most (89.7%) had one or more chronic diseases, two-thirds with hypertension, nearly half of them with arthritis and one-third with diabetes. On average they had 2.21 chronic diseases / person and took an average of 2.86 types of different drugs. Most participants achieved incomplete literacy level, had regular medical consultation, but rarely had been admitted to hospital. They had medium incomes level and exhibited mild self-efficacy scores. A summary of the baseline characteristics of participants is offered in Table 2.

Reliability

The mean HES score for the sample was 3.5 (SD 0.73), every question was in the range between 3.22-3.79 (SD 1.03-1.22). Coefficient of kurtosis (-0.764 to 0.077) and skewness (1.843-3.058) showed a normal distribution. HES exhibited excellent internal consistency with Cronbach's $\alpha= 0.89$ and >0.81 for full scale and corrected items to total correlation respectively. Floor and ceiling effects were small ($<20\%$), suggesting that HES has significant power to measure health empowerment level of older adults (Table 3). Test-retest reliability was assessed in 23 participants evaluated by the same researcher with an interval of 3 months, and intra-class correlation coefficient (ICC) was 0.92 ($p= 0.001$) suggesting good stability along time. With the half-splitting analysis of the scale, first and second halves showed a good reliability coefficient of 0.86 and 0.91 respectively.

Validity

Construct validity was demonstrated by Kaiser-Meyer-Olkin sampling (KMO = 0.890) and Bartlett's test of sphericity ($\chi^2_{(634)} = 5425.72$; $p < 0.001$) which showed that sample size was suitable for conducting factor analysis (least factor load method) and the correlation matrix had not occurred by chance. A single factor solution was judged the more acceptable, explaining 52.4% of the variance. Confirmatory Factor Analysis on three putative models (with one, two and three factors) showed the best fitness index for the one factor scale with CFI, GFI and NNFI ≥ 0.90 , and RMSEA ≤ 0.06 as good fit values (Table 4)

Concurrent validity

Significant Pearson's correlations were found between Total and Items' HES scores and the other three scales (GSES, SRDES and MDDES) (Table 5).

Content validity

After the survey, three gerontologists, two licensed nurses with gerontology training, and the author rated each question as a valid measure of the construct using a five-point Likert scale (1= strongly disagree, 5= strongly agree). Based on their responses, an item (I-CVI) and scale (S-CVI) content validity index scores of 0.98 and 1.0 respectively, were accepted as high.

Table 2. Demographic characteristics of subject (N=648)

	Male	Female	%	F (p)
	(n= 188) (29%)	(n=460) (71%)		
Age (years)				
60-69	28	68	15.0	
70-79	144	353	77.0	
80-89	12	31	6.6	
≥ 90	2	6	1.4	
Mean (SD)	75.5 (11.4)	74.1 (11.9)		0.34 (<0.01)
Schooling (years)				
None	31	74	16.0	
Elementary school	80	194	42.0	
Middle school	26	65	14.0	
High school	33	84	18.0	
College	16	38	8.0	
Graduate School	2	5	1.0	
Mean (SD)	9.5 (2.4)	7.1 (1.9)		0.04 (<0.01)
Marital Status				
Single	39	96	21.0	
Married	130	318	69.0	
Other	19	46	10.0	
Household Income/ month (thousand won)				
1000-1999	71	175	38.0	
2000-2999	86	211	46.0	
3000-3999	22	55	12.0	
4000-5000	9	23	5.0	
Number of chronic diseases				
Hypertension	111	271	58.9	
Arthritis	75	185	40.3	
Diabetes Mellitus	39	95	20.7	
Hyperlipidemia	62	79	17.1	
Osteoporosis	11	28	0.2	
Mean (SD)	3.5 (1.4)	2.1 (0.9)		0.14 (<0.01)
Number of visits to medical facilities (per one month)				
Mean (SD)	2.0	3.5		0.08 (<0.01)
Admission to medical facility in the past 5 years				
Yes	67	165	35.9	
No	121	295	64.1	
Mean (SD)	0.87 (0.11)	1.17 (0.54)		0.37 (<0.01)
General Self Efficacy Scale				
Mean (SD)	79.3 (21.3)	75.3 (20.5)		0.36 (<0.01)
Making Decisions Empowerment Scale				
Mean (SD)	97.2 (24.6)	96.1 (22.1)		0.69 (<0.01)
Swedish Rheumatic Disease Empowerment Scale				
Mean (SD)	65.2 (11.3)	62.5 (16.3)		0.31 (<0.01)
Participants (N= 648)				

Table 3. Statistics, corrected item-total correlation, floor-ceiling effects for Health Empowerment Scale (n= 648).

Item content	Mean	SD	Skewness	Kurtosis	Corrected Item-Total Correlation	α if Item deleted	Floor effect (% scoring 1)	Ceiling effect (% scoring 5)
I know what part(s) of taking care of my health that I am dissatisfied with.	3.22	1.12	0.071	-1.094	0.617	0.711	2.8	13.8
I am able to turn my health goals into a workable plan.	3.33	1.22	-0.213	-0.988	0.607	0.799	6.7	21.4
I can try out different ways of overcoming barriers to my health goals	3.22	1.10	-0.077	-0.903	0.663	0.798	4.5	11.7
I can find ways to feel better about having health.	3.75	1.03	-0.578	-0.254	0.614	0.749	2.2	24.6
I Know the positive ways I cope with health-related stress.	3.63	1.09	-0.527	-0.563	0.632	0.774	2.4	21.5
I can ask for support for having and caring for my health when I need it.	3.49	1.30	-0.78	-1.157	0.586	0.629	6.7	27.7
I know what helps me stay motivated to care for my health.	3.62	1.07	-0.595	-0.247	0.795	0.831	4.1	20.1
I know enough about myself as a person to make health care choices that are right for me.	3.79	1.07	-0.764	0.058	0.628	0.769	4.1	28.1

Discussion

In conclusion, the HES showed excellent reliability and validity for assessing the health-related empowerment of elderly individuals. First, the mean of the HES was 3.51, which shows that senior citizens have an empowerment level above medium on a score of 3, which represents “neither agree nor disagree” on the Likert scale. The same results were found in other studies using similar instruments such as the Empowerment for Self Care Scale²⁰ (Mean= 3.65, SD= 0.40), or the Patient Empowerment Scale (Mean= 3.68, SD= 0.53)²¹. Both the floor and ceiling effects were small according to usual criteria²². The ceiling effect refers to patients who start with higher empowerment abilities than the average patient and lack room for improvement, while floor effect means the opposite as these subjects have lower empowerment skills than average and so may show greater (and biased) improvements. As a consequence, HES was balanced enough to correctly measure health empowerment interventions outcomes without skewing the end results. HES Cronbach's α was good (0.89) compared to DES-SF reliability (0.84). Corrected item-total correlation was high for all questions (>0.81) although the lower value was for question 6. A possible explanation could be that participants regarded supporters to be acquaintances such as family members or friends, and many elders reported that they should not ask their family for help because they did not want to be a burden for their family. Instead they build on the sense that empowerment should render them independent from their family. So item 6 content should add health care providers or health care system to family and friends as putative support providers. Content validity of the HES was acceptable (I-CVIs= 1.0, S-CVI= 0.98). Construct validity was

supported by significant Pearson correlations between the HES total scores and the SRDES ($r= 0.69$), GSES ($r= 0.77$) and MDES ($r= 0.70$). This result shows that there is a strong correlation between empowerment and self-efficacy, supported by previous researches⁶ confirming that self-efficacy is both a component and a result of empowerment²³. It has been argued that health profile of the future elderly population in Latin America will be less predictable due to factors associated with demographic past that haunt them for a long time and make them more vulnerable, even if economic and institutional conditions turn out to be better than what they are likely to be²⁴. The number of chronic conditions will probably increase with age and would be higher among females than among males; levels of self-reported diabetes and obesity will be higher than those found in the US; along with more deteriorated health and functional status in the region. According to these assertions, it seems a valuable strategy to strengthen elder citizens' health promotion activities in our region as a way to reduce inequalities in health care access and improve chronic diseases outcomes such as diabetes or obesity²⁵. Although exhibiting its usefulness in measuring empowerment, several questions must be further developed and tested. Some limitations that can be endowed to this research are that this instrument measures empowerment skills at an individual level, excluding the organizational and community ones, which may be important in the case of the frail or disabled elder. Another limitation that should be considered is that, despite the fact that western cultures share common tenets, sometimes subtle differences between them may account for some issues like sentence comprehension or resource availability which could make the questionnaire not

Table 4. Psychometric results from exploratory and confirmatory factor analysis.

Factor	Eigenvalue	Commonalities	Rank	Initial α	% variance	% variance accumulated	Items Number
1	9.91	1.00	8.11-9.98	0.885	62.4	62.4	6
2	0.87	0.54	0.73-0.90	0.886	3.2	65.6	7
3	0.45	0.43	0.34-0.51	0.887	2.1	67.7	8
Model	Df	Chi2	GFI	CFI	NNFI	RMSEA	Items Number
1 factor	904	156.2	0.991*	0.963*	0.925	0.054*	8
2 factor	904	399.9	0.643	0.793	0.633	0.79	8
3 factor	904	234.9	0.661	0.654	0.701	0.78	8

* $p < 0.001$

Table 5. Coefficient correlations between the 8 items of the HES and the other scales

HES	GSES	<i>p</i>	SRDES	<i>p</i>	MDES	<i>p</i>
Item 1	0.64	0.001	0.67	0.01	0.71	0.05
Item 2	0.51	0.02	0.63	0.43	0.72	0.12
Item 3	0.76	0.05	0.64	0.04	0.88	0.001
Item 4	0.62	0.08	0.76	0.07	0.59	0.03
Item 5	0.75	0.05	0.59	0.06	0.75	0.06
Item 6	0.54	0.09	0.77	0.06	0.69	0.08
Item 7	0.71	0.02	0.78	0.05	0.87	0.003
Item 8	0.84	0.006	0.65	0.04	0.69	0.006
Total Score	0.77	0.005	0.69	0.06	0.70	0.003

General Self efficacy Scales (GSES), Swedish Rheumatic Disease Empowerment Scale (SRDES), and the Making Decisions Empowerment Scale (MDES).

directly suitable in Latin America culture, highlighting the need to take cultural differences between countries into account when adapting questionnaires. It may be suitable to perform a predictive assessment, evaluating participants' health one year later and comparing it with the present health status.

Conclusion

The HES scale possesses acceptable validity and reliability. Considering its brevity and ease of administration, the HES can be used as an outcome measure for the empowerment of Spanish-speaking senior citizens.

Conflict of interest

Not conflict of interest

References

- McAllister M, Dunn G, Payne K, Davies L, Todd C. Patient empowerment: The need to consider it as a measurable patient-reported outcome for chronic conditions. *BMC Health Services Research*. 2012; 12(1): 157.
- Schulman-Green D, Jaser S, Martin F, Alonzo A, Grey M, McCorkle R, *et al*. Processes of self-management in chronic illness. *J Nurs Scholarsh*. 2012; 44(2): 136–44.
- Laverack G. Improving health outcomes through community empowerment: a review of the literature. *J Health Popul Nutr*. 2011; 24(1): 113–20.
- Cattaneo LB, Chapman AR. The process of empowerment: a model for use in research and practice. *Am Psychol*. 2010; 65(7): 646–59.
- Mola E. Patient empowerment, an additional characteristic of the European definitions of general practice/family medicine. *Eur J Gen Pract*. 2013; 19: 128–31.
- Martin P, Tamblyn R, Ahmed S, Tannenbaum C. An educational intervention to reduce the use of potentially inappropriate medications among older adults (EMPOWER study): protocol for a cluster randomized trial. *Trials*. 2013; 14(1): 80.
- Shearer NB, Fleury J, Ward KA, O'Brien AM. Empowerment interventions for older adults. *West J Nurs Res*. 2012; 34(1): 24–51.
- Arvidsson S, Bergman S, Arvidsson B, Fridlund B, Tingström P. Psychometric properties of the Swedish Rheumatic Disease Empowerment Scale, SWE-RES-23. *Musculoskeletal Care*. 2012; 10(2): 101–9.
- Small N, Bower P, Chew-Graham CA, Whalley D, Protheroe J. Patient empowerment in long-term conditions: development and preliminary testing of a new measure. *BMC Health Services Research*. 2013; 13(1): 263.
- Anderson RM, Fitzgerald JT, Gruppen LD, Funnell MM, Oh MS. The diabetes empowerment scale-short form (DES-SF) *Diabetes Care*. 2013; 26(5): 1641–2.
- Hansson L, Björkman T. Empowerment in people with a mental illness: reliability and validity of the Swedish version of an empowerment scale. *Scand J Caring Sci*. 2005; 19(1): 32–8.
- Sanjuán SP, Pérez GAM, Bermúdez MJ. The general self-efficacy scale psychometric data from the Spanish adaptation. *Psicothema*. 2000; 12(Suppl 2): 509–13.
- Sousa VD, Rojjanasrirat W. Translation, adaptation and validation of instruments or scales for use in cross-cultural health care research: a clear and user-friendly guideline. *J Eval Clin Pract*. 2011; 17(2): 268–74.
- Barrio-Cantalejo IM, Simón-Lorda P, Melguizo M, Escalona I, Marijuán MI, Hernando P. Validación de la Escala INFLESZ para evaluar la legibilidad de los textos dirigidos a pacientes. *Anales Sis San Navarra*. 2008; 31(2): 135–52.
- Machin D, Campbell MJ, Tan SB, Tan SH. Sample size tables for clinical studies. Hoboken, NJ: John Wiley & Sons; 2011.
- Altman DG. How large a sample? In: Gore SM, Altman DG (eds). *Statistics in Practice*. London, UK: British Medical Association; 1982.
- Comrey A, Lee HB. *A first course in factor analysis*. 2nd Ed. Hillsdale, NJ: Erlbaum; 1973.
- Hu L, Bentler PM, Kano Y. Can test statistics in covariance structure analysis be trusted? *Psychological Bulletin*. 1992; 112: 351–62.
- Schweizer, K Some guidelines concerning the modeling of traits and abilities in test construction. *Eur J Psychol Assess*. 2010; 26(1): 1–2.
- Dobova SV, Espinosa P, Infante C, Aguirre R, Rodríguez L, Olivares R. Adaptation and validation of scales to measure self-efficacy and empowerment for self-care in Mexican climacteric stage women. *Salud Publica Mex*. 2013; 55(3): 257–66.
- Tu YC, Wang RH, Yeh SH. Relationship between perceived empowerment care and quality of life among elderly residents within nursing homes in Taiwan: A questionnaire survey. *Int J Nurs Stud*. 2006; 43(6): 673–80.
- Mokkink LB, Terwee CB, Patrick DL, Alonso J, Stratford PW, Knol DL, *et al*. The COSMIN checklist for assessing the methodological quality of studies on measurement properties of

health status measurement instruments: an international Delphi study. *Qual Life Res.* 2010; 19(4): 539–49.

23. Andrew N, Peterson NA. Empowerment theory: clarifying the nature of higher-order multidimensional constructs. *Am J Community Psychol.* 2014; 53(1-2): 96–108.

24. Palloni A, McEniry M. Aging and health status of elderly in Latin America and the Caribbean: preliminary findings. *J Cross Cult Gerontol.* 2007; 22(3): 263–85.

25. Cleghorn GD, Lussier J, Velez M, Canovitch M, Licciardello M, Stanlic S. Educating and Empowering Elders: Improving the Health of Senior Latino Diabetics through Community Collaboration. *J Health Disparities Res Practice.* 2008; 2(2): 1–16.