





Adjusting the water retention curve for retractable soils

Luana da Silva, Jackson Adriano Albuquerque, Letícia Sequinatto & Diego Bortolini

Departamento de Solos, Universidade do Estado de Santa Catarina, Lages, Brasil. slv.luana@gmail.com, jackson.irai@gmail.com, letisequinatto@gmail.com, diegobertanbortolini@gmail.com

Received: July 29th, 2020. Received in revised form: January 19th, 2021. Accepted: June 22th, 2021.

Abstract

The soil water retention curve (SWRC) relates moisture to soil water retention energy (matrix potential). The calculation of the volumetric humidity considers the sample volume. In retractable soils, this volume varies according to the drying or wetting of the soil, which can result in errors in the calculated moisture. The objective of this study is to quantify the volume variation in retractable soils and to elaborate the SWRC via the traditional method, which does not consider soil retraction, and a second method, called adjusted, that considers the phenomenon of soil retraction. Soil samples have been collected in horizons A and B from six soil profiles Thus, for retractable soils, it is recommended that the adjustment of the SWRC be carried out considering the actual volume of the soil (retracted), which varies for each matrix potential applied. This adjustment reduces errors, mainly in determining the permanent wilt point and available water.

Keywords: retraction; available water; estimation.

Ajuste de las curvas de retención de agua para suelos retráctiles

Resumen

La curva de retención de agua del suelo (SWRC) relaciona la humedad con la energía de retención de agua (potencial de matriz). En suelos retráctiles, este volumen varía de acuerdo con el secado o la humectación del suelo, lo que puede provocar errores en la humedad calculada. El objetivo de este estudio es cuantificar la variación de volumen en los suelos retráctiles y elaborar el SWRC a través del método tradicional, que no considera la retracción del suelo, y un segundo método, que considera la retracción. Se han recolectado muestras de suelo en los horizontes A y B de seis perfiles de suelo. Por lo tanto, para suelos retráctiles, se recomienda que el ajuste del SWRC se lleve a cabo considerando el volumen real del suelo (retraído), que varía para cada potencial de matriz aplicado. Este ajuste reduce los errores, principalmente al determinar el punto de marchitez permanente y el agua disponible.

Palabras clave: retraccíon; agua disponible; estimar.

1. Introduction

Soil quality is fundamental for the development of life, especially for its function of gradually retaining and releasing water for streams and rivers, as well as for plants. The capacity of the soil to retain water is assessed through the water retention curve (SWRC), which relates the potential matrix energy (ψ m) with the volumetric moisture (Uv). It is used to analyze infiltration, profile redistribution, availability to plants and irrigation management [23].

SWRC is specific to each soil, being influenced by granulometry, mineralogy, organic matter content [14] and soil structure, which alter the distribution and size of pores. The structure, in turn, is modified during the drying of the soil, especially in those that have a retractable character, forming pronounced vertical cracks in the exposed face of the profile [7]. Retraction/expansion is a behavior observed more intensely in Vertisolos [2] but also in soils with a retractable character [20], which are representative, mainly in the Southern Region of Brazil [17,18,22]. These authors reported that drying reduced 40% of the volume of retractable soils in southern Brazil, and 50% in a Vertisol in Rio Grande do Sul [19]. This reduction in the volume of soil during drying can introduce an error in determining the volumetric moisture of the soil.

Therefore, it is necessary to quantify the contraction when the soil is subjected to drying during the analysis of SWRC. This can

How to cite: da Silva, L., Albuquerque, J.A., Sequinatto, L. and Bortolini, D., Adjusting the water retention curve for retractable soils.. DYNA, 88(218), pp. 136-142, July - September, 2021.

be accomplished by combining the determination of SWRC [3] with the soil retraction index [17,18]. Thus, for each ψ m applied to the sample, the moisture (Uv) retained and the volume of the soil retracted are determined. The retracted volume is measured by filling the voids that arise during the retraction of the soil mass with fine sand, with a diameter less than 0.0425 mm [17,18].

The hypothesis of this work is that, in retractable soils, the soil moisture is underestimated at the lowest potentials of the SWRC, and thus, the available water is overestimated.

The goal of this work was to analyze the variations in the water retention curves in soils with retractable character and to propose a calculation procedure to adjust the SWRC that considers the retracted volume in each matrix potential (ψ m) applied to the soil.

2. Material and methods

2.1 Soils and collection site

The A and B horizons of Oxisols and Nitosols previously identified with a retractable character and the horizon A of a Vertisol have been sampled, used as a comparison, since their retraction/expansion characteristic is known to come from the

Table 1.

Soil identification and location with the presence of retractable character and Vertisol.

presence of type 2:1 minerals [1]. Location, identification and characterization of the analyzed soils are presented in the Table 1 and Table 2.

In each soil, a profile has been exposed and horizons A and B have been separated, described [16] and samples with preserved structure have been collected in metal cylinders with a diameter of 0.06 m, height of 0.05 m and volume 0.0141 m³. At lab the soils in the cylinders have been prepared, removing the excess soil. Next, they have been saturated for 48 hours and submitted to the retraction and water retention analysis as described below.

2.2 Determination of soil retraction

In order to quantify the retraction, 21 samples have been collected in each horizon A and B in the six retractable soils plus 21 in the horizon A of Vertisolo (total of 273 cylinders). Shrinkage has been assessed using the sand ring method [17] [18]. Pre-tests demonstrated that the retraction only begun to be noticeable after ψ m of -10 kPa, thus measuring the retracted volume in ψ m between -10 to -1,500 kPa, as detailed below:

Identification	Classification	Place	Location
NB _{PAI} *	Typical Nitossolo Bruno Dystrophic	Painel – SC	27°53'42" S 50°07'45" W
LB _{VAC}	Typical Latossolo Bruno Dystrophic	Vacaria – RS	28°30'47"S 50°53'37" W
LV_{CN}	Dystrophic Red Latosol	Campos Novos – SC	27°22'35" S 51°05'27" W
NB _{PS}	Humic Nitossolo Bruno Dystrophic	Ponte Serrada - SC	26°51'23" S 52°02'33" W
NB _{CUR}	Humic Nitossolo Bruno Dystrophic	Curitibanos – SC	27°22'12" S 50°34'46" W
LB _{VAR}	Nitossolic Latossolo Bruno Dystrophic	Vargeão – SC	26°51'13" S 52°05'56" W
VE _{SL}	Typical Ortic Ebanic Vertisol	Santana do Livramento - RS	30°42'05"S 55°49'43" W

 $\label{eq:shared} $$^{NB_{Fai}: Reddish-brown Nitosol - Painel/SC; LB_{VAC}: Reddish-brown Latosol - Vacaria/RS; LV_{CN}: Red Latosol - Campos Novos/SC; NB_{FS}: Reddish-brown Nitosol - Ponte Serrada/SC; NB_{CUR}: Reddish-brown Nitosol - Curitibanos/SC; LB_{VAR}: Reddish-brown Latosol - Vargeão/SC; VE_{SL} Dark Vertisol - Santana do Livramento/RS Source: The Authors.$

Table 2.

Identification, lithology, Group, geological formation (Serra Geral), chronology, altitude, local relief and primary vegetation of the analyzed soils.

Identifi****	Lithology	Group	Chr.	Altitude, m	Terrain Place	Vegetation Primary
NB _{PAI} *	Basalt	São Bento	J/K	1150	Wavy	Mixed Ombrophilous Forest
LB _{VAC**}	Basalt	São Bento	J/K	1000	Soft wavy	Field
LV_{CN}^{*}	Basalt	São Bento	J/T	939	Soft wavy	Native Field
NB_{PS}^{*}	Dacite	São Bento	J/T	1065	Soft wavy	Mixed Ombrophilous Forest
NB_{CUR}^{*}	Basalt	São Bento	J/K	1018	Soft wavy	Native Field
LB _{VAR} *	Dacite	São Bento	J/K	1043	Soft wavy	Mixed Ombrophilous Forest
VE _{SL} ***	Basalt	Escobar	C/I	233	Flat to Wavy	Native Field

* [6]; ** [5]; *** [13].

**** NBPai: Reddish-brown Nitosol - Painel/SC; LBVAC: Reddish-brown Latosol - Vacaria/RS; LVCN: Red Latosol - Campos Novos/SC; NBPS: Reddish-brown Nitosol - Ponte Serrada/SC; NBCUR: Reddish-brown Nitosol - Curitibanos/SC; LBVAR: Reddish-brown Latosol - Vargeão/SC; VESL Dark Vertisol – Santana do Livramento/RS

***** Chronology: J/K=Jurassic/Cretaceous; J/T= Jurassic/Triassic; C/I= Cretaceous/ Inferior Source: The Authors.

- a) The volumetric cylinders have been saturated, weighed and placed on Richards' sand table and chamber (as determined ψm);
- b) The applied ψ m were of -10 kPa (on a sand table), -33,

-100 -500, -1000 and -1,500 kPa (in Richards Chamber). After each ψ m was applied, three samples have been taken from Richards chambers in order to determine the retracted volume [17,18];

- c) After determining the shrinkage, the samples have been dried in an oven at 105°C;
- d) The retraction index for each ψm has been calculated [17] [18]:

$$IR = 1 - [(va)/(vr)]^{1/3}$$
(1)

where: RI = retraction index; va = volume of the ring (cm³); and vr = retracted volume (cm³).

2.3 Soil water retention curve

In order to quantify the retraction, 21 samples have been collected in each horizon A and B in the six retractable soils plus 21 in the horizon A of Vertisolo (total of 39 cylinders).

- a) The volumetric cylinders have been saturated, weighed and placed on sand table and next in Richards' chamber;
- b) On the sand table [9] ψ m have been applied at 1, -6 and -10 kPa;
- c) Then, they were taken to Richards' Chamber [15] at a ψ m of 33, 100, 500, 1,000 and 1,500 kPa;
- After each applied ψm the moisture reached equilibrium, and the samples have been weighed on a precision scale;
- Afterwards, the samples have been dried in an oven at 105°C, and the retained volumetric moisture in each ψm has been calculated.

The volumetric humidity, in each ψ m, has been calculated using two procedures. First, the volume of retained water in each ψ m has been divided by the initial volume of soil (equal to the volume of the volumetric cylinder = 0.0141 m³). In the second, the volume of retained water in each ψ m has been divided by the volume of soil retracted (that is, a variable volume for each applied tension).

Thus, for each soil and horizon, two water retention curves have been adjusted and the parameters obtained: field capacity (FC), retained volumetric humidity at ψ m of -10 kPa; permanent wilting point (PWP), retained volumetric moisture at ψ m of -1,500 kPa; and available water (AW), retained volumetric humidity between ψ m of 10 and -1,500 kPa.

To adjust the SWRC, the model proposed by [24] has been used:

$$\theta = \theta_r + \frac{\left(\theta_s - \theta_r\right)}{\left(1 + \left(\alpha h\right)^n\right)^n} \tag{2}$$

Where: θ = volumetric humidity; θ r = residual volumetric humidity; θ s = volumetric moisture of the saturated soil; h = matrix potential (ψ m); α , m, n = empirical parameters for adjusting the equation.

2.4 Granulometry

The determination of the granulometric distribution of the soil, for the quantification of sand (2-0.053 mm), silt (0.053-0.002 mm) and clay (<0.002 mm) has been performed using the Pipette method [4], with three replications.

2.5 Soil density and porosity

After collecting soil with the volumetric cylinders, the excess soil was removed, saturated for 48 hours, taken to the sand table at a tension of -6 kPa, and taken to the oven

to dry at 105°C until reaching constant weight. After each stage the samples were weighed. With these determinations (saturated mass, -6 kPa and kiln-dried) soil density (Sd), macroporosity (Macro), microporosity (Micro) and total soil porosity (TP) have been calculated [20]. The particle density (Pd) has been determined by volumetric flask procedure [20] in samples with altered structure and the ground soil in agate gral.

3. Results and discussion

3.1 Soil characterization

The studied soils are classified in the clayey and very clayey textural classes (Table 3), i.e., low sand content and high clay content, meeting the specifications for the retractable character. The granulometry has been influenced by the source material as well as by the weathering process that occurs in the analyzed soils.

The more clayey texture resulted in a lower density of retractable soils, between 0.80 and 0.96 g cm³ on horizon A and between 0.98 and 1.21 g cm⁻³ on horizon B (Table 4).

In the Vertisol, the density of horizon A was of 1.10 g cm⁻³. We know that clay soils with densities less than 1.25 g cm³ hardly present restrictions to the growth and development of roots [12]. In the case of Vertissolo, even with density not restrictive to root growth, its mineralogical characteristics provide physical limitations such as cracking during drier periods, high plasticity and stickiness, and low hydraulic conductivity and infiltration rate [10].

Table 3.

Textural classification and average levels of sand, silt, and clay from the analyzed soils.

Soil	Horizon	Texture	Clay	Silt	Sand
				g kg	-1
NB _{PAI}	А	Clayey	600	300	100
NB _{PAI}	В	Very clayey	730	50	220
LB _{VAC}	А	Clayey	590	300	100
LB _{VAC}	В	Very clayey	640	280	80
LV _{CN}	А	Clayey	560	310	130
LV _{CN}	В	Clayey	580	250	170
NB _{PS}	А	Clayey	600	320	80
NB _{PS}	В	Very clayey	770	210	20
NB _{CUR}	А	Clayey	600	330	70
NB _{CUR}	В	Very clayey	590	290	20
LB _{VAR}	А	Clayey	450	320	230
LB _{VAR}	В	Very clayey	740	200	60
VEst	А	Clavey	565	377	58

NBPAI: Reddish-brown Nitosol - Painel/SC; LBVAC: Reddish-brown Latosol - Vacaria/RS; LVCN: Red Latosol - Campos Novos/SC; NBPS: Reddish-brown Nitosol - Ponte Serrada/SC; NBCUR: Reddish-brown Nitosol - Curitibanos/SC; LBVAR: Reddish-brown Latosol - Vargeão/SC; VESL: Dark Vertisol - Santana do Livramento/RS Source: The Authors.

Table 4.

Soil density (Sd), particle density (Pd), total porosity (TP), macroporosity (Macro), microporosity (Micro) of analyzed soils.

Soil	Hor	Sd	Pd	TP	Macro	Micro	MOS
		g (cm ⁻³		cm ³ cm ⁻³	3	g kg ⁻¹
NB _{PAI}	А	0.91	2.64	0.68	0.16	0.52	75
NB _{PAI}	В	1.21	2.66	0.60	0.10	0.50	47
LB _{VAC}	А	0.94	2.60	0.70	0.18	0.52	92
LB _{VAC}	В	1.06	2.65	0.66	0.17	0.49	74

LV_{CN}	Α	0.82	2.57	0.67	0.30	0.37	65
LV _{CN}	В	1.02	2.60	0.63	0.16	0.47	63
NB _{PS}	Α	0.80	2.61	0.66	0.26	0.40	74
NB _{PS}	В	0.98	2.66	0.63	0.12	0.51	44
NB _{CUR}	Α	0.90	2.52	0.64	0.19	0.45	72
NB _{CUR}	В	1.07	2.60	0.61	0.17	0.44	40
LB _{VAR}	Α	0.96	2.49	0.61	0.14	0.47	95
LB _{VAR}	В	1.01	2.69	0.66	0.21	0.45	26
VE _{SL}	Α	1.10	2.22	0.62	0.07	0.55	109

* NB_{PAI}: Reddish-brown Nitosol - Painel/SC; LB_{VAC}: Reddish-brown Latosol - Vacaria/RS; LV_{CN}: Red Latosol - Campos Novos/SC; NB_{PS}: Reddish-brown Nitosol - Ponte Serrada/SC; NB_{CUR}: Reddish-brown Nitosol - Curitibanos/SC; LB_{VAR}: Reddish-brown Latosol - Vargeão/SC; VE_{SL}: Dark Vertisol – Santana do Livramento/RS. Source: The Authors.

The particle density (Pd) of horizon A varied between 2.49 to 2.66 g cm⁻³, in horizon B between 2.60 to 2.69 g cm⁻³ (Table 4), while in Vertisol it was lower, 2.22 g cm³. In general, Pd in horizon A was lower than in horizon B, because in those the MOS content is higher, a component that has a density around 1.20 g cm⁻³ [8] and acts in structuring of the soil. On horizon A of the LB_{VAR}, which is the horizon with the highest SOM content, the Pd was lower, 2.49 g cm⁻³.

The TP of most soils and horizons was high, generally exceeding 0.60 cm³ cm⁻³ (Table 4), mainly in horizon A, due to the MOS content and the effect of this component on aggregation. With the exception of LBVar, which can be attributed to the predominance of strong, very small, granular structure (coffee powder) on horizon B of this soil. The high TP was due to the magnitude of the microporosity, which ranged from 0.41 to 0.51 cm⁻³ for Nitosols and from 0.37 to 0.52 cm³ cm⁻³ for Oxisols. Macro was smaller and varied between 0.10 and 0.30 cm³ cm⁻³, being generally larger in horizon A. The Vertissolo presents a high TP of 0.62 cm³ cm⁻³, also with a predominance of Micropores (0.55 cm³ cm⁻³) in relation to macropores (0.07 cm³ cm⁻³), similar to that found by [11] for a Vertissolo from RS.

3.2 Characterization of soil retraction

The retraction differed among the soils as shown in Fig. 1. The Vertissolo RI was higher than the other retractable soils, a characteristic already visually observed during the description of the soil in the field. Within soils with a retractable character, the horizons A of NBPS and LBVAC presented higher RI, while horizons A and B of the LVCN presented lower RI. According to [21], the mineralogical composition of horizons A and B of the clay fraction was similar for the analyzed soils, in general with intense reflexes related to kaolinite, and with less intensity related to the presence of type 2:1 clay minerals or chlorite and oxides, goethite and/or hematite.



Soil horizons

Figure 1. Soil* retraction index, determined using the ring filling method with sand - RI.

* NB_{PAI}: Reddish-brown Nitosol - Painel/SC; LB_{VAC}: Reddish-brown Latosol - Vacaria/RS; LV_{CN}: Red Latosol - Campos Novos/SC; NB_{PS}: Reddish-brown Nitosol - Ponte Serrada/SC; NB_{CUR}: Reddish-brown Nitosol - Curitibanos/SC; LB_{VAR}: Reddish-brown Latosol - Vargeão/SC; VE_{SL}: Dark Vertisol – Santana do Livramento/RS.

Source: The Authors.

While analyzing the retraction evolution with drying, it has already been detected at ψ m -10 and -100 kPa, with the exception of Vertissolo. On soils NBPAI A, LBVAC B, NBCUR A e B and NBPS A and B, it has been detected at a ψ m of -10 kPa, while soils LBVAC A and LB_{var} A and B have been detected at a ψ m of -100 kPa. In the case of Vertissolo, the retraction was only noticeable at 1,000 kPa (Table 5). Upon reaching a ψ m of -1,500 kPa, retraction varied between 3,7 (LBVACA) and 8,7% (NBPS B). The greatest retraction occurred in ψ m below -1,500 kPa, as can be observed by the large difference between the retracted volume between -1,500 kPa and the dry soil at 105°C. The soils LBVAC A and NBPS A showed the largest retracted volume at the end of drying, close to 46%, while LVCN A and B presented the lowest retraction, close to 20%.

Table 5.

Soil retraction (%) as a function of water matrix potential during the drying of the analyzed soils.

Identifica	-10	-33	-100	-500	-1000	-1500	105°
tion	kPa	kPa	kPa	kPa	kPa	kPa	С
LV _{CN} A	0	2.2	2.3	3.5	5.2	6.4	24
LV _{CN} B	0	2.5	2.5	3.4	4.2	5.1	20
NB _{PAI} A	2.3	2.9	2.9	3.3	5.7	7.2	40
NB _{PAI} B	0	2.9	3.0	3.1	4.7	6.4	33
LB _{VAC} A	0	0	2.5	2.6	2.7	3.7	47
LB _{VAC} B	1.4	1.4	2.2	2.2	2.9	4.2	29
NB _{CUR} A	1.6	1.6	1.6	2.2	4.3	6.0	37
NB _{CUR} B	1.6	2.7	2.7	2.9	5.4	5.6	39
NB _{PS} A	1.8	2.2	2.2	4.9	5.1	5.1	46
NB _{PS} B	2.4	2.9	3.1	3.4	8.7	8.7	40
LB _{VAR} A	0	0	1.0	1.8	1.9	3.7	44
LB _{VAR} B	0	0	2.7	2.9	3.8	5.6	43
VE _{SL}	0	0	0	0	2.2	4.8	51

* NB_{PAI}: Reddish-brown Nitosol - Painel/SC; LB_{VAC}: Reddish-brown Latosol - Vacaria/RS; LV_{CN}: Red Latosol - Campos Novos/SC; NB_{PS}: Reddish-brown Nitosol - Ponte Serrada/SC; NB_{CUR}: Reddish-brown Nitosol - Curitibanos/SC; LB_{VAR}: Reddish-brown Latosol - Vargeão/SC; VE_{SL}: Dark Vertisol - Santana do Livramento/RS Source: The Authors.

As shown by Table 5, we observed that the retraction started at a macroscopic level, that is, when the water is retained between a ψ m from -10 to -100 kPa. However, it gets intensified in ψ m less than -100 kPa, suggesting that the retraction is related to processes that occur in soil microstructures, more precisely when the removal of water occurs in pores smaller than 0.2 μ m in diameter, normally related to the pores drained at voltages above -1500 kPa. It is important to note that the water retained can be located in capillary pores (with equivalent diameters mentioned above), as well as adsorbed on the surfaces of mineral and organic particles.

As noted above, all soils have retracted, to a lesser or greater degree. Thus, the water retention curves calculated by the standard procedure differed from the adjusted retention curves considering soil retraction (Fig. 2). The water retention curves calculated by the two procedures begin to differ from each other in ψ m less than -100 kPa. However, in Vertisolo this difference occurs only for ψ m less than -1000 kPa, that is, in the driest branch of the curve.



Figure 2 - Soil water retention curves adjusted by standard method (LVCN(A)) and adjusted by volume of soil retracted (LVCN(A)adj). Source: The Authors.

Table 6.

Parameters of the water retention curve of van Genuchten model, for horizons A and B of the analyzed soils, adjusted by the standard method and adjusted by the method that considers the volume of soil retracted.

	Ψ	Ψ-			
Identification	saturation	1,500 kPa	α	n	m
$LV_{CN}(A)$	0.634	0.424	2.445	1.336	0.252
LV _{CN} (A)aj	0,634	0.451	2.758	1.399	0.285
$LV_{CN}(B)$	0.642	0.448	2.602	1.244	0.196
LV _{CN} (B)aj	0.642	0.471	3.839	1.255	0.203
$NB_{PAI}(A)$	0.650	0.434	2.075	1.313	0.238
NB _{PAI} (A)aj	0.650	0.465	2.829	1.330	0.248
$NB_{PAI}(B)$	0.671	0.448	5.432	1.237	0.192
NB _{PAI} (B)aj	0.671	0.477	8.969	1.244	0.196
$LB_{VAC}(A)$	0.646	0.322	7.122	1.265	0.209
LB _{VAC} (A)aj	0.646	0.334	8.128	1.270	0.212
$LB_{VAC}(B)$	0.660	0.354	5.377	1.260	0.206
LB _{VAC} (B)aj	0.660	0.369	6.153	1.264	0.209
NB _{CUR} (A)	0.693	0.413	4.391	1.271	0.213
NB _{CUR} (A)aj	0.693	0.438	4.890	1.297	0.229
NB _{CUR} (B)	0.658	0.49	6.207	1.298	0.230
NB _{CUR} (B)aj	0.658	0.432	7.847	1.317	0.245
$NB_{PS}(A)$	0.733	0.390	4.007	1.284	0.221
NB _{PS} (A)aj	0.733	0.405	5.116	1.271	0.213
$NB_{PS}(B)$	0.668	0.405	6.653	1.267	0.211
NB _{PS} (B)aj	0.668	0.426	11.500	1.246	0.197
LB _{VAR} (A)	0.685	0.442	0.424	1.335	0.258
LB _{VAR} (A)aj	0.685	0.458	0.505	1.341	0.254
LB _{VAR} (B)	0.676	0.349	2.293	1.326	0.246
LB _{VAR} (B)aj	0.676	0.368	2.447	1.351	0.260
VE _{SL}	0.627	0.451	2.978	1.222	0.182
VE _{SL} aj	0.627	0.472	3.281	1.265	0.210

Source: The Authors.

We suggest that the change in the mass of the retractable soils is related, at first, to the soil structure, in a rearrangement of the microporosity. In the Vertissols, the retraction of the soil mass occurs close to -1000 kPa and, therefore, is related to very small pores, possibly formed by very small clay particles, which may be related to its mineralogy, since smectite has smaller dimensions than kaolinite [2].

The adjustment parameters of the van Genuchten equation α , n and m were higher in standard SWRC than in adjusted SWRC considering soil retraction. This increase indicates that there has been a reduction in water loss at the highest Ψ m (highest α value), that is, the SWRC is less inclined than the larger Ψ m, and that the water content decreases more rapidly at the end of the SWRC (values of n and m greater) that is, it is more inclined (Fig. 2). It is clear that, with the adjustment, considering the volume of soil retracted, the humidity in the PWP (-1,500 kPa) is higher and consequently the AW is lower, since the field capacity is not altered.

While comparing the curves obtained using the standard method with the adjusted curves considering the volume of soil retracted (Fig. 2), the FC of some soils have not differed or differed little (Fig. 3). However, we observed that for PWP these differences were greater than for FC. The smallest difference in FC occurs, as the retraction of the soil begins to become noticeable only after Ψ m of -10 kPa, and expands in the drier branch of the curve, close to the PWP. As a result, the higher PWP in the adjusted SWRC causes the available water content (AW) to decrease (Fig. 4). In some soils, the



Figure 3. Volumetric humidity in the field capacity (FC) and in the permanent wilt point (PWP) for horizons A and B of the retractable soils, calculated by the standard method in relation to the method adjusted by the soil retraction.

Source: The Authors.



Soil horizons

Figure 4. Available water content calculated by default method and calculated by the method adjusted by the soil retraction, expressed in content, for horizons A and B of retractable soils. Source: The Authors.

Table 7.

Difference between the available volumetric moisture calculated by the default method in relation to the one calculated by the method adjusted by soil retraction.

Soils	AW (m ³ m ⁻³)	AW (%)
LV _{CN} A	0.027	41
LV _{CN} B	0.023	27
NB _{PAI} A	0.019	25
$NB_{PAI}B$	0.029	39
LB _{VAC} A	0.012	15
$LB_{VAC}B$	0.008	9
NB _{CUR} A	0.017	20
NB _{CUR} B	0.015	26
NB _{PS} A	0.006	6
NB _{PS} B	0.009	13
LB _{VAR} A	0.016	12
LB _{VAR} B	0.020	18
VE _{SL}	0.022	24

Source: The Authors.

difference between methods was up to 40% in AW. This is more important when the crops are experiencing a period of water deficit or when the crop is more susceptible to water shortages.

For those who are irrigating and keeping the soil more humid, the difference in the adjustments is smaller, because in potentials between -10 and -100 kPa the difference between the curves is small. However, for crops without irrigation, when the soil is drying out, the amount of water available should be calculated considering the volume of soil retracted, which is less than the default method. Thus, for retractable soils, we suggest adjusting the water retention curve considering the volume of soil retracted (adjusted method), in order to obtain the volumetric humidity closer to what would actually be available to the plants.

4. Conclusions

The water retention curves differ depending on soil retraction, a difference that starts around -10 kPa and increases in the drier branch of the curves. The most pronounced shrinkage starts with the water leaving the pores smaller than 2 μ m and intensifies in the porosity range close to 0.1 μ m.

In soils with a retractable character, the SWRC must be adjusted considering the volume of soil retracted. With this adjustment procedure, the field capacity remains similar to the default method, however the permanent wilt point is higher and the available volumetric humidity is lower.

References

- Bertolazi V.T., Concentração de argilominerais 2:1 em solos subtropicais, Disertación Maestría em Ciencias del Suelo, Universidade Federal de Santa Maria, Santa Maria, Brasil, 2013, 34 P.
- [2] Coulombe, C.E., Dixon, J.B. and Wilding, L.P., Mineralogy and chemistry of Vertisolse, en: Ahmad, N. and Mermut, A., Eds., Vertosols and technologies for their management, Elsevier Science, Amsterdam, Netherlands, 1996, pp.115-200.
- [3] Dane, J.H. and Hopmans, J.W., Pressure plate extractor, en: Dane, J.H. and Topp, C., Eds., Methods of soil analysis: physical methods, Soil Science Society of America, Madison, USA, 2002. pp. 688-690.
- [4] Day, P.R., 1965. Pipette Method, en: Dane, J.H. and Topp, C., Eds., Methods of soil analysis. Part 4. Physical methods, Soil Science

Society of America, Madison, USA, 2002. pp. 643-670.

- [5] Empresa Brasileira de Pesquisa Agropecuária-EMBRAPA. Centro Nacional de Pesquisa de Solos. Guia de excursão de estudos de solos nos estados do Rio Grande do Sul. Santa Catarina e Paraná, Curitiba, Embrapa Florestas, 2000.
- [6] Empresa Brasileira de Pesquisa Agropecuária-EMBRAPA, Empresa de Pesquisa Agropecuária-EPAGRI. VIII RCC Reunião Nacional de Correlação e Classificação de Solos (Santa Catarina), 1^m ed., Rio de Janeiro, Embrapa, 2008.
- [7] Empresa Brasileira de Pesquisa Agropecuária-EMBRAPA. Centro Nacional de Pesquisa de Solos. Sistema Brasileiro de Classificação de Solos – 5^{ta} ed. Rev. Ampl. – Brasília. Embrapa, 2018.
- [8] Ferreira, M.M., Caracterização física do solo, em: van Lier, Q.J. Ed., Física do Solo, Sociedade Brasileira de Ciência do Solo, Viçosa, Brasil, 2010. pp. 1-27.
- [9] Leamer, R.W., and Shaw, B.A., Simple apparatus for measuring noncapillary porosity in extensive scale. Journal of American Society of Agronomy, 33, pp. 1003-1008, 1941.
- [10] Moustakas, N.K.A., Study of Vertisol genesis in North Eastern Greece. Catena, 92, pp. 208-15, 2012, DOI: 10.1016/j.catena.2011.12.011
- [11] Pértile, P., Propriedades mecânicas em micro e mesoescala de solos do Rio Grande do Sul. Tesis Doctorado em Ciencias del Suelo, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brasil, 2015.
- [12] Reichert, J.M., Reinert, D.J. and Braida, J.A., Qualidade dos solos e sustentabilidade de sistemas agrícolas, Revista Ciência & Ambiente, 27, pp. 29-48, 2003.
- [13] Reinert, D.J., Reichert, J.M., Azevedo, A.C., Dalmolin, R.S.D. and Pedron F.A., Principais solos da depressão central e campanha do Rio Grande do Sul: Guia de Excursão, 2ª. ed, Gráfica Universitária, Santa Maria, Brasil, 2007.
- [14] Ribeiro, K.D., Menezes, S.M., Mesquita, M.G.B.F. and Sampaio, F.M.T., Propriedades físicas do solo, influenciadas pela distribuição de poros, de seis classes de solos da região de Lavras-MG, Ciência Agrotécnica, 31, pp. 1167-1175, 2007, DOI: 10.1590/S1413-70542007000400033
- [15] Richards, L., Methods of measuring moisture tension, Soil Science, 58(1), pp. 95-112, 1949.
- [16] Santos, R.D., Santos, H.G., Ker, J.C., Anjos, L.H.C. and Shimizu, S.H., Manual de descrição e coleta de solo no campo. 7^a ed., Sociedade Brasileira de Ciência do Solo, Viçosa, Brasil, 2015.
- [17] Silva, L., Sequinatto, L., Almeida, J.A. and Bortolini, D., Methods for quantifying shrinkage in Latossolos (Ferralsols) and Nitossolos (Nitisols) in southern Brazil, Revista Brasileira de Ciência do Solo, 41, art. e0160364, 2017, DOI: 10.1590/18069657rbcs20160364
- [18] Silva, L., Caráter retrátil de Nitossolos e Latossolos do Sul do Brasil, Disertación Maestría em Ciencias del Suelo, Universidade do Estado de Santa Catarina, Lages, Brasil, 2015, 131 P.
- [19] Silva, L., Retração do solo e relação com as propriedades físicohídricas de Latossolos e Nitossolos do Sul do Brasil, Tesis Doctorado em Ciencias del Suelo, Universidade do Estado de Santa Catarina, Lages, Brasil, 2019.
- [20] Teixeira, P.C., Donagemma, G.K., Fontana, A. and Teixeira, W.G., Manual de métodos de análise de solo, 3^{ra} ed., Rev. e Ampl, Brasília, Embrapa, 2017.
- [21] Testoni, S., Mineralogia da fração argila de Latossolos e Nitossolos Brunos e Vermelhos com caráter retrátil do sul do Brasil, Disertación Maestría em Ciencias del Suelo, Universidade do Estado de Santa Catarina, Lages, Brasil, 2015,
- [22] Testoni, S.A., Almeida, J.A., Silva, L. and Andrade, G.R.P., Clay mineralogy of Brazilian oxisols with shrinkage properties. Revista Brasileira Ciência do Solo; 41, art. e0160487, 2017. DOI: 10.1590/18069657rbcs20160487
- [23] Tormena, C.A. and Silva, A.P., Incorporação da densidade no ajuste de dois modelos à curva de retenção de água no solo, Revista Brasileira de Ciência do Solo, 26, pp 305-314. 2002, DOI: 10.1590/S0100-06832002000200003
- [24] van Genuchten, M.T.H., A Closed-Form equation for predicting the hidraulic conductivity of unsaturated soils, Soil Science Society American Journal, 44, pp 892-897, 1980.

L. da Silva, Eng in Agronomist in 2012 from the Universidade Tecnológica Federal do Paraná, Pato Branco, Brazil, MSc. in Soil Science in 2015 from the Universidade do Estado de Santa Catarina, Phd in Soil Science in 2019 from Universidade do Estado de Santa Catarina. From 2019 to 2021, she worked for the SEED Santa Santa Catarina, Brazil. ORCID: 0000-0002-0051-9297

J.A. Albuquerque, Eng in Agronomist in 1990 from the Universidade Federal de Santa Maria, Santa Maria, Brazil, MSc. in Soil Science in 1994 from the Universidade Federal de Santa Maria, Phd in Soil Science in 1998 from Universidade Federal do Rio Grande do Sul. From 1997 to actualy, he worked for the Universidade do Estado de Santa Catarina, Brazil. ORCID: 0000-0002-7387-0336

L. Sequinatto, Eng in Agronomist in 2004 from the Universidade Federal de Santa Maria, Santa Maria, Brazil, MSc. in Soil Science in 2007 from the Universidade Federal de Santa Maria, Phd in Soil Science in 2010 from Universidade Federal do Rio Grande do Sul. From 2011 to actualy, she worked for the Universidade do Estado de Santa Catarina, Brazil. ORCID: 0000-0001-7389-9780

D. Bortolini, Eng in Agronomist in 2010 from the Universidade Tecnológica Federal do Paraná, Pato Branco, Brazil, MSc. in Agronomist in 2012 from the Universidade Tecnológica Federal do Paraná, Phd in Soil Science in 2016 from Universidade do Estado de Santa Catarina. From 2017 to 2020, she worked for the Brazilian Institute of Geography and Statistics, Brazil, he currently works in CELESC, Brazil. ORCID:0000-0003-0895-1401

142