

Implementation of a Non-Invasive Bioprospecting Protocol for Isolation of *Lactobacillus* from Feces of Hens Under Foraging Conditions

Simón Robledo-Cardona¹, Sabina Ramírez-Hincapié² and Javier Correa-Álvarez³

Received: 19-04-2018 | Accepted: 31-10-2018 | Online: 23-11-2018

doi:10.17230/ingciencia.14.28.4

Abstract

In animal production, probiotics seek to replace the use of antibiotics, while diminishing mortality and morbidity rates to raise productivity. Probiotics constitute a natural alternative that, in contrast with antibiotics, neither produces pathogen resistance, nor leaves chemical residues in the final product. Several bacteria, including some belonging to the genus *Lactobacillus* have been described as probiotics with high potential. A non-invasive bioprospecting protocol aimed for the isolation and characterization of lactobacilli from chicken feces was established. Fecal samples were collected from the ground. These were diluted and cultured in LAB selective medium. Colonies were identified by three methods: Gram stain, MALDI-TOF MS and sequencing of 16S rRNA gene. An initial probiotic potential of lactobacilli isolates was determined via antagonism tests using five enteropathogen reference strains: *Staphylococcus aureus*, *Enterococcus*

¹ Universidad EAFIT, srobled1@eafit.edu.co, <https://orcid.org/0000-0003-1055-0375>, Medellín, Colombia.

² Universidad EAFIT, sramir30@eafit.edu.co, <https://orcid.org/0000-0001-5258-1239>, Medellín, Colombia.

³ Universidad EAFIT, jcorre38@eafit.edu.co, <https://orcid.org/0000-0001-9009-823X>, Medellín, Colombia.

faecium, *Candida albicans*, *Pseudomonas* spp. and *Salmonella* spp. 24 isolates belonging to four *Lactobacillus* species were identified by MALDI-TOF MS. BLAST of 16S rRNA gene of eight randomly selected isolates, confirmed MALDI-TOF MS identification. Five of these eight isolates inhibited the growth of at least one of the pathogenic strains used, three isolates of *Lactobacillus plantarum* and two of *Lactobacillus salivarius*. Our protocol achieved 21 lactobacilli per 100 isolates performance, greatly surpassing the normal percentage of lactobacilli in chicken gut microbiome, that so, its implementation would facilitate the isolation and identification of new probiotic strains from feces.

Keywords: Bioprospecting; probiotics; lactic acid bacteria (LAB); antagonism test.

Implementación de un protocolo de bioprospección no invasivo para el aislamiento de *Lactobacillus* a partir de heces de gallinas en condiciones de forrajeo

Resumen

En la producción animal, los probióticos son una alternativa para reemplazar el uso de antibióticos, ya que disminuyen las tasas de morbilidad y mortalidad, aumentando al mismo tiempo la productividad. Así mismo, los probióticos constituyen una alternativa natural que, en contraste con los antibióticos, no generan patógenos resistentes ni dejan residuos químicos en el producto final. Una variedad de bacterias, pertenecientes al género *Lactobacillus*, han sido descritas como probióticos con alto potencial. Se estableció un protocolo de bioprospección no invasivo dirigido al aislamiento y caracterización de lactobacilos a partir de heces de gallinas. Las muestras fecales fueron colectadas a partir del suelo. Luego fueron diluidas y sembradas en medio selectivo para bacterias ácido lácticas MRS. Las colonias fueron identificadas por tres métodos: tinción de Gram, MALDI-TOF MS y secuenciamiento del gen ARNr 16s. Para caracterizar inicialmente el potencial probiótico de ocho de los aislados de lactobacilos obtenidos se realizaron pruebas de antagonismo usando cinco cepas referencia de patógenos: *Staphylococcus aureus*, *Enterococcus faecium*, *Candida albicans*, *Pseudomonas* spp. y *Salmonella* spp. 24 aislados distribuidos en cuatro especies de *Lactobacillus* fueron identificadas por MALDI-TOF MS. La identificación por MALDI-TOF MS fue confirmada mediante el secuenciamiento del gen ARNr 16s de ocho aislados seleccionados aleatoriamente. Cinco aislados, tres identificados como *Lactobacillus plantarum* y dos como *Lactobacillus salivarius*, inhibieron el crecimiento de al menos uno de los patógenos seleccionados. El protocolo logró un desempeño de 21 lactobacilos por 100 aislados, superando con creces la representación normal de los

lacobacilos en el microbioma gastrointestinal de las gallinas, de manera que su implementación facilitaría los esfuerzos de aislamiento e identificación de nuevas cepas probióticas a partir de heces.

Palabras clave: Bioprospección; probióticos; bacterias ácido lácticas; prueba de antagonismo.

1 Introduction

Antibiotics are widely used in the livestock industry for treatment of infections, disease prevention (prophylaxis), and as growth promoters in sub-therapeutic dosage. Their use began during the fifties in response to the rapidly growing demand of meat worldwide [1]. Nevertheless, it is now well known that antibiotic use in livestock selects resistant pathogens, which may become an overlapped resistance reservoir between animals and humans, ensuing spread in the environment and causing an impact in public health [2].

The first evidence of this reality emerged in 1994, when glycopeptide resistant enterococcus (GRE) were isolated from livestock in Great Britain, Germany and Denmark [3],[4],[5],[2]. Thereafter, the occurrence of GRE infections increased dramatically in hospitals all around Europe and the United States, followed by the fact that these bacteria did not respond to any antibiotics available then, making GRE infections virtually untreatable.

The European Union banned the use of some antibiotics in livestock production, creating a unique opportunity for scientists to research into the effects of selective pressure removal on antimicrobial resistance spread and occurrence [2]. Results proved that after antimicrobial use in livestock ceased, the prevalence of antibiotic-resistant pathogens was significantly reduced, with little to no changes in productivity and animal health [6],[2]. In this context, there has been a fast emergence of alternatives to antibiotics in livestock production within the last three decades [1]. One promising alternative consists in the use of probiotics as prevention tools from infectious diseases, and as growth promoters.

Probiotics are live microorganisms, for the most part bacteria that once ingested, bring about health benefits beyond their inherent nutritional value [7]. These microorganisms have physiological characteristics

that allow them to modify the ecology and the functional properties of the gut microbiome, which in turn, plays a fundamental role in preserving the homeostasis of the digestive system [8].

The gut of domestic animals hosts a wide variety of dense and complex microbial communities composed of bacteria, yeast, protozoa, archaea, and viruses [8]. The range of activities performed by the gut microbiota (GM) is also wide and diverse: vitamin synthesis, bioconversion of toxic compounds, immune system stimulation, and maintenance of gut peristalsis. It also plays a role as a barrier that blocks pathogen colonization of the intestine walls [8],[9].

In the last four decades, a considerable amount of research has been committed to the characterization of the digestive ecosystem. These efforts have led to the description of key bacteria that play a role in the healthy functioning of GM [8],[9].

The possibility of using key bacteria in animal feed as probiotics, that would potentially help to maintain a pathogen resistant gut, has produced considerable interest during the last few years; also taking into account the spread of antibiotic-resistant pathogens and the presence of chemical growth promoters left in meat from substances added to animal feed [8].

The most commonly used probiotics in monogastric animal (birds and pigs) feed are bacteria belonging to the genus *Lactobacillus*, *Enterococcus*, *Pediococcus*, and *Bacillus* [8]. Multiple effects have been attributed to their use in poultry. In broilers, an increased resistance to infectious diseases produced by *Salmonella*, *Escherichia coli*, and *Clostridium perfringens* has been described [10],[11],[12],[13],[14],[15], as well as an increased bone health and fat percentage [16], and a reduction of behaviors related to thermal stress and inflammatory processes [17]. Other benefits described for laying hens consists of increasing feed rates and productivity [18],[19],[20], and improvement of egg quality (reduced level of yolk cholesterol, increased shell thickness and higher egg weight [8](Chaucheyras-Durand & Durand, 2009)) (Chung et al., 2015; Kurtoglu et al., 2004; Mazanko et al., 2017; Tang et al., 2015; Xu et al., 2006).

However, beneficial effects of probiotics have not only been described for poultry, and according to ReportLinker¹ ® (2017), the probiotics market

¹<https://goo.gl/XYJFFw>

for animal feed has a projected value of USD 1,746.5 billion by the year 2022.

Lactic acid bacteria, especially Gram positive bacteria of the genus *Lactobacillus*, possess a high probiotic potential, and naturally inhabit the gut of mammals and birds [21],[22]. Nevertheless, lactobacilli make up less than 1% of the chicken GM [23]. Thus, the main objective of this work consisted in the design and implementation of a bioprospecting protocol for the isolation of lactobacilli from poultry feces, based on morphological and molecular identification, that could increase *Lactobacillus* isolation percentage compared to its representation in the GM. The protocol ends in an initial characterization of the probiotic potential inherent to the isolates obtained using antagonism tests.

The composition of the GM of chickens that inhabit in conditions which allow foraging behavior is different from the GM of chickens that are born in industrial conditions, confined in cages and under a traditional feed. Likewise, the varied diet of foraging chickens imply a healthier and more diverse gut ecosystem that may grant protection against certain pathogens (as the dietary effects on GM that have been observed in humans [24]). Therefore, the chickens that provided the samples live in artisanal production conditions that allow foraging behavior, which provides a varied diet of invertebrates, human food, corn, seeds, fruit, etc., without being restricted to a small cage.

Gram stain is a widely used technique for morphological identification of bacteria. However, to achieve species-level identification more sophisticated tools are required. MALDI-TOF mass spectrometry (MALDI-TOF MS) provides a quick and precise technique for bacterial identification that has been lately used in clinical laboratories around the globe [25]. It is based on laser evaporation of colonies, which generates volatile particles that are sucked by vacuum, with different time flights, into a detector, creating a spectrum unique to each species of bacteria. Another robust method used to identify bacteria is the sequencing of the 16S rRNA gene, a conserved locus whose sequence can be subjected to homology analysis for species assignment [26].

Antagonism tests are a common way to assess the probiotic potential of bacterial isolates. These tests are *in vitro* assays where interacting bacte-

ria exhibit behavioral changes. Particularly, antagonism tests can provide information on the inhibitory effects of isolates against indicator organisms (e.g. pathogenic bacteria to be controlled), thus allowing the selection of promising isolates[27].

The importance of probiotics research aimed for poultry industry lays on the increasing consumption rates of poultry products, and therefore, the need to improve product quality and reduce costs. This need is especially conspicuous in Colombia, where according to the National Federation of Colombian Poultry Farmers (FENAVI, according to its Spanish acronym), the annual consumption of poultry meat per person went from 14.2 kgs in the year 2000, to 32.3 kgs for 2017, doubling in less than 20 years. During the same period, egg consumption per person has increased by 111 units/year.

2 Materials and methods

2.1 Sample processing

Under the collection framework permit granted by the National Environment Licensing Authority (in Spanish, Autoridad Nacional de Licencias Ambientales or ANLA), fecal samples were collected in three Colombian municipalities (Bahía Solano, Chocó: $n = 10$; Itagüí, Antioquia: $n = 30$; and Rionegro, Antioquia: $n = 14$), from chickens that inhabited in artisanal production conditions, which allowed foraging behavior. Samples were stored in Falcon tubes, and were transported in fridges. One gram of each sample was diluted fourfold in sterile water, and each dilution was then cultured on Merck's ® solid MRS selective medium, which has a low pH and certain nutrients that benefit lactic acid bacteria growth.

2.2 Identification

The isolates that grew on selective medium were identified by three methods. First, a Gram stain was used to select colonies that met the morphologic characteristics of lactobacilli (lengthened Gram positive bacilli).

The colonies selected by Gram stain were then identified by a bio-Mérieux® MALDI-TOF MS spectrometer with a 337 nm nitrogen laser (fixed focus) (VITEK® MS). Colonies were identified based on spectra made of protein flight times, whose weight ranged between 1 and 500 kDa. This range includes ribosomal and some housekeeping proteins. Spectra were compared with the VITEK® MS V3.2.0 CE-marked database, using MYLA® middleware software, to obtain identity of colonies. High fidelity identifications were preserved at -70 °C in Merck's ® MRS medium with 20% of glycerol.

To confirm MALDI-TOF MS identification, eight isolates representing all of the *Lactobacillus* species found by mass spectrometry, were randomly selected. For each colony, a total DNA extraction and PCR amplification of the 16S rRNA gene were performed.

DNA was extracted using Ultra Clean ® microbial DNA isolation kit (MO BIO laboratories, inc). DNA was amplified via PCR using Taq Polymerase, recombinant (5 U/L) (Thermo Fisher Scientific, inc). Primers: 27F (AGAGTTTGATCMTGGCTCAG) and 907R (CCGTCAATTCMTTTRAGTTT) (Lane, 1991). The obtained amplicons were sequenced by Sanger method, and consensus sequences were created using Geneious ®. To determine the isolate species, 16S rRNA gene sequences were aligned to NCBI database using BLAST [28].

2.3 Antagonism tests

Antagonism test were performed in order to initially assess the probiotic potential of the same isolates selected for 16S rRNA gene identification. These were tested against five pathogenic strains provided by The Universidad EAFIT strain collection: *Staphylococcus aureus*, *Enterococcus faecium*, *Candida albicans*, *Pseudomonas* spp., and *Salmonella* spp. The eight isolates were cultured in Petri dishes, on solid MRS Merck ® medium

along a negative control strain *Lactobacillus reuteri* (ATCC® 23272). 24 hours later, the five pathogenic strains were inoculated in 1% semi-solid LB broth (lysogenic broth, Merck ®) which was then poured over the MRS solid medium, with already visible lactobacillus colonies that grew overnight. The Petri dishes containing both growth media were incubated at 37 °C for 24 hours. After this incubation period, the presence of inhibition zones was visually established according to the presence or absence of pathogen growth surrounding the vicinity of lactobacillus colonies (Figure S1).

3 Results and discussion

3.1 Identified isolates

114 colonies were obtained from the initial cultures of the diluted fecal samples, and after Gram stain, 49 of the 114 original colonies met the morphologic characteristics of lactobacilli. These 49 colonies were submitted to MALDI-TOF MS, and by this method, 31 were identified with 99% precision. The 31 isolates belonged to the order Lactobacillales, distributed in three families (Enterococcaceae, Streptococcaceae, and Lactobacillaceae), four genera (*Enterococcus*, *Pediococcus*, *Streptococcus*, and *Lactobacillus*), and in eight species of bacteria (Figure 1). 24 isolates belonged to the genus *Lactobacillus* were distributed in four species (Figure 1), which accounts for 77.4% of the 31 isolates identified by mass spectrometry and 21% of the original 114 colonies. This suggests an effective selection by the MRS medium, and by morphological selection using Gram stain, given the fact that the majority of isolates belonged to the genus *Lactobacillus*, and all of them are grouped under the order Lactobacillales and the family Lactobacillaceae (lactic acid, non-sporulating, Gram positive bacteria [29], fulfilling the necessary physiological and morphological characteristics in order to grow in MRS medium and be selected by Gram stain. Furthermore, the family Lactobacillaceae represents less than 1% of the total microbial population of the chicken gut [23] and our bioprospecting protocol was able to augment Lactobacillaceae representation up to 21% of the fecal samples.




Figure 1: Identification process and results. Fecal samples were collected on three locations with different sample size: Bahia Solano (BS), Itagüí (I), Rionegro (R). After four dilutions, 114 colonies grew in selective medium. 49 were singled out by Gram stain for further identification by MALDI-TOF MS. This method identified the species of 31 (63%) isolates with 99% precision (P.), with the exception of 14 isolates assigned to a group of three closely related species, *Lactobacillus plantarum/paraplantarum/pentosus* (*). To confirm MS identification, NCBI-BLAST of 16S rRNA gene was performed with >97% identity results matching MALDI-TOF MS identification.

The exact species for 14 of the 24 *Lactobacillus* isolates couldn't be identified by MALDI-TOF MS. These 14 isolates were assigned to a group of three closely related species, *Lactobacillus plantarum/paraplantarum/pentosus*. In order to confirm MALDI-TOF MS results, eight isolates representing the four lactobacillus species previously identified (including *Lactobacillus plantarum/paraplantarum/pentosus*) were randomly selected for 16S rRNA gene identification. All NCBI-BLAST results of the gene sequences confirmed MALDI-TOF MS identification (Figure 1). Furthermore, within the eight isolates selected, three of them grouped by mass spectrometry in *L. plantarum/paraplantarum/pentosus* cluster (S6, S8, and S14) were identified with 99% of identity (E-value < 0.1) as *L. plantarum*, by majority of matches with 16S rRNA gene sequences BLAST.

3.2 Antagonism tests

Table 1 summarizes antagonism tests results between eight *Lactobacillus*, previously selected for 16S rRNA gene identification, and five pathogenic strains that are commonly found in the gut of chickens.

Table 1: Antagonism tests. Eight *Lactobacillus* isolates tested against five pathogenic strains. +: inhibition zone present; - : inhibition zone absent; EF: *E. faecium*; CA: *C. albicans*; S: *Salmonella* spp.; SA: *S. aureus*; P: *Pseudomonas* spp.

Species	Code	EF	CA	S	SA	P
<i>L. plantarum</i>	S6	+	-	-	-	+
<i>L. plantarum</i>	S8	-	-	+	-	-
<i>L. plantarum</i>	S14	-	-	+	-	-
<i>L. brevis</i>	S31	-	-	-	-	-
<i>L. brevis</i>	S107	-	-	-	-	-
<i>L. salivarius</i>	S11	+	+	+	+	+
<i>L. salivarius</i>	S12	+	+	-	+	+
<i>L. crispatus</i>	S26	-	-	-	-	-

Five of the isolates selected, three corresponding to *L. plantarum*, and two identified as *L. salivarius*, showed a degree of inhibitory effect against at least one pathogenic strain. S11 and S12, the two *L. salivarius* isolates, showed the broadest spectrum of inhibition, hindering the growth of all pathogenic strains (S11) or of all with the exception of *Salmonella* spp. (S12). In agreement with these results, *L. salivarius* anti-infective ability has already been proved in mice against *Listeria monocytogenes* [30], against *Salmonella enteritidis* infections in chickens (Pascual et al., 1999), hindering of biofilm formation by *C. albicans* in children with caries (krzysciak et al., 2017), in vitro growth inhibition of *Helicobacter pylori* [31], and other bacteria belonging to the genera *Bacillus*, *Enterococcus*, *Staphylococcus*, *Clostridium*, *Pseudomonas*, *Streptococcus*, *Escherichia*, and *Bifidobacterium* [32].




Figure 2: Inhibition of *S. aureus* growth surrounding *L. salivarius* colonies (S11 and S12). No inhibition surrounded *L. reuteri* ATCC ® 23272 control colony (R).

Our results suggest S11 and S12 ability to produce either one type of broad spectrum antimicrobial compound, or several types of compounds that inhibit the growth of the variety of pathogen strains tested, all markedly different: *S. aureus* (Figure 2) and *E. faecium* (Gram positive), *Salmonella* spp. and *Pseudomonas* spp. (Gram negative), in addition to a yeast (*C. albicans*). One possible explanation for S11 and S12 broad spectrum inhibitory activity is production of several types of bacteriocin (selective broad spectrum antimicrobial peptides) [33].

4 Conclusions

For this study, we designed a precise and efficient bioprospecting protocol directed to lactobacilli isolation from poultry feces. Based on four techniques of filtering and identification of colonies (in order: selective medium, Gram stain, MALDI-TOF MS, and 16S rRNA gen sequence BLAST), this protocol allowed the robust identification of 24 *Lactobacillus* isolates and four species (along with other four lactic acid bacteria species) from an initial sample of 114 colonies that grew on MRS selective medium. Thus, with a rate of 21% (24/114) isolation, this protocol highly increases *Lactobacillus* yield compared to its representation of less than 1% in the chicken GM (21-fold).

The protocol ends with a preliminary assessment of a possible probiotic potential of the isolates using antagonism tests, and our results showed positive for five isolates belonging to two *Lactobacillus* species, meaning that these isolates inhibited the growth of at least one pathogen strain tested. From these results a probiotic potential hierarchy can be proposed: with S11 and S12 in the first and second place, respectively (*L. salivarius* isolates); followed by S6 in third place, with S8 and S14 sharing the fourth place (*L. plantarum* isolates); finally, sharing the last place are S31, S107 (*L. brevis* isolates), and S26, *L. crispatus* single isolate. Positive antagonism results could also be increased by performing tests to a larger sample of isolates and pathogen strains.

Given the superior inhibition ability observed for *L. salivarius* isolates, we propose that these bacteria could produce either a wider variety of antimicrobial substances or a single molecule with a broad antimicrobial spectrum, and that this production often reaches the minimal inhibitory concentration (MIC). On the other hand, we suggest several hypothesis that explain that *L. brevis* and *L. crispatus* isolates did not show antagonistic properties: it is possible that both isolates produce antimicrobial compounds but their quantities do not reach the MIC; these isolates lost the genetic potential to produce antimicrobial compounds; antimicrobial compounds are produced but they are not effective against the pathogenic strains tested; growth conditions hinder expression of antimicrobial compounds.

A way to confirm these hypotheses is by directly analyzing antimicro-

bial production by a quantification technique (e.g. HPLC) or indirectly by a transcriptomic analysis directed to the identification of transcripts associated with antimicrobial compounds. Bacteriocin production could be confirmed by quantifying the sensitivity of protease activity, or by a proteomics approach that would allow identification of the different types of bacteriocins produced.

The probiotic potential assessment done in this study is preliminary, however, due to the reduced number of isolates and pathogen strains tested, as well as the constant conditions in which these tests were conducted (pH, salts, and temperature). Furthermore, pathogen inhibition in a Petri dish is just one of the measurable characteristics that imply benefits to the host of the probiotic bacterium. Inhibition of pathogen adhesion to gut epithelia [34], host immune system induction [35], influence on nutrients uptake [36], and resistance of the probiotic bacterium to bile salts and low pH [37], are just some characteristics that make up a high potential probiotic. Then, to achieve a more holistic probiotic potential characterization, it is required to measure several parameters that take into account the mentioned properties *in vitro* and *in vivo*.

Acknowledgements

This study was made possible thanks to the financial support of Universidad EAFIT, and its openness in lending all of its resources and installations. Special thanks to Jaime Robledo and Nora Cardona, both senior medical researches that guided our young curious minds through the hole process of this work. We also have a great gratitude towards the Labmédico Laboratorio Médico de Referencia staff, who trusted two undergraduate students with the utterly expensive MALDI-TOF MS machine and materials, normally used to analyze clinical patients samples. Finally, we would like to thank Carolina Agudelo for checking our final English draft.

References

- [1] H. K. Allen, U. Y. Levine, T. Looft, M. Bandrick, and T. A. Casey, "Treatment, promotion, commotion: antibiotic alternatives in food-producing animals," *Trends in microbiology*, vol. 21, no. 3, pp. 114–119, 2013. [Online]. Available: <https://doi.org/10.1016/j.tim.2012.11.001> 95
- [2] H. C. Wegener, "Antibiotics in animal feed and their role in resistance development," *Current opinion in microbiology*, vol. 6, no. 5, pp. 439–445, 2003. [Online]. Available: <https://doi.org/10.1016/j.mib.2003.09.009> 95
- [3] F. M. Aarestrup, "Occurrence of glycopeptide resistance among enterococcus faecium isolates from conventional and ecological poultry farms," *Microbial Drug Resistance*, vol. 1, no. 3, pp. 255–257, 1995. [Online]. Available: <https://doi.org/10.1089/mdr.1995.1.255> 95
- [4] J. Bates, J. Z. Jordens, and D. T. Griffiths, "Farm animals as a putative reservoir for vancomycin-resistant enterococcal infection in man," *Journal of Antimicrobial Chemotherapy*, vol. 34, no. 4, pp. 507–514, 1994. [Online]. Available: <https://doi.org/10.1093/jac/34.4.507> 95
- [5] I. Klare, H. Heier, H. Claus, G. Böhme, S. Marin, G. Seltmann, R. Hakenbeck, V. Antanassova, and W. Witte, "Enterococcus faecium strains with vana-mediated high-level glycopeptide resistance isolated from animal food-stuffs and fecal samples of humans in the community," *Microbial Drug Resistance*, vol. 1, no. 3, pp. 265–272, 1995. 95
- [6] J. Stephenson, "Antibiotics in animal feed," *JAMA*, vol. 290, no. 11, pp. 1443–1443, 2003. 95
- [7] F. Guarner and G. Schaafsma, "Probiotics." *International journal of food microbiology*, vol. 39, no. 3, pp. 237–238, 1998. 95
- [8] F. Chaucheyras-Durand and H. Durand, "Probiotics in animal nutrition and health," *Beneficial microbes*, vol. 1, no. 1, pp. 3–9, 2009. [Online]. Available: <https://doi.org/10.3920/BM2008.1002> 96
- [9] M. Vanbelle, E. Teller, and M. Focant, "Probiotics in animal nutrition: a review," *Archives of Animal Nutrition*, vol. 40, no. 7, pp. 543–567, 1990. [Online]. Available: <https://doi.org/10.1080/17450399009428406> 96
- [10] P. Banjeree and N. Pradhan, "Live yeasts a good alternative to agp in broiler chickens," *World Poultry*, vol. 22, no. 8, pp. 32–34, 2006. [Online]. Available: https://www.poultryworld.net/PageFiles/27778/001_boerderij-download-WP6926D01.pdf 96

- [11] J. Higgins, S. Higgins, J. Vicente, A. Wolfenden, G. Tellez, and B. Hargis, “Temporal effects of lactic acid bacteria probiotic culture on salmonella in neonatal broilers,” *Poultry Science*, vol. 86, no. 8, pp. 1662–1666, 2007. [Online]. Available: <https://doi.org/10.1093/ps/86.8.1662> 96
- [12] S. Higgins, J. Higgins, A. Wolfenden, S. Henderson, A. Torres-Rodriguez, G. Tellez, and B. Hargis, “Evaluation of a lactobacillus-based probiotic culture for the reduction of salmonella enteritidis in neonatal broiler chicks,” *Poultry Science*, vol. 87, no. 1, pp. 27–31, 2008. [Online]. Available: <https://doi.org/10.3382/ps.2007-00210> 96
- [13] R. La Ragione, A. Narbad, M. Gasson, and M. J. Woodward, “In vivo characterization of lactobacillus johnsonii f9785 for use as a defined competitive exclusion agent against bacterial pathogens in poultry,” *Letters in Applied Microbiology*, vol. 38, no. 3, pp. 197–205, 2004. 96
- [14] R. M. La Ragione and M. J. Woodward, “Competitive exclusion by bacillus subtilis spores of salmonella enterica serotype enteritidis and clostridium perfringens in young chickens,” *Veterinary microbiology*, vol. 94, no. 3, pp. 245–256, 2003. [Online]. Available: [https://doi.org/10.1016/S0378-1135\(03\)00077-4](https://doi.org/10.1016/S0378-1135(03)00077-4) 96
- [15] G. Tellez, C. Pixley, R. Wolfenden, S. Layton, and B. Hargis, “Probiotics/direct fed microbials for salmonella control in poultry,” *Food Research International*, vol. 45, no. 2, pp. 628–633, 2012. [Online]. Available: <https://doi.org/10.1016/j.foodres.2011.03.047> 96
- [16] F. Yan, W. Wang, R. Wolfenden, and H. Cheng, “The effect of bacillus subtilis based probiotic on bone health in broiler chickens,” *Poultry Science*, vol. 95, p. 40, 2016. [Online]. Available: <https://doi.org/10.1093/jas/sky092> 96
- [17] T. Inatomi, “Growth performance, gut mucosal immunity and carcass and intramuscular fat of broilers fed diet containing a combination of three probiotics,” *Sci Postprint*, vol. 1, p. e00052, 2015. 96
- [18] V. Kurtoglu*, F. Kurtoglu, E. Seker, B. Coskun, T. Balevi, and E. Polat, “Effect of probiotic supplementation on laying hen diets on yield performance and serum and egg yolk cholesterol,” *Food additives and contaminants*, vol. 21, no. 9, pp. 817–823, 2004. [Online]. Available: <https://doi.org/10.1080/02652030310001639530> 96
- [19] C. Pineda-Quiroga, R. Atxaerandio, I. Zubiria, I. Gonzalez-Pozuelo, A. Hurtado, R. Ruiz, and A. Garcia-Rodriguez, “Productive performance and cecal microbial counts of floor housed laying hens supplemented with

- dry whey powder alone or combined with *pediococcus acidilactici* in the late phase of production,” *Livestock Science*, vol. 195, pp. 9–12, 2017. [Online]. Available: <https://doi.org/10.1016/j.livsci.2016.11.007> 96
- [20] M. Yörük, M. Gül, A. Hayirli, and M. Macit, “The effects of supplementation of humate and probiotic on egg production and quality parameters during the late laying period in hens,” *Poultry Science*, vol. 83, no. 1, pp. 84–88, 2004. [Online]. Available: <https://doi.org/10.1093/ps/83.1.84> 96
- [21] R. Fuller, “Ecological studies on the lactobacillus flora associated with the crop epithelium of the fowl,” *Journal of Applied Bacteriology*, vol. 36, no. 1, pp. 131–139, 1973. [Online]. Available: <https://doi.org/10.1111/j.1365-2672.1973.tb04080.x> 97
- [22] J. Patterson and K. Burkholder, “Application of prebiotics and probiotics in poultry production,” *Poultry science*, vol. 82, no. 4, pp. 627–631, 2003. [Online]. Available: <https://doi.org/10.1093/ps/82.4.627> 97
- [23] B. B. Oakley, H. S. Lillehoj, M. H. Kogut, W. K. Kim, J. J. Maurer, A. Pedroso, M. D. Lee, S. R. Collett, T. J. Johnson, and N. A. Cox, “The chicken gastrointestinal microbiome,” *FEMS microbiology letters*, vol. 360, no. 2, pp. 100–112, 2014. [Online]. Available: [10.1111/1574-6968.12608](https://doi.org/10.1111/1574-6968.12608) 97, 100
- [24] P. J. Turnbaugh, V. K. Ridaura, J. J. Faith, F. E. Rey, R. Knight, and J. I. Gordon, “The effect of diet on the human gut microbiome: a metagenomic analysis in humanized gnotobiotic mice,” *Science translational medicine*, vol. 1, no. 6, pp. 6ra14–6ra14, 2009. [Online]. Available: <https://doi.org/10.1126/scitranslmed.3000322> 97
- [25] E. Carbonnelle, C. Mesquita, E. Bille, N. Day, B. Dauphin, J.-L. Beretti, A. Ferroni, L. Gutmann, and X. Nassif, “Maldi-tof mass spectrometry tools for bacterial identification in clinical microbiology laboratory,” *Clinical biochemistry*, vol. 44, no. 1, pp. 104–109, 2011. [Online]. Available: <https://doi.org/10.1016/j.clinbiochem.2010.06.017> 97
- [26] J. M. Janda and S. L. Abbott, “16s rRNA gene sequencing for bacterial identification in the diagnostic laboratory: pluses, perils, and pitfalls,” *Journal of clinical microbiology*, vol. 45, no. 9, pp. 2761–2764, 2007. [Online]. Available: <https://doi.org/10.1128/JCM.01228-07> 97
- [27] U. Schillinger and F. K. Lücke, *Applied and environmental microbiology*, vol. 55, no. 8, pp. 1901–1906, 1989. 98
- [28] T. A. Tatusova and T. L. Madden, “Blast 2 sequences, a new tool for comparing protein and nucleotide sequences,” *FEMS microbiology*

- letters, vol. 174, no. 2, pp. 247–250, 1999. [Online]. Available: [https://doi.org/10.1016/S0378-1097\(99\)00149-4](https://doi.org/10.1016/S0378-1097(99)00149-4) 99
- [29] K. Sonomoto and A. Yokota, *Lactic acid bacteria and bifidobacteria: current progress in advanced research*. Horizon Scientific Press, 2011. 100
- [30] S. C. Corr, Y. Li, C. U. Riedel, P. W. O’Toole, C. Hill, and C. G. Gahan, “Bacteriocin production as a mechanism for the antiinfective activity of lactobacillus salivarius ucc118,” *Proceedings of the National Academy of Sciences*, vol. 104, no. 18, pp. 7617–7621, 2007. [Online]. Available: <https://doi.org/10.1073/pnas.0700440104> 103
- [31] K. A. Ryan, P. Daly, Y. Li, C. Hooton, and P. W. O’Toole, “Strain-specific inhibition of helicobacter pylori by lactobacillus salivarius and other lactobacilli,” *Journal of Antimicrobial Chemotherapy*, vol. 61, no. 4, pp. 831–834, 2008. [Online]. Available: <https://doi.org/10.1093/jac/dkn040> 103
- [32] C. Dunne, L. Murphy, S. Flynn, L. O’Mahony, S. O’Halloran, M. Feeney, D. Morrissey, G. Thornton, G. Fitzgerald, C. Daly *et al.*, “Probiotics: from myth to reality. demonstration of functionality in animal models of disease and in human clinical trials,” in *Lactic Acid Bacteria: Genetics, Metabolism and Applications*. Springer, 1999, pp. 279–292. 103
- [33] S. Messaoudi, M. Manai, G. Kergourlay, H. Prévost, N. Connil, J.-M. Chobert, and X. Dousset, “Lactobacillus salivarius: bacteriocin and probiotic activity,” *Food microbiology*, vol. 36, no. 2, pp. 296–304, 2013. [Online]. Available: <https://doi.org/10.1016/j.fm.2013.05.010> 103
- [34] M. Velraeds, H. Van der Mei, G. Reid, and H. J. Busscher, “Inhibition of initial adhesion of uropathogenic enterococcus faecalis by biosurfactants from lactobacillus isolates.” *Applied and environmental microbiology*, vol. 62, no. 6, pp. 1958–1963, 1996. [Online]. Available: [https://doi.org/10.1016/S0090-4295\(97\)00065-4](https://doi.org/10.1016/S0090-4295(97)00065-4) 105
- [35] C. M. Galdeano and G. Perdigon, “The probiotic bacterium lactobacillus casei induces activation of the gut mucosal immune system through innate immunity,” *Clinical and Vaccine Immunology*, vol. 13, no. 2, pp. 219–226, 2006. [Online]. Available: <https://doi.org/10.1128/CVI.13.2.219-226.2006> 105
- [36] G. Storelli, A. Defaye, B. Erkosar, P. Hols, J. Royet, and F. Leulier, “Lactobacillus plantarum promotes drosophila systemic growth by modulating hormonal signals through tor-dependent nutrient sensing,” *Cell metabolism*, vol. 14, no. 3, pp. 403–414, 2011. [Online]. Available: <https://doi.org/10.1016/j.cmet.2011.07.012> 105

- [37] F. L. Tulini, L. K. Winkelströter, and E. C. De Martinis, "Identification and evaluation of the probiotic potential of *Lactobacillus paraplantarum* ft259, a bacteriocinogenic strain isolated from brazilian semi-hard artisanal cheese," *Anaerobe*, vol. 22, pp. 57–63, 2013. [Online]. Available: <https://doi.org/10.1016/j.anaerobe.2013.06.006> 105

Supplementary data




Figure S1: Antagonism tests diagram. A: atop view of a Petri dish with inhibition zones surrounding *Lactobacillus* colonies. B: side view of a Petri dish with semi-solid LB medium on top and solid MRS medium at the bottom.