

# Elimination of Yaws in Colombia: An Epidemiological Milestone Within Reach

## Eliminación del pian en Colombia: un hito epidemiológico al alcance

Alfonso J. Rodríguez-Morales<sup>1,\*</sup>, Juan Carlos Sepúlveda-Arias<sup>2</sup>, Wilmer E. Villamil-Gomez<sup>3</sup>

**Keywords:** frambesia, Yaws, *Treponema pallidum*, Colombia

**Palabras clave:** frambesia, Pian, *Treponema pallidum*, Colombia

Yaws, also known as frambesia or pian, is a chronic, non-venereal treponematoses caused by *Treponema pallidum* subspecies *pertenue*<sup>1,2</sup>. It is transmitted through direct skin-to-skin contact with infectious lesions and primarily affects children living in impoverished, rural, and humid tropical settings<sup>3</sup>. The disease evolves through three stages. The primary stage presents as a painless papule ("mother yaw") that may ulcerate<sup>4</sup>. The secondary stage is characterized by multiple disseminated papillomatous or ulcerative lesions, often accompanied by periostitis<sup>5</sup>. If untreated, patients may enter a latent phase, and approximately 10% can progress to tertiary disease, with destructive lesions of skin, bones, and cartilage that may lead to permanent disability and social stigma<sup>6,7</sup>.

Diagnosis relies on clinical suspicion supported by serological testing, combining non-treponemal tests (VDRL or RPR) with treponemal tests (TPHA or FTA-ABS). The differential diagnosis for yaws lesions is broad and includes syphilis, leishmaniasis, leprosy, and Buruli ulcer, as well as non-infectious causes. Because serological assays cannot differentiate yaws from other treponematoses, the epidemiological context is essential for interpretation<sup>8</sup>. Treatment is simple and highly effective: a single oral dose of azithromycin is currently recommended, with benzathine penicillin as an alternative.

Prevention depends on early case detection, contact tracing and treatment, sustained surveillance, and improvements in social determinants such as hygiene, housing conditions, and access to health services<sup>9</sup>.

Historically, yaws was widespread across tropical regions worldwide<sup>10</sup>. In the mid-20th century, tens of millions of people were estimated to be infected globally<sup>11</sup>. Large-scale mass treatment campaigns with penicillin, coordinated internationally between the 1950s and 1960s, led to dramatic reductions in prevalence<sup>12</sup>. However, incomplete surveillance and waning political commitment allowed the disease to persist in pockets of poverty in several regions. Yaws was introduced into the Americas during the colonial period and became established in vulnerable communities<sup>13,14</sup>. The first case of the disease arrived in the Caribbean from Africa in 1502, and its incidence increased after 1512, with the spread of slavery and the beginning of sugarcane cultivation. In Colombia, by the early 20th century, it was recognized as an endemic disease, particularly affecting rural and coastal populations<sup>15</sup>. By the early 1930s, authorities estimated approximately 70,000 active cases nationwide, with the highest burden concentrated along the Pacific coast, but also present in other areas of the country (e.g., Antioquia, Valle del Cauca)<sup>16-23</sup>.

The introduction of penicillin marked a turning point in Colombia's epidemiological history of yaws. In 1950, the country launched a national anti-yaws campaign based on acti-

1 Faculty of Health Sciences, Universidad Científica del Sur, Lima, Peru. Grupo de Investigación Biomedicina, Faculty of Medicine, Fundación Universitaria Autónoma de las Américas-Institución Universitaria Visión de las Américas, Pereira 660007, Colombia. Associate Editor, Infectio. <https://orcid.org/0000-0001-9773-2192>

2 Grupo Infección e Inmunidad, Facultad de Ciencias de la Salud, Universidad Tecnológica de Pereira, Pereira 660003, Colombia. Editor-in-Chief, Infectio. <https://orcid.org/0000-0002-5410-5997>

3 Centro de Investigación en Ciencias de la Vida, Universidad Simón Bolívar, Barranquilla, Colombia. Comité de Medicina Tropical, Zoonosis y Medicina del Viajero, Asociación Colombiana de Infectología (ACIN). Comité de Enfermedades Tropicales, Asociación Panamericana de Infectología (API). <https://orcid.org/0000-0002-5492-3455>

\* Autor para correspondencia:  
Correo electrónico: [arodriguezmo@cientifica.edu.pe](mailto:arodriguezmo@cientifica.edu.pe)

Recibido: 18/02/2026; Aceptado: 18/02/2026

Cómo citar este artículo: A.J. Rodríguez-Morales, *et al.* Elimination of Yaws in Colombia: An Epidemiological Milestone Within Reach. Infectio 2026; 30(2): 111-114

ve case finding and house-to-house treatment. Over three years, more than 111,000 individuals with active yaws and over 125,000 of their contacts received treatment. The impact was immediate and profound. Reported cases fell dramatically from more than 50,000 in 1950 to fewer than 6,000 by 1960. By 1973, only a few hundred cases were reported in endemic areas, and by 1983, the official annual report documented just 31 cases nationwide. This decline represents one of the earliest and most successful public health interventions against a neglected tropical disease in Colombia. However, as transmission decreased, surveillance activities were gradually reduced. Routine notification eventually ceased, and the absence of reported cases was not accompanied by formal verification of a transmission interruption<sup>15,20,22</sup>.


The epidemiological trajectory of yaws in Colombia reflects a classic pattern: hyperendemicity in the early 20th century, rapid decline following mass treatment, and subsequent epidemiological silence (Figure 1). While the near disappearance of reported cases strongly suggests that transmission was interrupted, designating elimination as a public health success does not automatically imply that it is a verified epidemiological status. Today, Colombia is considered to be in the elimination phase (<https://www.minsalud.gov.co/salud/publica/PET/Paginas/pian.aspx>). The country no longer reports routine cases of yaws, yet a formal demonstration of interruption of transmission remains necessary to achieve certification. This requires structured surveillance activities capable of detecting any residual or re-emergent transmission<sup>2,24</sup>.

The primary epidemiological challenge lies in distinguishing true absence of disease from absence of detection (<https://www.minsalud.gov.co/salud/publica/PET/Paginas/pian.aspx>). Yaws predominantly affects children under 15 years of age, especially those living in remote, socioeconomically disad-


vantaged settings. Consequently, surveillance efforts must focus on historically endemic territories and vulnerable populations. Colombia has recently reinforced its institutional commitment by adopting a comprehensive national plan for the elimination and sustained elimination of communicable diseases and priority conditions for the period 2025–2031 ([https://www.minsalud.gov.co/Normatividad\\_Nuevo/Resolucion%20No%20119%20de%202026.pdf](https://www.minsalud.gov.co/Normatividad_Nuevo/Resolucion%20No%20119%20de%202026.pdf)). This framework aligns with regional and global initiatives to eliminate multiple communicable diseases by 2030.

The national plan establishes the elimination of transmission and sustained elimination as explicit objectives (<https://www.minsalud.gov.co/sites/rid/Lists/BibliotecaDigital/RIDE/VS/PP/ET/plan-nacional-eliminacion-transmision-pian-2025-2031.pdf>). It emphasizes integration into primary health care, intersectoral coordination, territorial governance, community participation, and systematic monitoring and evaluation (Figures 2 and 3). These elements are particularly relevant for diseases such as yaws, which are deeply linked to poverty, environmental conditions, and health system access. Under international guidelines, verification of yaws elimination requires evidence of zero indigenous cases over a defined period, supported by sensitive and reliable surveillance systems (Figure 3). Annual surveys of children in previously endemic areas are recommended during the post-zero-case phase, as children are the most sensitive indicator group for ongoing transmission ([https://www.minsalud.gov.co/Normatividad\\_Nuevo/Resolucion%20No%20119%20de%202026.pdf](https://www.minsalud.gov.co/Normatividad_Nuevo/Resolucion%20No%20119%20de%202026.pdf)).

Achieving certification of elimination depends fundamentally on epidemiological vigilance. Strengthened surveillance must include training healthcare workers to recognize compatible skin lesions, ensuring the availability of appropriate serological testing, and promoting accurate case registration within national health information systems (Figure 3). Active



**Figure 1.** Historical situation of yaws in Colombia. Based on data from the Ministry of Health of Colombia (<https://www.minsalud.gov.co/salud/publica/PET/Paginas/pian.aspx>).




**Figure 2.** Infographics of the Ministry of Health of Colombia about the importance of yaws in the country. (Source: <https://www.minsalud.gov.co/sites/rid/Lists/BibliotecaDigital/RIDE/VS/PP/ET/importancia-pian-salud-publica.pdf>)

case finding in historically affected municipalities is essential, particularly in remote rural and Pacific coastal regions where transmission was once concentrated. Additionally, integrating yaws surveillance into broader neglected tropical disease platforms can enhance efficiency and sustainability. Colombia's ongoing investments in primary health care, territorial management, and community-based strategies provide an operational foundation for these activities<sup>6</sup>.

Accurate documentation is critical. Even isolated suspected cases must be thoroughly investigated, with laboratory confirmation and epidemiological assessment. The absence of confirmed cases, supported by systematic surveillance data, will constitute the evidence base required for formal certification. From an epidemiological perspective, elimination of yaws in Colombia appears highly plausible. The extraordinary reduction from tens of thousands of cases in the mid-20th century to near-zero reports by the 1980s strongly suggests that sustained transmission was likely interrupted decades ago. Nevertheless, certification demands rigorous demonstration<sup>2,25</sup>.

Beyond the technical process, verified elimination would represent a symbolic and substantive achievement. It would confirm that a disease historically rooted in poverty, exclusion, and limited access to health services can be effectively controlled and ultimately eliminated through coordinated public health action. Moreover, Colombia's success would contribute to the broader regional and global goal of yaws eradication. Unlike many infectious diseases, yaws has no known animal reservoir, no vector, and a highly effective single-dose treatment. These characteristics make eradication biologically feasible, provided surveillance and response systems remain vigilant. The elimination of yaws in Colombia is therefore not merely the conclusion of a historical campaign; it is a test of the country's capacity to sustain epidemiological rigor in the absence of visible disease. Through strengthened surveillance, targeted verification activities, and continued commitment to equity in health, Colombia stands on the threshold of formally consigning yaws to history. The final step is clear: document the absence, verify the interruption, and sustain the achievement<sup>13,26</sup>.



**Figure 3.** Infographics from Colombia's Ministry of Health on the appropriate recording/notification of yaws cases in the country. (Source: <https://www.minsalud.gov.co/sites/rid/Lists/BibliotecaDigital/RIDE/VS/PP/ENT/registro-casos-pian-colombia.pdf>)

## References

- Mitjà O, Marks M, Konan DJ, et al. Global epidemiology of yaws: a systematic review. *Lancet Glob Health* 2015;3:e324-31.
- Zoni AC, Saboyá-Díaz MI, Castellanos LG, Nicholls RS, Blaya-Novakova V. Epidemiological situation of yaws in the Americas: A systematic review in the context of a regional elimination goal. *PLoS Negl Trop Dis* 2019;13:e0007125.
- Barton A, Pospíšilová P, Beiras CG, et al. Effect of repeated mass drug administration on the transmission of yaws: a retrospective genomic epidemiology study. *Lancet Microbe* 2026;7:101229.
- Marks M, Yin YP, Chen XS, et al. Metaanalysis of the Performance of a Combined Treponemal and Nontreponemal Rapid Diagnostic Test for Syphilis and Yaws. *Clin Infect Dis* 2016;63:627-33.
- Vicar EK, Simpson SV, Mensah GI, Addo KK, Donkor ES. Yaws in Africa: Past, Present and Future. *Diseases* 2025;13.
- Marks M. Advances in the Treatment of Yaws. *Trop Med Infect Dis* 2018;3.
- Román GC, Román LN. Occurrence of congenital, cardiovascular, visceral, neurologic, and neuro-ophthalmologic complications in late yaws: a theme for future research. *Rev Infect Dis* 1986;8:760-70.
- Giacani L, Lukehart SA. The endemic treponematoses. *Clin Microbiol Rev* 2014;27:89-115.
- Mitjà O, Houine W, Moses P, et al. Mass treatment with single-dose azithromycin for yaws. *N Engl J Med* 2015;372:703-10.
- Mitjà O, Asiedu K, Mabey D. Yaws. *Lancet* 2013;381:763-73.
- Rinaldi A. Yaws eradication: facing old problems, raising new hopes. *PLoS Negl Trop Dis* 2012;6:e1837.
- Asiedu K, Fitzpatrick C, Jannin J. Eradication of yaws: historical efforts and achieving WHO's 2020 target. *PLoS Negl Trop Dis* 2014;8:e3016.
- Marks M, Mitjà O, Vestergaard LS, et al. Challenges and key research questions for yaws eradication. *Lancet Infect Dis* 2015;15:1220-5.
- Guderian RH, Anselmi M, Calvopiña M, Cooper PJ, Mancero T. El pian en la provincia de Esmeraldas, Ecuador. *Biomédica* 1995;15:137-43.
- Vélez Tobón GJ, Estrada Orrego V. Deformaciones y contagios. La lucha contra el pian en Colombia en la primera mitad del siglo XX. *Trashumante Revista Americana de Historia Social* 2021;0:28-51.
- Lopez Narvaez G. [Yaws and the anti-yaws campaign in Colombia]. *Rev Fac Med Univ Nac Colomb* 1956;24:294-342.
- Hopkins DR, Florez D. Pinta, yaws, and venereal syphilis in Colombia. *Int J Epidemiol* 1977;6:349-55.
- Uribe WR. Yaws in Colombia. *Rev Infect Dis* 1985;7 Suppl 2:S276-7.
- Hopkins DR, Florez D. Comparison of RPR 'teardrop' card test, VDRL and FTA-ABS tests results on sera from persons with suspected yaws in Colombia. *Br J Vener Dis* 1977;53:218-20.
- VÉLEZ TOBÓN GJ, ESTRADA ORREGO V. La medicina social en zonas rurales: una historia de la lucha contra el pian en Antioquia, 1918-1941. *Anuario Colombiano de Historia Social y de la Cultura* 2022;49:97-126.
- Guerra A G. El pian en el Litoral Pacífico Colombiano: Cómo debe orientarse su campaña. *Oficina Sanitaria Panamericana* 1941;20:895-7.
- Restrepo M, Restrepo BN, Orozco B, Padilla JC, Cuervo CM, Alvarez G. Encuesta seroepidemiológica de pian en la Costa Pacífica colombiana. *Biomédica* 2001;21:155-61.
- Vargas Cuéllar PI. El pian en el Departamento del Valle del Cauca, Colombia. *Oficina Sanitaria Panamericana* 1941;20:897-914.
- Castañeda-Hernández DM, Rodríguez-Morales AJ. Panorama de las enfermedades transmisibles en Colombia desde la perspectiva del Plan Decenal de Salud Pública 2012-2021. *Infectio* 2015;19:141-3.
- Dofitas BL, Kalim SP, Toledo CB, Richardus JH. Yaws in the Philippines: first reported cases since the 1970s. *Infect Dis Poverty* 2020;9:1.
- Fitzpatrick C, Asiedu K, Solomon AW, et al. Prioritizing surveillance activities for certification of yaws eradication based on a review and model of historical case reporting. *PLoS Negl Trop Dis* 2018;12:e0006953.