

The relationship between atmospheric condition and human mortality associated with coarse material particulate in Bogotá (Colombia)

Relación entre condición atmosférica y mortalidad humana asociada con material particulado grueso en Bogotá (Colombia)

Relação entre condição atmosférica e mortalidade humana associada com material Particulado grosso em Bogotá (Colômbia)

Carlos Alfonso Zafra-Mejía^{a*} | Juan Pablo Rodríguez-Miranda^b | Hugo Alexander Rondón-Quintana^c

^a <https://orcid.org/0000-0002-4061-4897>

^b <https://orcid.org/0000-0002-3761-8221>

^c <https://orcid.org/0000-0003-2946-9411>

Universidad Distrital Francisco José de Caldas, Bogotá, Colombia

- **Fecha de recepción:** 2020-05-12
- **Fecha concepto de evaluación:** 2020-07-11
- **Fecha de aprobación:** 2020-07-14
<https://doi.org/10.22335/rlct.v12i3.1237>

Para citar este artículo / To reference this article / Para citar este artigo: Zafra-Mejía, C.A., Rodríguez-Miranda, J.P., & Rondón-Quintana, H.A. (2020). The relationship between atmospheric condition and human mortality associated with coarse particulate matter in Bogotá (Colombia). *Revista Logos Ciencia & Tecnología*, 12(3), 57-68. <https://doi.org/10.22335/rlct.v12i3.1237>

ABSTRACT

This article studies the relationship between atmospheric condition (AC) and human mortality rate associated with coarse particulate matter (PM₁₀) in a high-altitude mega-city (Bogotá-Colombia). Information was collected from three automatic monitoring stations equipped with measuring instruments for PM₁₀, temperature, solar radiation and wind speed. The sampling period lasted six years. The results showed the best possible scenario for the maximum hourly concentrations of PM₁₀ (52.3-135 µg/m³). These events occurred during daytime periods where the predominant AC was between unstable and very unstable. The risk from exposure to PM₁₀ showed that February>March>January were the highest risks. These months showed PM₁₀ concentrations 35.9% higher than those observed during the months of lower risk (August>June>July). A higher mortality rate (+2.0%) was suggested in urban sectors with less atmospheric instability (AI) and predominance of impervious cover compared to sectors with higher AI and predominance of vegetated cover.

Keywords: Bogotá, air pollution, atmospheric stability, megacity, public health, PM₁₀



RESUMEN

Este artículo estudia la relación entre condición atmosférica (CA) y tasa de mortalidad humana asociada con el material particulado grueso (PM_{10}) en una mega-ciudad de elevada altitud (Bogotá-Colombia). Se recolectó información de tres estaciones automáticas de monitoreo equipadas con instrumentos de medición para PM_{10} , temperatura, radiación solar y velocidad del viento. El periodo de muestreo duró seis años. Los resultados mostraron el mejor escenario posible para las máximas concentraciones horarias de PM_{10} (52.3-135 $\mu\text{g}/\text{m}^3$). Estos eventos ocurrieron durante periodos diurnos donde la CA predominante estuvo entre inestable-muy inestable. El riesgo por exposición a PM_{10} mostró que febrero>marzo>enero fueron los de mayor riesgo. Estos meses mostraron concentraciones de PM_{10} un 35.9% más elevadas que aquellas observadas durante los meses de menor riesgo (agosto>junio>julio). Se sugirió una mayor tasa de mortalidad (+2,0%) en sectores urbanos con menor inestabilidad atmosférica (IA) y predominancia de cobertura impermeable, en comparación con sectores de mayor IA y predominancia de cobertura vegetada.

Palabras clave: Bogotá, contaminación atmosférica, estabilidad atmosférica, megaciudad, salud pública, PM_{10}

RESUMO

Este artigo estuda a relação entre condição atmosférica (CA) e taxa de mortalidade humana associada com o material particulado grosso (PM_{10}) em uma megacidade de elevada altitude (Bogotá-Colômbia). Coletou-se informação de três estações automáticas de monitoramento equipadas com instrumentos de medição para PM_{10} , temperatura, radiação solar e velocidade do vento. O período de amostragem teve uma duração de seis anos. Os resultados revelaram o melhor cenário possível para as máximas concentrações horárias de PM_{10} (52.3-135 $\mu\text{g}/\text{m}^3$). Estes eventos ocorreram durante períodos diurnos onde a CA predominante esteve entre instável e muito instável. O risco por exposição a PM_{10} mostrou que fevereiro>março>janeiro foram os de maior risco. Estes meses mostraram concentrações de PM_{10} um 35.9% mais elevadas que aquelas observadas durante os meses de menor risco (agosto>junho>julho). Sugeriu-se uma maior taxa de mortalidade (+2,0%) em setores urbanos com menor instabilidade atmosférica (IA) e predominância de cobertura impermeável, em comparação com setores de maior IA e predominância de cobertura vegetada.

Palavras-chave: Bogotá, contaminação atmosférica, estabilidade atmosférica, megacidade, saúde pública, PM_{10}

Atmospheric aerosols are of global importance because they affect the climate via direct and indirect radiative forcing and adversely impact the human health and ecosystems (Dahari, Latif, Muda & Hussein, 2020). Air pollution continues to be a growing problem for human health in terms of short and long-term effects and is a burden on public health systems around the world (Maesano et al., 2020; Yang et al., 2020). One of the main pollutants that contributes to the deterioration of air quality is particulate matter (PM). The increase in the urban PM is mainly associated with the concentration of industrial activities and the increase in the number of motor vehicles (Terrouche et al., 2016; Zafra-Mejía, Gutiérrez-Malaxechebarria & Hernández-Peña, 2019). Because of this,

many Latin American countries have decided to strengthen the active monitoring of air pollution in urban areas in order to control atmospheric pollutants (Palacio, Zafra & Rodríguez, 2014; Franck, Leitte & Suppan, 2014). The impacts of PM on human health are associated with a reduction in cardiopulmonary functions and with an increase in mortality from cardiovascular disease, the occurrence of asthma in children, and the risk of cancer (Hou, An, Tao & Sun, 2016; Maesano et al., 2020).

Dockery et al. (1993) reported that diseases such as bronchitis and chronic asthma were positively correlated with PM concentrations. In a European study conducted in 15 western European cities on air pollution and its effects on

public health, it was reported that an increase of $50 \mu\text{g}/\text{m}^3$ in PM_{10} concentrations could cause a 2.1% increase in the human mortality rate (Katsouyanni et al., 1996). Another study conducted with the support of the United States Environmental Protection Agency (US EPA) on the effects of air pollution on the respiratory health of people in Chinese mega-cities (Guangzhou, Wuhan, and Chongqing), it was observed that the PM_{10} concentration was directly related to the rate of childhood lung dysfunction (Liu, Xu & Yang, 2018). Lacasaña et al. (1999) and Hernández et al. (2013) reported that the effects on human health from extreme events of air pollution were short-term. This study reported latency periods of up to seven days.

Studies reported that the AC plays an important role in the transportation and dispersion of PM_{10} , and is significantly related to temperature variation at altitude (thermal gradient) and wind speed (Srinivas, Venkatesan, Somayaji & Indira, 2009), with this last variable also depending on land surface characteristics (Chambers et al., 2015; Zafra, Ángel & Torres, 2017). Related to this, Lee, Ho, Lee, Choi and Song, (2013) reported that in Seoul (Korea), under extreme atmospheric stability conditions (thermal inversion) the PM_{10} concentrations tended to increase significantly ($\text{PM}_{10} > 100 \mu\text{g}/\text{m}^3$). Vecchi, Marcazzan and Valli, (2007) reported a 13% increase in PM_{10} concentrations in Milan (Italy) under prevailing conditions of nighttime atmospheric stability (low dispersion), despite a reduction in the emission sources (traffic, domestic heating, and industries). This study also reported that, under conditions of daytime atmospheric instability (high dispersion), the PM_{10} concentrations tended to decrease.

As can be seen, it is relevant that urban public health control agencies study the possible relationship between PM concentrations and the atmospheric condition (AC) that prevails in a given area. In this way, it is also relevant to study the possible relationship between AC and the human mortality rate associated with PM concentrations.

This study was conducted in the city of Bogotá, Colombia, a high-altitude Latin American mega-city with a population of 8.85 million inhabitants in 2015. It is located in a large inter-mountain valley in the eastern Andes ($04^{\circ}36'35''\text{N}$ - $74^{\circ}04'54''\text{W}$) at an average altitude of 2600 masl. Its tropical mountain climate is characterized by large temperature variations per hour (maximum variation = 12°C). The city is recognized as the urban centre with the highest population density (26,000 inhabitants/ km^2) and the third-highest air pollution level in

Latin America (Sarmiento, Hernández, Medina, Rodríguez, & Reyes, 2015). According to Montoya, Cepeda and Eslava, (2004), atmospheric instability conditions during the day are recorded in the city due to the increase of solar radiation until noon and afternoon. During the night, the wind speed is very low, which produces stable and neutral atmospheric conditions. Therefore, due to the interaction between the degree of pollution and the particular climatic characteristics of the city, the development of this study is of interest when it comes to understanding the behaviour of PM_{10} and the associated mortality rate according to atmospheric conditions.

This article aims to study the relationship between AC and the human mortality rate associated with PM_{10} concentrations in this high-altitude megacity of Bogotá (Colombia). The study was developed from the available PM_{10} information per hour between the years 2007-2012 from three automatic monitoring stations located throughout the city. This study hopes to increase knowledge about: (1) the behaviour of PM_{10} contamination in high-altitude megacities in developing countries; (2) the possible relationship between AC and the daily mortality rate associated with PM_{10} in megacities.

■ Materials and methods

Data collection

The three automatic monitoring stations were located in Kennedy (S1), Barrios Unidos (S2) and Guaymaral (S3) in Bogotá, Colombia. These stations were selected because they showed the maximum (S1) and minimum (S2 and S3) concentrations of PM_{10} during the study period. Additionally, the selected stations did not have any significant changes in land surface coverage, but they had information about climatic variables allowing for a determination of the AC. The tropical mountain climate showed an average daily temperature of between $13.3 - 14.3^{\circ}\text{C}$ and wide temperature variations per hour with variations between $7.2 - 19^{\circ}\text{C}$. Table I shows the main characteristics of each monitoring station. The locations of the monitoring stations are shown in Figure 1.

PM_{10} sampling system

The sampling period lasted for six years (01/01/2007-12/31/2012). This time interval was selected because the areas surrounding the monitoring stations showed no significant changes in land surface coverage (impervious, vegetated, non-vegetated, and water-bodies; see Table I).

Table 1. Characteristics of the location zones of each monitoring station

Characteristics	S1 (Kennedy)	S2 (Barrios Unidos)	S3 (Guaymaral)
Coordinates	4°37'30.18"N 74°9'40.80"W	4°39'30.48"N 74°5'2.28"W	4°47'1.52"N 74°2'39.06"W
Altitude (masl)	2580	2577	2580
Average daily PM ₁₀ (µg/m ³) ^a	85.9	40.0	34.9
Average annual rainfall (mm) ^a	521	1084	832
Average daily wind speed ^b	2.2	1.35	1.0
Prevailing wind direction ^a	SW	W	SE
Average daily temperature (°C) ^a	14.3	14.3	14.2
Type of zone	Urban	Urban	Suburban
Land use ^b	R-I-C	R-IN-I	R-IN
Land surface coverage: Impervious/Vegetated/Non-vegetated/Water-bodies (%) ^c	85.8/10.9/2.90/0.40	56.5/39.3/0.50/3.70	17.9/78.4/2.83/0.87
PM ₁₀ sampler location (m) ^d	7.0	4.6	4.0
Type of monitoring station	Background	Background	Background
Population density (Inhabitants/ha)	400	30	< 1

Note: ^a During the study period; ^b R = Residential, I = Industrial, C = Commercial, and IN = Institutional; ^c Within a 1600-m radius with center in the monitoring station; ^d With respect to the land surface.



Figure 1. Location of monitoring stations in Bogotá, Colombia (Google Maps, 2017). S1 = Kennedy, S2 = Barrios Unidos, and S3 = Guaymaral

Thus, during the sampling period, the possible effects of the variation of land surface coverage in the AC (vertical gradient of temperature and wind speed) at each sampling

site was minimized. The hourly PM₁₀ sampling system consisted of continuous-monitoring equipment which used beta ray attenuation (Met One Instruments, BAM 1020). The sampling protocol followed the guidelines set forth by the United States Environmental Protection Agency in EPA/625/R-96/010a-IO-1.2 (U.S. EPA, 1999). The equipment's constant flow rate was 16.7 l/min. The lowest detection limit was 3.6 µg/m³ and 1.0 µg/m³ for hourly and daily sampling intervals, respectively. Resolution in the measurement was 0.24 µg in a range of 1 mg. The precision was ± 8% for hourly intervals and ± 2% for daily intervals.

Atmospheric condition

For each monitoring station, the AC was determined based on the Pasquill (1961), Turner (1964), and Gifford (1976) methodologies, using hourly data on wind speed and solar radiation. The prevalence of the AC was determined following the methodologies proposed by Zoras, Triantafyllou and Deligiorgi, (2006) and Chambers et al. (2015) according to their hourly frequency. Differences in hourly frequency between monitoring stations were evaluated by applying the Kruskal-Wallis test, which is a non-parametric alternative test to a one-way ANOVA. The CA data distribution was validated with the Shapiro-Wilk test, identifying a non-normal distribution ($df = 24$; p values < 0.001). The CA states were defined on

a quantitative scale according to the categories: 1 = stable, 2 = slightly stable, 3 = neutral, 3.5 = neutral to slightly unstable, 4 = slightly unstable, 4.5 = slightly unstable to unstable, 5 = unstable, 5.5 = unstable to very unstable and 6 = very unstable. Finally, a graphical comparison between PM_{10} concentrations and the AC was presented for each monitoring station.

Air quality standards

The standards selected to assess air quality in the study areas were Colombian Resolution No. 610/2010 (MAVDT, 2010) and air quality guidelines of the World Health Organization (WHO, 2005). These standards established that the maximum permissible levels allowed for PM_{10} are 100 and 50 $\mu\text{g}/\text{m}^3$ for an exposure time of 24 hours, respectively. The frequency of exceedance of these maximum permissible levels during the entire study period was analysed for daily and monthly time scales.

Following the daily concentrations of PM_{10} and the reference legislation, quartiles (Q) were established in order to assess the risk to public health. Average daily PM_{10} concentrations were calculated by month and later organized in order of precedence. The three months that exhibited the highest PM_{10} concentration (greatest public health risk) were assigned to the first quartile (Q1). The quartiles Q2, Q3 and Q4 were assigned according to PM_{10} concentration in order of precedence. Thus, the three months with the lowest average PM_{10} concentration were assigned to the last quartile (Q4), representing the lowest risk group for public health.

The increase in the human mortality rate related to PM_{10} for the study areas was calculated. This calculation was based on the guidelines established by WHO (WHO, 2005). The results of studies at a global level suggested that the public health risks related to short-term exposures to PM_{10} were probably similar in cities in developed and developing countries, with an increase in the daily mortality rate around 0.50% for each 10 $\mu\text{g}/\text{m}^3$ increase in PM_{10} (WHO, 2005). Thus, this study assumed an increase in the daily mortality rate of 0.50%. These increases in the daily mortality rate were calculated from the maximum 24-h limit established by WHO for PM_{10} (50 $\mu\text{g}/\text{m}^3$).

PM_{10} data analysis

For all monitoring stations, the hourly variation of PM_{10} concentration was evaluated for the entire study period ($n = 24$, per monitoring station). The daily variation of PM_{10} concentrations was evaluated using a 24-hour

moving average. The non-normal distribution of PM_{10} data was determined with the Shapiro-Wilk test (p -values < 0.048). Spearman's correlation coefficient (r_s) analysis was carried out to assess the relationship between monitoring stations according to the hourly and daily PM_{10} concentration. Additionally, to assess the hourly variation of PM_{10} concentrations at each monitoring station, a standard deviation was used. All statistical analyses were carried out using IBM-SPSS V.19[®] software.

A graphical comparison between the hourly variation of average PM_{10} concentration and average AC was carried out to study the influence of the hourly cycles of PM_{10} emissions from motor vehicles and industries located across the entire study area. Finally, the hourly and daily PM_{10} concentrations of monitoring stations located in areas with predominantly impervious land surface coverage (S1) were compared with the concentrations of stations located in areas with predominately vegetated land surface coverage (S2 and S3). The previous analysis also considered the prevailing AC with relation to the type of land surface coverage (LSC). In this study, four types of urban LSC were considered: vegetated (trees and grasslands), non-vegetated (bare soil), impervious (roof tops, pavements, and footpaths), and water bodies (rivers, lakes, and wetlands). In the selected monitoring stations, there was no LSC variation during the PM_{10} monitoring period.

Results

PM_{10} concentrations

For all monitoring stations, an increase in the average hourly PM_{10} concentration was observed at 5 a.m. and a decrease between 11 a.m. and noon. Peaks in PM_{10} concentrations were observed between 8 - 9 a.m. (Figure 2). These peaks in PM_{10} concentrations coincided with the highest intensity of traffic in Bogotá, which precedes the commencement of standard work activities. During the study period, the results showed, on average, a similar trend in PM_{10} concentrations per hour for all monitoring stations. A Spearman's correlation analysis for all monitoring stations showed medium positive relationships, Spearman's r_s between 0.410 - 0.506 (Table 2). The results probably showed uniform behaviour in the hourly cycles of PM_{10} emission from motor vehicles and industries located across the entire study area. The results also showed a similar trend for the daily PM_{10} concentrations generated in this study (24-hour moving average).




Figure 2. Average hourly PM₁₀ concentration during the entire study period (n = 24, per monitoring station). S1 = Kennedy, S2 = Barrios Unidos, and S3 = Guaymaral

A Spearman’s correlation analysis for all monitoring stations showed positive relationships with a tendency for medium values, Spearman’s *rs* between 0.355 - 0.507. The results also showed uniform behaviour in the daily cycles of PM₁₀ emission from motor vehicles and industries located across the entire study area. On average, S1 displayed the highest daily PM₁₀ concentrations with daily concentrations of 115% and 146% higher than those displayed in S2 and S3, respectively.

Table 2. Correlations (Spearman’s *rs*) between monitoring stations for hourly and daily PM₁₀ concentrations

Monitoring stations	S1	S2	S3
Hourly (n = 70080, per monitoring station)			
S1	1.00		
S2	0.410 (sig. = 0.046)	1.00	
S3	0.506 (sig. = 0.012)	0.483 (sig. = 0.017)	1.00
Daily, 24-hour moving average (n = 70079, per monitoring station)			
S1	1.00		
S2	0.447 (sig. < 0.001)	1.00	
S3	0.355 (sig. < 0.001)	0.507 (sig. < 0.001)	1.00

Air quality standards

An initial assessment of daily PM₁₀ concentrations was conducted in relation to the public health risk limits established by Colombian legislation (MAVDT, 2010) and WHO (2005) for a 24-hour exposure time. The results showed that S1 station was the most critical from a public health viewpoint. This monitoring station exceeded the Colombian daily limit 31% of the time during the study period, but especially during the month of February, where the average daily PM₁₀ concentration was 107.8

µg/m³ (Figure 3). During that month, the daily legislative limit was exceeded 54.1% of the time. In contrast, the daily PM₁₀ concentrations recorded at the S2 and S3 stations complied with the Colombian limit 99.5% and 100% of the time during the study period respectively. According to the daily limit established by WHO, the results showed on a monthly average that PM₁₀ concentrations in the S1 and S2 stations represented a public health risk during all months of the year (PM₁₀ > 71 µg/m³) and in February (PM₁₀ = 52.2 µg/m³), respectively. The monitoring stations S1 and S2 exceed this daily limit 92.7% and 31.7% of the time during the study period. S3 station showed the lowest public health risk. This station exceeds the daily limit 13.0% of the time during the study period.

The order of precedence established by quartiles (Q) to assess the risk per daily exposure to PM₁₀ showed on average that the first three months of the year were Q1 in all monitoring stations. February was highlighted as the highest risk for public health, followed in order of precedence for the months of March and January (Figure 3). In contrast, the Q4 months or lowest risk for public health were in order of precedence July, June, and August. A similar monthly trend was observed for all monitoring stations during the study period.




Figure 3. Average daily PM₁₀ concentrations at the monitoring stations (S1, S2, and S3). Month: 1 = January, 2 = February, 3 = March, 4 = April, 5 = May, 6 = June, 7 = July, 8 = August, 9 = September, 10 = October, 11 = November, 12 = December.

PM = PM₁₀ concentration

During the month of greatest risk to public health (February), variations in the concentration of PM₁₀ between the days of the week concerning legislative limits referenced for an exposure time of 24 hours were studied. On average, the results showed that at station S1, PM₁₀ concentrations represented a health risk during most days of the week in relation to the Colombian limit,

except for the period of exposure between Sunday and Monday ($PM_{10} = 96.2 \mu\text{g}/\text{m}^3$). The previous trend was similar at station S1 and S2 from the daily limit established by WHO (Figure 4). On average, station S3 was the only one that complied with the legislative limits referenced in this study during February.




Figure 4. Average daily PM_{10} concentration during February at the monitoring stations (S1, S2, and S3). Day: 1 = Monday, 2 = Tuesday, 3 = Wednesday, 4 = Thursday, 5 = Friday, 6 = Saturday, and 7 = Sunday. PM = PM_{10} concentration

Atmospheric condition

The prevailing AC for stations S1, S2, and S3 between 6 - 18 hours (daytime) was slightly unstable (AC = 4; occurrence frequency for 24 h, f-24 h = 19.5%), unstable (AC = 5; f-24 h = 22.7%), and unstable (AC = 5; f-24 h = 24.5%), respectively. As for the prevailing AC between 18 - 6 h (night time), it was stable (AS = 1): 35.0% (f-24 h), 50.0% (f-24 h) and 49.2% (f-24 h), respectively. On average, for the entire study area, the results indicated that the prevailing AC during the daytime (6 - 18 h) was slightly unstable and unstable (AC between 4 and 5.5; f-24 h = 46.1%). At night time (18 - 6 h), the prevailing AC was stable (AS = 1; f-24 h = 45.1%). A Kruskal-Wallis test for stations S1, S2, and S3 showed that there were no significant hourly variations in AC ($p\text{-value} = 0.827$). In other words, there was similar behaviour in the hourly AC during the study period for all monitoring stations. Figure 5 shows the average hourly AC during the sampling period from the quantitative scale used in this study.

However, on an average monthly basis the results showed comparatively that the order of precedence for the degree of daytime atmospheric instability was as follows: S3 (AC = 5.5) > S2 (AC between 5 and 5.5) > S1 (AC between 4 and 5). The results also showed that the order of precedence for PM_{10} concentration was as follows: S1 ($90.2 \mu\text{g}/\text{m}^3$) > S2 ($41.3 \mu\text{g}/\text{m}^3$) > S3 ($35.6 \mu\text{g}/\text{m}^3$). Thus, the results indicated that the daily PM_{10} concentrations were

the lowest at the monitoring station that associated the highest degree of daytime atmospheric instability (see S3 in Figure 6).




Figure 5. Average hourly AC at the study areas (S1, S2, and S3). AC: 1 = stable, 2 = slightly stable, 3 = neutral, 3.5 = neutral to slightly unstable, 4 = slightly unstable, 4.5 = slightly unstable to unstable, 5 = unstable, 5.5 = unstable to very unstable, 6 = very unstable. PM = Average hourly PM_{10} concentration

Discussion

Urban PM_{10} pollution

On an hourly and daily basis, the results suggested a similar trend in PM_{10} concentrations between all monitoring stations (Spearman's r_s between 0.355 - 0.507; see Table 2). This trend was associated with the uniform behaviour




Figure 6. Monthly comparison between PM_{10} concentration and day-time atmospheric stability. Month: 1 = January, 2 = February, 3 = March, 4 = April, 5 = May, 6 = June, 7 = July, 8 = August, 9 = September, 10 = October, 11 = November, 12 = December. AC: 1 = stable, 2 = slightly stable, 3 = neutral, 3.5 = neutral to slightly unstable, 4 = slightly unstable, 4.5 = slightly unstable to unstable, 5 = unstable, 5.5 = unstable to very unstable, 6 = very unstable. PM = average daily PM_{10} concentration

in the PM_{10} emission cycles by mobile (vehicles), fixed (industries), and natural sources in the study areas. The monitoring stations selected in this study covered 21.9 km (in a straight line) with relation to the 33 km length of the city, from north to south. Therefore, the results suggested that this uniform trend in PM_{10} emission cycles per hour and per day was probably similar in all sectors of the city of Bogotá.

On an hourly timescale, the results allowed us to identify the most critical time interval for PM_{10} from the point of view for urban public health. On average, this time interval was from 6 - 11 a.m., with the maximum PM_{10} concentrations between 8 - 9 a.m. (Figure 2). Thus, public health surveillance and control agencies should establish guidelines for sensitive population groups (children and older adults) during these time periods. For example, there should be a restriction of outdoor physical activities at educational institutions for children - especially around the monitoring station S1, where the average hourly PM_{10} concentrations

were 115% and 144% higher in relation to that observed at S2 and S3, respectively.

The reference legislation made it possible to demonstrate that the most critical monitoring station within the public health framework was S1. On average, this station exceeded the Colombian daily limit (24-h exposure time) during February by 7.8%, and the WHO daily limit for all months of the year by 80.4% (Fig. 3). Based on the WHO Air Quality Guide (WHO, 2005), the daily PM_{10} concentration observed in February at the S1 monitoring station (average = $107.8 \mu\text{g}/\text{m}^3$) could increase the short-term mortality rate by 2.9% with relation to the mortality rate observed for a concentration of $50 \mu\text{g}/\text{m}^3$ (WHO maximum permissible level). The WHO (2005) reported that the adverse effects on human health could not be ruled out below the established maximum permissible level, but that compliance with these standards supposes a significant reduction of the risks on public health. On average, for all months, in the area of influence around the S1 monitoring station, there was probably an increase of 2.0% in the daily mortality rate associated with PM_{10} . S2 and S3 monitoring stations complied with the Colombian limit during all months, and the S3 station was the only one that complied with the WHO limit throughout the year.

The order of precedence established by quartiles to assess the monthly risk for exposure to PM_{10} in Bogotá was as follows: Q1 (February > March > January), Q2 (May > December > October), Q3 (November > September > April), and Q4 (August > June > July). On average, during the Q1 months the daily PM_{10} concentrations were shown to be 11.5%, 16.8%, and 35.9% higher than those observed in the Q2, Q3 and Q4 months respectively (Figure 3). In the present study, the results suggested that public health surveillance and control agencies should plan and implement different strategies according to the monthly Q value. For example, there should be more demanding strategies of PM_{10} control and surveillance during the months of greatest public health risk (Q1) in comparison to the months of lowest public health risk (Q4). On average, the short-term mortality rate during the Q1 months could be increased by between 0.46% - 1.32% with relation to the short-term mortality rate of Q4 months. This was based on an increase in the short-term mortality rate proposed by WHO Air Quality Guide (WHO, 2005).

Additionally, variation of the PM_{10} concentration during each day of the week for the most critical month (February) was evaluated from all monitoring stations.

On average, the results showed an increase in daily PM_{10} concentrations as the week progressed, reaching its maximum level during the exposure period between Thursday and Saturday (Figure 4). PM_{10} concentrations during this exposure period exceeded the Colombian daily limit at station S1 (15.6%), and the WHO daily limit at the stations S1 and S2 ($S1 = 131\%$; $S2 = 12.9\%$). Monitoring station S3 complied with all legislative limits of reference. A similar trend was observed in daily PM_{10} concentrations for other months. Therefore, the results suggested that this exposure period is the highest risk during the week in relation to the possible PM_{10} impact on human health.

On a weekly basis, the highest daily PM_{10} concentrations tended to occur during the exposure period between Friday and Saturday (Figure 4). On average, concentrations during this exposure period were higher ($S1 = 14.3\%$, $S2 = 13.3\%$, and $S3 = 16.8\%$) compared to the concentrations observed during the exposure period between Sunday and Friday. This trend was probably due to the fact that in Bogotá there was a restriction on vehicular movement due to traffic congestion from Monday to Friday. This traffic restriction prevented the use of 40% of the city's vehicles between 6 and 9 a.m., and 3 and 7 p.m. In other words, there was no traffic restriction on Saturdays; which probably explained the increase in PM_{10} concentrations on this day.

AC influence

With relation to the influence of AC, the results showed the best possible scenario from a public health viewpoint for maximum hourly PM_{10} concentrations. Namely, the maximum concentrations recorded by the monitoring stations S1 ($135 \mu\text{g}/\text{m}^3$), S2 ($62.4 \mu\text{g}/\text{m}^3$) and S3 ($52.3 \mu\text{g}/\text{m}^3$) occurred during daytime periods (8 - 9 a.m.), when the prevailing AC was between unstable and very unstable (Figure 5). This suggested high PM_{10} dispersion and a probable reduction in hourly concentration during these extreme air pollution episodes. Otherwise, the maximum PM_{10} concentrations in Bogotá during daytime would probably have been higher.

The results showed that during night time the prevailing ACs were stable ($AC = 1$). This suggested a low PM_{10} dispersion during these time periods and therefore a probable increase in hourly concentrations. Again, the best possible scenario in the public health framework for monitoring stations S1 and S2 occurred given that during

night time periods the lowest hourly PM_{10} concentrations were observed (Figure 5). On average, night time PM_{10} concentrations at monitoring stations S1 and S2 were 20.1% and 4.61% lower than those observed in daytime. However, hourly PM_{10} concentrations at the S3 station were different. On average, at this monitoring station, the night time PM_{10} concentrations were 1.39% higher than those observed in daytime, despite a probable decrease in the emission sources of PM_{10} such as traffic and industries. The results suggested that the increase in night time PM_{10} concentrations at station S3 was probably associated with the prevailing ACs, which were stable (low PM_{10} dispersion). Lee et al. (2013) and Vecchi et al. (2007) reported a similar trend under prevailing conditions of night time atmospheric stability ($AC = 1$) in the cities of Seoul (Korea) and Milan (Italy).

From the above, the analyses showed the hourly variation of PM_{10} concentrations between monitoring stations using station S3 as a reference, and considering that this monitoring station tended to show the highest degree of atmospheric instability in the daytime (S3, AC between 5 - 5.5; S2 and S1, AC between 4 - 5.5). The results showed that during the night time there were no differences between monitoring stations in the hourly AC ($AC = 1$). Hourly PM_{10} concentrations at the S1 and S2 monitoring stations were 144% and 13.3% higher than those observed at S3 station (Figure 5). Additionally, at this last monitoring station there was less difference between daytime and night time concentrations of PM_{10} (difference = 1.39%) in comparison with the monitoring stations S1 (difference = 20.9%) and S2 (difference = 4.61%). This trend was also supported by a lower hourly variation than the average PM_{10} concentrations at monitoring station S3, evaluated by the standard deviation (S3, $\sigma = 6.27 \mu\text{g}/\text{m}^3$; S2, $\sigma = 8.18 \mu\text{g}/\text{m}^3$; S1, $\sigma = 17.8 \mu\text{g}/\text{m}^3$). Thus, the results suggested that the monitoring stations located in urban areas where there were atmospheric conditions of greater instability tended to show lower concentrations and variations of hourly PM_{10} .

Daily PM_{10} concentrations showed a trend like that described above for hourly PM_{10} concentrations. Monthly, the results showed, on average, that the daily PM_{10} concentrations were lower in the monitoring station associated with the highest degree of atmospheric instability (S3). The order of precedence in the degree of atmospheric instability for monitoring stations was as

follows: $S3 > S2 > S1$ (Figure 6). The order of precedence for daily PM_{10} concentrations was as follows: $S1 > S2 > S3$. Results suggested an inverse relationship between PM_{10} concentration and atmospheric instability.

Finally, in the present study, a relationship was observed between atmospheric instability and the type of land surface coverage (LSC). The results showed that urban areas of greater atmospheric instability (lower PM_{10} concentration) were characterized by a predominantly vegetated LSC ($S2 = 86.2\%$; $S3 = 74.6\%$; see Table 1). In contrast, urban areas of lower atmospheric instability (greater PM_{10} concentration) were associated with a predominantly impervious LSC ($S1 = 68.9\%$). Studies in urban areas have reported a significant influence of a vegetated LSC on PM_{10} concentrations. For example, Chen et al. (2015) found that the presence of trees in Wuhan (China) reduced PM_{10} concentrations from between 7% - 15%. Similarly, Shan et al. (2007) and Islam et al. (2012) reported a reduction of total suspended particulates ($<100 \mu m$) in urban areas by 30% (Shanghai, China) and 55% (Khulna, Bangladesh), respectively. A vegetated LSC also acted as an effective sink for gaseous and particulate atmospheric pollutants (Fowler et al. 2004).

Conclusions

The results suggest that public health surveillance and control agencies should pay more attention to urban sectors with lower atmospheric instability, because these urban sectors show higher concentrations (between 131% - 144%) and variations of PM_{10} compared to urban sectors of greater atmospheric instability. On average, urban sectors with lower atmospheric instability show an increase in the daily mortality rate of 2.0% with relation to the mortality rate for a daily PM_{10} concentration of $50 \mu g/m^3$ (WHO limit). According to WHO (2005), an increase in the daily mortality rate of 5% requires immediate corrective measures. The type of urban LSC can probably be used as a public health indicator, since in urban sectors where there is evidence of greater increases in the daily mortality rate by PM_{10} (lower atmospheric instability), an impervious LSC predominates instead of a vegetated LSC.

Finally, the findings of this study serve as a reference point to deepen our knowledge about the behaviour of PM_{10} concentrations and the influence of AC, allowing us to develop and implement different strategies of surveillance

and control for public health in megacities. However, the following limitations form part of this study, requiring further attention. First, the number of years for the PM_{10} time series (six years) was limited due to LSC changes in the areas covered by the monitoring stations; for analysis of the influences on the AC upon PM_{10} concentrations, the same LSC distribution is required throughout the study period. Second, the monitoring stations do not have similar characteristics regarding the distance and emission intensity of the PM_{10} sources (highways and industries). Third, PM_{10} concentrations and AC for the entire study area are evaluated using data from only three monitoring stations, so the number of stations could be increased.

Acknowledgements

The author wishes to thank the Universidad Distrital Francisco José de Caldas (Colombia) and the Environment Secretary of Bogotá (Colombia). Finally, the author is very grateful for the helpful comments and suggestions offered by reviewers.

Conflict of interest

The author states that there is no conflict of interest.

References

- Chambers, S. D., Wang, F., Williams, A. G., Xiaodong, D., Zhang, H., Lonati, G.,... Allegrini, I. (2015). Quantifying the influences of atmospheric stability on air pollution in Lanzhou, China, using a radon-based stability monitor. *Atmospheric Environment*, 107, 233-243. <https://doi.org/10.1016/j.atmosenv.2015.02.016>
- Chen, X., Pei, T., Zhou, Z., Teng, M., He, L., Luo, M., & Liu, X. (2015). Efficiency differences of roadside greenbelts with three configurations in removing coarse particles (PM_{10}): A street scale investigation in Wuhan, China. *Urban Forestry & Urban Greening*, 14(2), 354-360. <https://doi.org/10.1016/j.ufug.2015.02.013>
- Dahari, N., Latif, M.T., Muda, K., & Hussein, N. (2020). Influence of meteorological variables on suburban atmospheric $PM_{2.5}$ in the southern region of peninsular Malaysia. *Aerosol and Air Quality Research*, 20, 14-25. <https://doi.org/10.4209/aaqr.2019.06.0313>

- Dockery, D. W., Pope, C. A., Xu, X., Spengler, J. D., Ware, J. H., Fay, M. E.,... Speizer, F. E. (1993). An association between air pollution and mortality in six U.S. cities. *New England Journal of Medicine*, 329(24), 1753-1759. <https://doi.org/10.1056/NEJM199312093292401>
- Fowler, D., Skiba, U., Nemitz, E., Choubedar, F., Branford, D., Donovan, R., & Rowland, P. (2004). Measuring Aerosol and Heavy Metal Deposition on Urban Woodland and Grass Using Inventories of ^{210}Pb and Metal Concentrations in Soil. In R. K. Wieder, M. Novák, & M. A. Vile (Eds.), *Biogeochemical Investigations of Terrestrial, Freshwater, and Wetland Ecosystems across the Globe* (pp. 483-499). Springer Netherlands. https://doi.org/10.1007/978-94-007-0952-2_33
- Franck, U., Leitte, A. M., & Suppan, P. (2014). Multiple exposures to airborne pollutants and hospital admissions due to diseases of the circulatory system in Santiago de Chile. *The Science of the Total Environment*, 468-469, 746-756. <https://doi.org/10.1016/j.scitotenv.2013.08.088>
- Gifford, F. A. (1976). *Turbulent diffusion-typing schemes: A review*. http://inis.iaea.org/Search/search.aspx?orig_q=RN:8296495
- Hernández, E., Martín, F., & Valero, F. (1992). Statistical forecast models for daily air particulate iron and lead concentrations for Madrid, Spain. *Atmospheric Environment. Part B. Urban Atmosphere*, 26(1), 107-116. [https://doi.org/10.1016/0957-1272\(92\)90042-Q](https://doi.org/10.1016/0957-1272(92)90042-Q)
- Hou, Q., An, X., Tao, Y., & Sun, Z. (2016). Assessment of resident's exposure level and health economic costs of PM_{10} in Beijing from 2008 to 2012. *Science of the Total Environment*, 563-564, 557-565. <https://doi.org/10.1016/j.scitotenv.2016.03.215>
- Islam, M. N., Rahman, K.-S., Bahar, M. M., Habib, M. A., Ando, K., & Hattori, N. (2012). Pollution attenuation by roadside greenbelt in and around urban areas. *Urban Forestry & Urban Greening*, 11(4), 460-464. <https://doi.org/10.1016/j.ufug.2012.06.004>
- Katsouyanni, K., Schwartz, J., Spix, C., Touloumi, G., Zmirou, D., Zanobetti, A., ... Anderson, H. R. (1996). Short term effects of air pollution on health: A European approach using epidemiologic time series data: the APHEA protocol. *Journal of Epidemiology and Community Health*, 50(Suppl 1), S12-S18.
- Lacasaña, M., Aguilar, C., & Romieu, I. (1999). Evolución de la contaminación del aire e impacto de los programas de control en tres megaciudades de América Latina. *Salud Pública de México*, 41(3), 203-215.
- Lee, S., Ho, C.H., Lee, Y. G., Choi, H.J., & Song, C.K. (2013). Influence of transboundary air pollutants from China on the high- PM_{10} episode in Seoul, Korea for the period October 16–20, 2008. *Atmospheric Environment*, 77, 430-439. <https://doi.org/10.1016/j.atmosenv.2013.05.006>
- Liu, W., Xu, Z., & Yang, T. (2018). Health effects of air pollution in China. *International Journal of Environmental Research and Public Health*, 15(7). <https://doi.org/10.3390/ijerph15071471>
- Maesano, C. N., Morel, G., Matynia, A., Ratsombath, N., Bonnetty, J., Legros, G.,...Annesi-Maesano, I. (2020). Impacts on human mortality due to reductions in PM_{10} concentrations through different traffic scenarios in Paris, France. *Science of The Total Environment*, 698, 134257. <https://doi.org/10.1016/j.scitotenv.2019.134257>
- MAVDT. (2010). *Resolución 610 de 2010 Ministerio de Ambiente, Vivienda y Desarrollo Territorial*. <https://www.alcaldiabogota.gov.co/sisjur/normas/Normal.jsp?i=39330>
- Montoya, G. de J., Cepeda, W., & Eslava, J. A. (2004). Características de la turbulencia y de la estabilidad atmosférica en Bogotá. *Revista de La Academia Colombiana de Ciencias Exactas, Físicas y Naturales*, 28(108), 327-335.
- Palacio, D. F., Zafra, C.A., & Rodríguez, J. P. (2014). Evaluation of the air quality by using a mobile laboratory: Puente Aranda (Bogotá D.C., Colombia). *Revista Facultad de Ingeniería Universidad de Antioquia*, 71, 153-166.
- Pasquill, F. (1961). The estimation of the dispersion of wind-borne material. *Meteorol. Mag.*, 90, 33-49.
- Sarmiento, R., Hernández, L. J., Medina, E. K., Rodríguez, N., & Reyes, J. (2015). Respiratory symptoms associated with air pollution in five localities of Bogotá, 2008-2011, a dynamic cohort study. *Biomédica*, 35(SPE), 167-176. <https://doi.org/10.7705/biomedica.v35i0.2445>
- Shan, Y., Jingping, C., Liping, C., Zhemin, S., Xiaodong, Z., Dan, W., & Wenhua, W. (2007). Effects of vegetation status in urban green spaces on particle removal in a street canyon atmosphere. *Acta Ecologica Sinica*, 27(11), 4590-4595. [https://doi.org/10.1016/S1872-2032\(08\)60007-4](https://doi.org/10.1016/S1872-2032(08)60007-4)
- Srinivas, C.V., Venkatesan, R., Somayaji, K. M., & Indira, R. (2009). A simulation study of short-range atmospheric dispersion for hypothetical air-borne effluent releases using different turbulent diffusion methods. *Air Quality, Atmosphere & Health*, 2(1), 21-28. <https://doi.org/10.1007/s11869-009-0030-6>
- Terrouche, A., Ali-Khodja, H., Kemmouche, A., Bouziane, M., Derradji, A., & Charron, A. (2016). Identification of sources of atmospheric particulate matter and trace metals in Constantine, Algeria. *Air Quality, Atmosphere & Health*, 9(1), 69-82. <https://doi.org/10.1007/s11869-014-0308-1>
- Turner, D. B. (1964). A diffusion model for an urban area. *Journal of Applied Meteorology*, 3(1), 83-91. [https://doi.org/10.1175/1520-0450\(1964\)003<0083:ADMFAU>2.0.CO;2](https://doi.org/10.1175/1520-0450(1964)003<0083:ADMFAU>2.0.CO;2)

- U.S.EPA. (1999). *Compendium of methods for the determination of inorganic compounds in ambient air*.
- Vecchi, R., Marcazzan, G., & Valli, G. (2007). A study on night time–daytime PM₁₀ concentration and elemental composition in relation to atmospheric dispersion in the urban area of Milan (Italy). *Atmospheric Environment*, 41(10), 2136-2144. <https://doi.org/10.1016/j.atmosenv.2006.10.069>
- WHO. (2005). *WHO | Air quality guidelines—Global update 2005*. WHO. https://www.who.int/phe/health_topics/outdoorair/outdoorair_aqg/en/
- Yang, J., Zhou, M., Zhang, F., Yin, P., Wang, B., Guo, Y.,... Liu, Q. (2020). Diabetes mortality burden attributable to short-term effect of PM₁₀ in China. *Environmental Science and Pollution Research*. <https://doi.org/10.1007/s11356-020-08376-1>
- Zafra, C., Ángel, Y., & Torres, E. (2017). ARIMA analysis of the effect of land surface coverage on PM₁₀ concentrations in a high-altitude megacity. *Atmospheric Pollution Research*, 8(4), 660-668. <https://doi.org/10.1016/j.apr.2017.01.002>
- Zafra-Mejía, C., Gutiérrez-Malaxechebarria, Á., & Hernández-Peña, Y. (2019). Correlation between vehicular traffic and heavy metal concentrations in road sediments of Bogotá, Colombia. *Revista Facultad de Medicina*, 67(2), 193-199. <https://doi.org/10.15446/revfacmed.v67n2.68269>
- Zoras, S., Triantafyllou, A. G., & Deligiorgi, D. (2006). Atmospheric stability and PM₁₀ concentrations at far distance from elevated point sources in complex terrain: Worst-case episode study. *Journal of Environmental Management*, 80(4), 295-302. <https://doi.org/10.1016/j.jenvman.2005.09.010>