

Pre-operative fasting guidelines: an update

**E. Søreide¹, L. I. Eriksson², G. Hirlekar³, H. Eriksson⁴,
S. W. Henneberg⁵, R. Sandin⁶, J. Raeder⁷ (task Force On
Scandinavian Pre-operative Fasting Guidelines, Clinical Practice Committee
Scandinavian Society Of Anaesthesiology And Intensive Care Medicine)**

Liberal pre-operative fasting routines have been implemented in most countries. In general, clear fluids are allowed up to 2 h before anaesthesia, and light meals up to 6 h. The same recommendations apply for children and pregnant women not in labour. In children <6 months, most recommendations now allow breast- or formula milk feeding up to 4 h before anaesthesia.

Recently, the concept of pre-operative oral nutrition using a special carbohydrate-rich beverage has also gained support and been shown not to increase gastric fluid volume or acidity. Based on the available literature, our Task Force has produced new consensus-based Scandinavian guidelines for pre-operative fasting. What is still not clear is to what extent the new liberal fasting routines should apply to patients with functional dyspepsia or systematic diseases such as diabetes mellitus. Other still controversial areas include the need for and effect of fasting in emergency patients, women in labour and in association with procedures done under 'deep sedation'.

We think more research on the effect of various fasting regimes in subpopulations of patients is needed before we can move one step further towards completely evidence-based pre-operative fasting guidelines.

Key words: Anaesthesia; general; gastric content; gastric emptying; preoperative fasting; pulmonary aspiration complications.

Anaesthesia-related pulmonary aspiration leading to respiratory failure (Aspiration Pneumonitis; Mendelson's syndrome) has been described in both elective and emergency surgical patients¹⁻⁷. With the hope of reducing the risk of this complication, rigid fasting routines before surgery have been enforced⁸. However, the scientific basis for these rigid fasting routines in elective patients has been challenged and found to be nonexistent⁹⁻¹¹. Based on this new information, several national anaes-

thesia societies now have accepted more liberal fasting rules for clear fluids (water, clear juices, coffee, tea)¹²⁻¹⁷. Intake of solids in the morning of elective surgery is still not recommended.

To what extent pre-operative fasting is of any importance in emergency cases is still a matter of uncertainty with variation in clinical practice.

Further, it is also not clear to what extent specific patient populations with suspected or proven

1. Department of Anaesthesia and IntensiveCare, Stavanger University Hospital, Stavanger, Norway.
2. Department of Anaesthesiology and Intensive Care, Karolinska University Hospital, Stockholm, Sweden.
3. Department of Anaesthesia and Intensive Care, Akureyri Hospital, Akureyri, Iceland.
4. Department of Anaesthesia and Intensive Care, Helsinki University Hospital, Helsinki, Finland.
5. Department of Anaesthesia, Rigshospitalet, Copenhagen, Denmark.
6. Department of Anaesthesiology and Intensive Care, Regional Hospital, Kalmar, Sweden and
7. Department of Anaesthesia, Ullevål University Hospital, Oslo, Norway.

Recibido para publicación, enero 18 de 2008, Aceptado para publicación, enero 28 de 2008.

delayed gastric emptying need to be exempt from the new fasting guidelines^{16,17}. Recently, a new method for pre-operative optimization of the elective patient, oral nutrition with carbohydrates^{16,18,19}, has been introduced, but so far has not been widely implemented into clinical practice.

This review aims to give an update on preoperative fasting and gastric content as a risk factor of pulmonary aspiration. We will focus on the development and experience with the new and more liberal clinical practice guidelines, but also present still controversial areas worth further research.

ANAESTHESIA-RELATED PULMONARY ASPIRATION: RISK FACTORS

Gastric emptying

The volume of the adult ventricle is approximately 1500 ml. The ventricle can be divided into two functional parts, i.e. the proximal and the distal part 20. The proximal part consists of the fundus, cardia and the upper part of the corpus, and acts as a reservoir for ingested food regulating the intra-gastric pressure (adaptive relaxation) and the speed of gastric emptying.

The distal part (Fig. 1) of the ventricle includes the lower part of the corpus, antrum and pylorus. The contractions of the distal part of the ventricle mix the larger solid food particles with gastric fluid. Only particles small enough (i.e. less than 1 mm) are allowed to pass via the pylorus to the duodenum.

In a normal situation, the gastric emptying of fluids is influenced by the pressure gradient between the stomach and the duodenum, and the volume, caloric density, pH and osmolality of the gastric fluid^{20,21}. In otherwise healthy patients, gastric fluid content is not increased in the immediate pre-operative period despite the theoretical negative impact of anxiety on gastric emptying^{22,23}. Gastric emptying of water and other inert, non-caloric fluids follows an extremely fast exponential curve with a mean half-time of 10 min^{20,21} (Fig. 2). Initially, glucose-loaded fluids empty a little slower, but after 90 min this difference is negligible^{20,21,23}.

In contrast, the gastric emptying curve for solids is linear^{20,21} (Fig. 2). Gastric emptying of solid food starts approximately 1 h after a meal. Within 2 h, approximately 50% of the solid food ingested is passed to the duodenum. The gastric emptying of solids is independent of the amount of food ingested but dependent on the caloric density of the meal.




Figure 1. The anatomical relationship between the gastro-oesophageal part of the digestive system and the upper airways and the lungs makes pulmonary aspiration likely during regurgitation or vomiting when the patient is anaesthetised. Previously published in the Doctoral Thesis 'Anaesthesia and gastric content. New methods and trends' (1995) by Eldar Soreide. Permission granted.




Figure 2. The gastric emptying of solids (thick dotted line) and clear fluids (thin line). Previously published in the Doctoral Thesis 'Anaesthesia and gastric content. New methods and trends' (1995) by Eldar Soreide. Permission granted.

Gastric emptying is slower in females than in males and slower in the elderly. In order to secure emptying of solids, longer fasting is needed²⁴.

In neonates and infants, clear fluids also follow first order kinetics and emptying of solids in a linear manner²⁵. Gastric emptying of human milk in mature neonates and infants is not complete after 2 h and at least 3 h seems to be required²⁶.

The optimal fasting period for human milk has not been established but it is more than 2 h and less than 5 h.

Pre-mature babies have a somewhat slower rate of gastric emptying, and cow's milk empties slower than human milk₂₆. Gastric emptying time for formulas vary with the content of the formula. One should be aware of the fact that there is a rather large variation in the composition of formula food between different regions/countries. Most paediatric anaesthesiologists now use the same 2-h limit for clear fluids as in adults, and recommend 4- to 6-h fasting after breast- and formula milk with the lower limit applied in children less than 6 months₂₅.

Delayed gastric emptying is found in numerous situations, and may be divided by aetiology into alterations in normal physiology, state of disease and intake of external agents, either drugs or substances for abuse. Pain and opioids are well-known reasons for delayed gastric emptying_{20,24}. Some systemic diseases are known to slow down the gastric emptying: among them most notably diabetes mellitus₂₇. Diabetes does affect gastric emptying much more for solids than for fluids_{20, 27}. Local gastrointestinal stasis (tumour or obstruction) may have the same effect.

Gastric emptying times for solids are delayed in smokers, but not with nicotine patch use_{28,29}. Habitual smokers have a small but statistical significant increase in gastric fluid volumes when compared with non-smokers, even when refraining from smoking₃₀. To what extent smoking affects gastric fluid emptying and volume is still controversial, but overall there seems to be good reasons for avoiding smoking immediately before anaesthesia_{24,30,31}. Recreational abuse of cannabinoids₃₂ and high doses of alcohol₃₃ also inhibit gastric emptying. Functional dyspepsia₃₄₋₃₆ is associated with a delay in gastric emptying. Obese patients seem to have a similar gastric emptying to nonobese patients, and pre-operative fluid intake does not increase gastric content₃₇. Studies on the impact of female hormones on gastric emptying have shown variable results_{20,24}. Pregnant females seem to have a normal gastric emptying rate, except for the first trimester, where a hormonal cause for slowing has been suggested₃₈. When in labour, gastric emptying will be slowed down and stay slow for at least 2 h afterwards₃₉.

Metoclopramide may improve gastric emptying in these patients but cannot assure emptying of the stomach content_{24,40}. The same goes for patients with pain or on opiate medication.

Gastric content and gastro-oesophageal reflux

The volume and acidity of the gastric content are a result of gastric secretion, oral intake and gastric emptying_{20,24}. For passive regurgitation and pulmonary aspiration to occur during anaesthesia, a certain gastric volume needs to be present. Studies₄₁ indicate that more than 200 ml is needed in an adult patient. In otherwise healthy elective patients much lower gastric fluid volumes in the range of 10-30 ml are found_{9-11,42}. In a few patients, higher volumes up to 200 ml may be found, irrespectively of intake of clear fluids or not. These outliers probably represent patients with an undetected gastric disorder such as functional dyspepsia₃₄₋₃₆. In patients with gastro-oesophageal reflux or if active vomiting occurs, even smaller gastric volumes may be propelled up and into the trachea_{2,7,43} (Fig. 1).

Patient and anaesthetic factors

Airway management problems frequently precipitate pulmonary aspiration₃₋₇. Air blown into the stomach and bucking and coughing due to light anaesthesia may all cause gastro-oesophageal reflux episodes. Obese patients, patients with known gastro-oesophageal reflux disease and patients with difficult airways are particular prone to pulmonary aspirations, independent of their gastric content.

Kruger et al.₇ found such patient factors together with poor judgement in choice and performance of anaesthetic method to be the most important factors predisposing clinical significant pulmonary aspiration, and not violations of fasting precautions. This certainly put our historic overemphasis on gastric content into perspective. It is important to differentiate between what happens when airway manipulation during a light stage of anaesthesia induces active vomiting or gastro-oesophageal reflux episodes independently of the volume of gastric content, and the situation with a distended stomach pouch and anaesthesia that causes the oesophageal sphincters to relax and passive flow (regurgitation) of gastric content into the upper airways and pulmonary aspiration₂. The anaesthetist is probably as an important factor as the gastric content.

NEW GUIDELINES FOR PRE-OPERATIVE FASTING

Clinically controlled studies and meta- analysis

Numerous controlled studies and meta-analysis have concluded that in otherwise healthy adults scheduled for elective surgery, oral intake of water

and other clear fluids (tea, coffee, soda water, apple and pulp-free orange juice) up to 2 h before induction of anaesthesia does not increase gastric fluid volume or acidity^{9-11,15,44}. The studies were performed in both male and females adults (the study was in adults), and in different countries⁴⁴. Hence, according to the evidence-based medicine classification⁴⁵, the present scientific evidence allows a Level 1 recommendation for more liberal fasting routines for clear fluids.

National and anaesthesia society guidelines

Based on the new data, most national anaesthesiology societies now recommend no more than 2-h fasting for clear fluids (water, tea, coffee, pulp-free fruit juices) in elective patients, both adults and children and including pregnant women not in labour¹²⁻¹⁷. Importantly, this does not apply to milk, any other fat-containing fluids, or solids. No complications associated with the new and more liberal fasting guidelines have been reported^{44,46}. To provide sufficient safety margins, the fasting period after intake of solids should not be less than 6 h^{47,48}. Further, although shown to affect gastric content not all national societies guidelines include information on the use of chewing gum, tobacco and pre-operative medications in the immediate pre-operative period. Chewing gum and tobacco use both increase gastric content, but to what extent the increase is of any clinical significance is very uncertain³⁰. Still, we think their Pre-operative

fasting guidelines use should be discouraged in the immediate preoperative period¹⁴.

Oral benzodiazepines are commonly used for premedication. Up to 150 ml of water together with oral medication up to 1 h before induction of anaesthesia is perfectly safe in adults⁴². Based on the prolonged gastric emptying seen with the use of opiates, it is reasonable to stop fluid intake 1 h before the use of opiate premedication⁴⁹.

New Scandinavian guidelines

Our Task Force aimed at making one combined but not too detailed practice guideline for pre-operative fasting for all the Scandinavian countries. We concluded that based on the current knowledge, a general recommendation of 2-h fasting for clear fluids and 6 h for solids in otherwise healthy elective patient is appropriate (Table 1). This guideline is non-controversial and valid both for children>1 years, adults and pregnant women not in labour. Similar to others, we define clear fluids as water, coffee, tea, pulp-free juice and soft drinks, but also included the pre-operative carbohydrate drink intended for pre-operative nutrition (Nutricia Preop®, Numico, The Netherlands)^{16,19,23}. The restrictions for solids include soups, yoghurt, sour milk or milk-containing drinks. We felt that our Scandinavian consensus-based clinical practice guidelines should not go into more detail but leave this to the national societies. Instead, we also decided to include topics still controversial or topics where more research is needed.

Table 1

Scandinavian guidelines for pre-operative fasting in elective patients.
Patients (adults as well as children) may drink clear fluids up to 2 h prior to general or regional anaesthesia Patient should not take solid food 6 h prior to induction of anaesthesia Breast-feeding should be stopped 4 h prior to induction of anaesthesia. The same applies to formula milk Adults may drink up to 150 ml of water with pre-operative oral medication up to 1 h before induction of anaesthesia, and children up to 75 ml. Use of chewing gum and any form of tobacco should be discouraged during the last 2 h prior to induction of anaesthesia
These guidelines also apply to elective Caesarean sections. Clear fluids are defined as non-particulate fluids without fat, for example, water, clear fruit juice, tea or coffee. Both cow's milk and powdered milk are treated as solid food. Patients with known or suspected delay in gastric emptying (diabetes mellitus, upper gastrointestinal pathology and symptoms) should be assessed on an individual basis.

CONTROVERSIAL TOPICS AND TOPICS FOR FUTURE RESEARCH

Patient groups exempt from the liberal fasting guidelines

While the new, liberal fasting guidelines can be safely used for the majority of elective patients, it is important to emphasise that pre-operative fasting is still strictly recommended for all emergency surgery cases. The delayed gastric emptying in emergency cases may be due to both the effect of pain per se, the opioids given or gastrointestinal obstruction_{2, 24}. Hence, fasting such patients will never make them 'fasted and elective'. The same applies to pregnant women in labour₅₀.

There are also elective patients where a significant delayed gastric emptying must be suspected. These include patients with gastrointestinal obstruction of any form, or cancer in the upper gastrointestinal tract. When it comes to choice of anaesthetic technique, patients with a known hiatus hernia have a greater risk of regurgitation and should be handled as 'at risk of regurgitation'. However, there is no clear evidence of slower gastric emptying or greater residual gastric volumes in these patients_{16, 17}.

There is a high prevalence of delayed gastric emptying and gastro-paresis in patients with upper gastrointestinal symptoms, which is not influenced by the presence of organic disease₃₄₋₃₆. Hence, such patients should probably be fasted after intake of solids for more than 6 h. How long, however, is not known. Although the effect on gastric emptying of fluids probably is much less, more controlled trials are needed in these patients₄₄.

In patients with systemic disease, the extent of gastric slowing may be highly variable depending on the severity of the disease_{20, 27}. Most investigations have been carried out in diabetes mellitus where the gastric slowing is due to polyneuropathy in the innervations of the gastrointestinal system with advanced disease. To what extent diabetic patients should be nil per mouth after midnight to secure gastric content in the normal range is still not known. Diabetes and other medical conditions do affect gastric emptying much more for solids than for fluids_{20, 24, 27}. Probably, a 2-h fasting period for clear fluids is also enough in patients with systemic diseases. More studies, however, are needed before a scientific validated answer can be given. In the mean time, we think it is wise to err on the conservative side when it comes to fasting after intake of solids in these patients.

Women going into labour have very prolonged gastric emptying times₃₉ and have an increased incidence of pulmonary aspirations compared with other patient groups₆. Despite this fact, most maternity wards encourage oral intake during labour₅₀. This may sound counter-productive for us as anesthesiologists, but to obstetricians, midwives and the women themselves, the small risk of an emergency Caesarean-section under general anaesthesia may not be a valid argument to impose unphysiological starvation during a natural process with a large need for calories₅₀. A trade-off that midwives and obstetricians may accept is to allow fluids but no solids during labour. Anaesthesiologists are not in the position to decide the fasting guidelines for women in labour. We need more data on the actual practice and possible adverse effects in Scandinavian maternity systems before we can move forward on this topic.

Sedation and need for pre-sedation fasting

An increasing number of surgical procedures are done with 'light, conscious or deep sedation' in various combinations with local and regional anaesthesia. Should these patients be included in the preoperative fasting guidelines? Sedation and analgesics tend to impair airway reflexes in proportion to the degree of sedation/analgesia achieved_{51, 52}. The available literature does not provide sufficient evidence to conclude that pre-procedure fasting results in a decreased incidence of adverse outcomes in patients undergoing either moderate or deep sedation. However, the American Society of Anesthesiologists recommends that patients undergoing sedation/analgesia for elective procedures should have the same restrictions as patients undergoing general anaesthesia₅₂. These guidelines are arbitrary and based upon consensus opinion.

In emergency situations, the potential for pulmonary aspiration of gastric contents must be considered. Green et al.₅₁ found that pulmonary aspiration during emergency department procedural sedation and analgesia had not been reported in medical literature. Therefore, there is little evidence to support specific fasting periods.

Steeds and Mather₅₃ surveyed the policy of preoperative fasting in connection with eye surgery under regional anaesthesia. Fifty per cent of the respondents felt that fasting was not necessary and mentioned hypoglycaemia, faint, thirst, nausea, headache, and dizziness as complications to prolonged starvation. Maltby and Hamilton₅₄ found no case of pulmonary aspiration in 30 000 patients

undergoing cataract surgery done under regional anaesthesia. Only 1% needed sedation. Since 1984, they have allowed breakfast before the procedure. Still, they cautioned against heavy sedation and conversion of regional block into general anaesthesia. Most eye surgery can be done with local anaesthesia only. It looks like pre-operative fasting ensures very little extra patient safety, and at the expense of patient comfort. Post-operative emesis which is detrimental after ophthalmic surgery might be reduced by shorter pre-operative fast. It seems like the key points are to make the ophthalmologists aware of the potential danger of heavy sedation and non-fasting and to make local guidelines that take into account the type of surgery, type of local and regional anaesthesia, the need for sedation and the possibility of having to convert a failed regional anaesthetic to a general one. We feel that more data on the current sedation practice in elective and emergency cases in Scandinavia are needed before we can produce specific recommendations on pre-procedural fasting in these situations.

Pre-operative fasting vs. oral nutrition

The metabolic implications of prolonged starvation vs. shorter fasting times are also important.¹⁸ Studies have indicated that the availability of carbohydrates and the metabolic setting of the fed state are important factors which improve post-operative recovery.¹⁶ The main objective of pre-operative carbohydrate treatment is to cause a change in metabolism that normally takes place when someone takes their breakfast. This will elicit an endogenous insulin release that turns off the overnight fasting state of the metabolism. A carbohydrate-rich (12.5%) clear beverage containing mainly polymers of carbohydrates to minimize the osmotic load and thus reduce the gastric emptying time has been tested.²³ Both in

healthy volunteers and in preoperative patients 400 ml passed the stomach within 90 min. Studies in more than 250 patients have shown that the median residual gastric volume is only approximately 20 ml.^{16,18} A small fraction of the patients had gastric volumes above 120 ml, the highest being 200 ml. When the 400-ml dose was divided into 2 200 ml, the last intake 2 h before the gastroscopy, the highest volume Pre-operative fasting guidelines found was 120 ml, with the averages approximately 35 ml.

Studies have found that this carbohydrate-rich pre-operative beverage both improves subjective well-being compared with a placebo (water) and may positively affect the post-operative recovery.^{16,19} From a patient safety point of view, it is important to notice that intake of up to 400 ml of the beverage does not produce negative effects on the gastric content compared with a similar intake of water.^{16,24} The effect of oral fluid intake on peri-operative urine output should also be included in future studies.⁴²

CONCLUSIONS

Free intake of clear fluids, including a specially designed beverage for oral carbohydrate nutrition, up until 2 h prior to anaesthesia for elective surgery is safe and improves subjective well-being. The new Scandinavian guidelines emphasize that the minimum fasting time after intake of solids should still be 6 h. Fasting in emergency patients cannot secure gastric emptying and should not delay surgical interventions. More studies are needed on preoperative fasting and gastric content in patients with systemic disease, such as diabetes mellitus and patients with upper gastrointestinal symptoms. Overall, the choice of anaesthetic technique and airway management seems to be as important as adherence to any fasting guidelines when it comes to reducing the chance of pulmonary aspiration.

Nota de Autorización: Las anteriores Guías de Ayuno Preoperatorio: Actualización, se publican del original en Inglés aparecidas en la Revista ACTA ANESTHESIOLOGICA SCAND. 2005; Sep 49 (8): 1041-7, con la debida autorización del Prof. Eldan Søreide PhD, Presidente de la Sociedad Escandinava de Anestesiología y del Editor en Jefe, Sven Eric Gisvold de la Revista "Acta Anestesiologica Escandinava".

REFERENCES

1. Edwards G, Morton HJV, Pask EA, Wylie WD. Deaths associated with anaesthesia. A report on 1000 cases. *Anaesthesia* 1956; 11: 194-220.
2. Bannister WK, Sattilaro AJ. Vomiting and aspiration during anesthesia. *Anesthesiology* 1962; 23: 251-64.
3. Olsson GL, Hallen B, Hambraeus-Jonzon K. Aspiration during anaesthesia: computer-aided study of 185,358 anaesthetics. *Acta Anaesthesiol Scand* 1986; 30: 84-92.
4. Mellin-Olsen J, Fasting S, Gisvold SE. Routine preoperative gastric emptying is seldom indicated. A study of 85,594 anaesthetics with special focus on aspiration pneumonia. *Acta Anaesthesiol Scand* 1996; 40: 1184-8.
5. Warner MA, Warner ME, Weber JG. Clinical significance of pulmonary aspiration in the perioperative period. *Anesthesiology* 1993; 78: 56-62.
6. Søreide E, Bjørnstad E, Steen PA. An audit of Aspiration Pneumonitis in gynaecological and obstetric patients. *Acta Anaesthesiol Scand* 1996; 40: 14-9.
7. Kluger MT, Short TG. Aspiration during anaesthesia: a review of 133 cases from the Australian Anaesthesia Incident Monitoring Study (AIMS). *Anaesthesia* 1999; 54: 19-26.
8. McIntyre JW. Evolution of 20th century attitudes to prophylaxis of pulmonary aspiration during anaesthesia. *Can J Anaesth* 1998; 45: 1024-30.
9. Maltby JR, Sutherland AD, Sale JP, Shaffer EA. Preoperative oral fluids: is a five-hour fast justified prior to elective surgery? *Anesth Analg* 1986; 65: 1112-6.
10. Philips S, Hutchinson S, Davidson T. Preoperative drinking does not affect gastric contents. *Br J Anaesth* 1993; 70: 6-9.
11. Søreide E, Strømskag KE, Steen PA. Statistical aspects in studies of preoperative fluid intake and gastric content. *Acta Anaesthesiol Scand* 1995; 39: 738-43.
12. Strunin L. How long should patients fast before surgery? Time for new guidelines (Editorial). *Br J Anaesth* 1993; 70: 1-3.
13. Erikson LI, Sandin R. Fasting guidelines in different countries. *Acta Anaesthesiol Scand* 1996; 40: 971-4.
14. Søreide E, Fasting S, Ræder J. New preoperative fasting guidelines. *Acta Anaesthesiol Scand* 1997; 41: 799.
15. American Task Force on Preoperative Fasting. Practice guidelines for preoperative fasting and the use of pharmacological agents to reduce the risk of pulmonary aspiration: application to healthy patients undergoing elective procedures. A report by the American Task Force on Preoperative Fasting. *Anesthesiology* 1999; 90: 896-905.
16. Ljungqvist O, Søreide E. Preoperative fasting. *Br J Surg* 2003; 90: 400-6.
17. Spies CD, Breuer JP, Wichmann M, Adolph M, Senkal M, Kampa U et al. Preoperative fasting. An update. *Anaesthesist* 2003; 52: 1039-45.
18. Soop M, Nygren J, Myrenfors Thorell A, Ljungqvist O. Preoperative carbohydrate treatment attenuates immediate postoperative insulin resistance. *Am J Physiol* 2001; 280: E576-E583.
19. Hausel J, Nygren J, Lagerkranser M, Hellström PM, Hammarqvist F, Lindh A et al. A carbohydrate rich drink reduces preoperative discomfort in elective surgery patients. *Anesth Analg* 2001; 93: 1344-50.
20. Read NW, Houghton LA. Physiology of gastric emptying and pathophysiology of gastroparesis. *Gastroenterol Clin North Am* 1989; 18: 359-73.
21. Brener W, Hendrix TR, McHugh R. Regulation of the gastric emptying of glucose. *Gastroenterology* 1983; 85: 76-82.
22. Haavik P, Søreide E, Hofstad P, Steen PA. Does preoperative anxiety influence gastric fluid volume and acidity? *Anesth Analg* 1992; 75: 91-4.
23. Nygren J, Thorell A, Jacobsson H, Schnell PO, Ljungqvist O. Preoperative gastric emptying: the effects of anxiety and carbohydrate administration. *Ann Surg* 1995; 222: 728-34.
24. Petring OU, Blake DW. Gastric emptying in adults: an overview related to anaesthesia. *Anaesth Intensive Care* 1993; 21: 774-81.
25. Cote CJ. Preoperative preparation and premedication. *Br J Anaesth* 1999; 83: 16-28.
26. Litman RS, Wu CL, Quinlivan JK. Gastric volume and pH in infants fed clear liquids and breast milk prior to surgery. *Anesth Analg* 1994; 79: 407-9.
27. Horowitz M, O'Donovan D, Jones KL, Feinle C, Rayner CK, Samsom M. Gastric emptying in diabetes: clinical significance and treatment. *Diabet Med* 2002; 19: 177-94.
28. Scott AM, Kellow JE, Shuter B, Nolan JM, Hoschl R, Jones MP. Effects of cigarette smoking on solid and liquid intragastric distribution and gastric emptying. *Gastroenterology* 1993; 104: 410-6.
29. Wong PW, Kadakia SC, McBiles M. Acute effect of nicotine patch on gastric emptying of liquid and solid contents in healthy subjects. *Dig Dis Sci* 1999; 44: 2165-71. E. Søreide et al.
30. Søreide E, Veel T, Holst-Larsen H, Steen PA. The effects of chewing gum on gastric content prior to induction of anesthesia. *Anesth Analg* 1995; 80: 985-9.
31. Schumacher A, Vagts DA, Noldge-Schomburg GF. Smoking and preoperative fasting - are there evidence-based guidelines? *Anaesthesiol Reanim* 2003; 28: 88-96.
32. Pertwee RG. Cannabinoids and the gastrointestinal tract. *Gut* 2001; 48: 859-67.
33. Bujanda L. The effects of alcohol consumption upon the gastrointestinal tract. *Am J Gastroenterol* 2000; 95: 3374-82.
34. Brock-Utne JG, Moshal MG, Downing JW, Spitaels JM, Stiebel R. Fasting Volume and acidity of stomach contents associated with gastrointestinal symptoms. *Anaesthesia* 1977; 32: 749-52.
35. Hveem K, Hausken T, Berstad A. Ultrasonographic assessment of fasting liquid content in the human stomach. *Scan J Gastroenterol* 1994; 29: 786-9.
36. Stanghellini V, Tosetti C, Horowitz M, De Giorgio R, Barbara G, Cogliandro R et al. Predictors of gastroparesis in outpatients with secondary and idiopathic upper gastrointestinal symptoms. *Dig Liver Dis* 2003; 35: 389-96.
37. Maltby JR, Pytka S, Watson NC, Cowan RA, Fick GH. Drinking 300 mL of clear fluid two hours before surgery has no effect on gastric fluid Volume and pH in fasting

- and non-fasting obese patients. *Can J Anaesth* 2004; 51: 111-5.
38. Levy DM, Williams OA, Magides AD, Reilly CS. Gastric emptying is delayed at 8-12 weeks' gestation. *Br J Anaesth* 1994; 73: 237-8.
39. Whitehead EM, Smith M, Dean Y, O'Sullivan G. An evaluation of gastric emptying times in pregnancy and the puerperium. *Anaesthesia* 1993; 48: 53-7.
40. Olsson GL, Hallen B. Pharmacological evacuation of the stomach with metoclopramide. *Acta Anaesthesiol Scand* 1982; 26: 417-20.
41. Tryba M, Zenz M, Mlasowsky B, Huchzermeyer H. Does a stomach tube enhance regurgitation during general anaesthesia. *Anaesthesist* 1983; 32: 407-9.
42. Søreide E, Holst Larsen H, Reite K, Mikkelsen H, Søreide JA, Steen PA. The effects of giving 25-450 ml of water with diazepam premedication 1-2 hours before general anesthesia. *Br J Anaesth* 1993; 71: 503-6.
43. Vanner RG, Goodman NW. Gastro-oesophageal reflux in pregnancy at term and after delivery. *Anaesthesia* 1989; 44: 808-11.
44. Brady M, Kinn S, Stuart P. Preoperative fasting for adults to prevent perioperative complications. In: *Cochrane Database Syst Rev* 2003;(4). Hoboken: John Wiley & Sons, 2003.
45. Sackett DL, Straus SE, Richardson WS, Rosenberg W, Haynes RB. *Evidence-Based Medicine. How to Practice and Teach EBM*. New York: Churchill Livingstone, 2000.
46. Fasting S, Søreide E, Ræder J. Changing preoperative fasting policies. Does national consensus help? *Acta Anaesthesiol Scand* 1998; 42: 1188-92.
47. Søreide E, Hausken T, Søreide JA, Steen PA. Gastric emptying of a light hospital breakfast. A study using ultrasonography. *Acta Anaesthesiol Scand* 1996; 40: 549-53.
48. Brock-Utne JG. Clear fluids, not breakfast, before surgery. *Acta Anaesthesiol Scand* 1996; 40: 507-8.
49. Maltby JR, Koehli N, Ewen A, Shaffer EA. Gastric fluid volume, pH, and gastric emptying in elective inpatients. Influences of narcotic-atropine premedication, oral fluid, and ranitidine. *Can J Anaesth* 1988; 35: 562-6.
50. Scrutton MJ, Metcalfe GA, Lowy C, Seed PT, O'Sullivan G. Eating in labour. A randomised controlled trial assessing the risks and benefits. *Anaesthesia* 1999; 54: 1017-9.
51. Green SM, Krauss B. Pulmonary aspiration risk during emergency department procedural sedation-an examination of the role of fasting and sedation depth. *Acad Emerg Med* 2002; 9: 35-42.
52. The American society of Anesthesiologists-Task Force on Sedation and Analgesia by Non-Anesthesiologists. Practice Guidelines for Sedation and Analgesia by Non-Anesthesiologists (Approved by the House of Delegates on October 17, 2001) [www.document]. URL <http://www.asahq.org>. [accessed on 20 February 2005].
53. Steeds C, Mather SJ. Fasting regimens for regional ophthalmic anaesthesia. A survey of members of the British Ophthalmic Anaesthesia Society. *Anesthesia* 2001; 56: 638-42.
54. Maltby JR, Hamilton RC. Preoperative fasting guidelines for cataract surgery under regional anaesthesia. *Br J Anaesth* 1993; 71: 167.