



Revista Colombiana de Anestesiología

Colombian Journal of Anesthesiology

www.revcolanest.com.co



Case report

Jet ventilation for the extraction of endobronchial foreign body in a patient with drained pneumothorax: A case report[☆]



Julieta Latorre, Nicolas Brogly*, Genaro Maggi, Fabiola Quinteros, Natalia Peña del Ser, Fernando Gilsanz

Anesthesiology Department, Hospital Universitario La Paz, Madrid, Spain

ARTICLE INFO

Article history:

Received 24 August 2015

Accepted 25 November 2016

Available online 24 February 2017

Keywords:

Pneumothorax

Anesthesia, general

Foreign bodies

Respiration, artificial

Anesthesia

ABSTRACT

Introduction: Even though foreign body aspiration (FBA) is rare in adult patients, they sometimes require the performance of rigid bronchoscopy for its extraction. Ventilation for this procedure is challenging, especially in patients with pulmonary disease.

Clinical, diagnostic evaluation and interventions: We described the case of a 71-year old man who presented with a FBA in the left upper lobe associated with a contralateral pneumothorax. After the placement of a pleural drainage, the foreign body was extracted, using rigid bronchoscopy under general anesthesia and high flow jet ventilation with no further hemodynamic or pulmonary complications, suggesting that this technique is safe in patients with pulmonary leaks.

Conclusion: In the case described, rigid bronchoscopy minimized the risk of aspiration of blood and detritus due to continuous flow of gas upward, and avoided the increase of the air leak through the pneumothorax thank to limited airway pressures, making it a clear indication.

© 2016 Published by Elsevier España, S.L.U. on behalf of Sociedad Colombiana de Anestesiología y Reanimación. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Ventilación jet para la extracción de un cuerpo extraño endobronquial en un paciente con neumotórax drenado: informe de caso

RESUMEN

Introducción: Aunque la aspiración de cuerpos extraños es rara en adultos, requiere en algunas ocasiones la realización de una broncoscopia rígida para su extracción. El modo ventilatorio para este procedimiento es un reto, especialmente en pacientes con enfermedad pulmonar.

Palabras clave:

Neumotórax

Anestesia general

Cuerpos extraños

[☆] Please cite this article as: Latorre J, Brogly N, Maggi G, Quinteros F, Peña del Ser N, Gilsanz F. Ventilación jet para la extracción de un cuerpo extraño endobronquial en un paciente con neumotórax drenado: informe de caso. Rev Colomb Anestesiología. 2017;45:140-142.

* Corresponding author at: Servicio de Anestesia y Reanimación, Hospital Universitario La Paz, Paseo de la Castellana 261, 28046, Madrid, Spain.

E-mail address: nicolas0brogly@hotmail.com (N. Brogly).

2256-2087/© 2016 Published by Elsevier España, S.L.U. on behalf of Sociedad Colombiana de Anestesiología y Reanimación. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Respiración artificial
Anestesia

Clínica, evaluación diagnóstica e intervenciones: Describimos aquí el caso de un paciente de 71 años con aspiración de un cuerpo extraño enclavado a nivel del lóbulo superior izquierdo asociado a un neumotórax contralateral. Después de insertar un tubo de tórax se extrajo el cuerpo extraño utilizando broncoscopia rígida bajo anestesia general y ventilación jet de alto flujo sin complicaciones hemodinámicas o pulmonares adicionales, sugiriendo que esta técnica es segura en pacientes con fugas pulmonares.

Conclusiones: En el presente caso clínico, la broncoscopia rígida minimizó el riesgo de aspiración de sangre y detritus gracias a la inyección de un flujo continuo de gas ascendente, y evitó así el incremento de la fuga a través del neumotórax por las presiones limitadas en la vía aérea, constituyendo una indicación clara de esta técnica.

© 2016 Publicado por Elsevier España, S.L.U. en nombre de Sociedad Colombiana de Anestesiología y Reanimación. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

High frequency jet ventilation (HFJV) is a recognized ventilation technique for surgical and endoscopic interventions on the bronchial tree, in particular after a foreign body aspiration (FBA). Even though few clinical cases were reported about its use in adults,¹ we report the case of a patient with a FBA complicated with a pneumothorax who was ventilated with HFJV for the FBA extraction.

Patient information

We report a case of a 71-year old male with a history of COPD and pulmonary silicosis.

Clinical findings, diagnostic assessment and interventions

He was admitted to the emergency department of our hospital for sudden dyspnea after choking while eating, resulting in an increased work of breathing, and desaturation despite oxygen administration with reservoir mask.

A right chest tube was placed after the diagnosis of a pneumothorax on the chest X-ray and permitted to obtain a clinical improvement. Due to the persistence of an atelectasis of the left upper lobe, a chest CT scan was performed and allowed to identify a round image of 5 mm in relation with a foreign body, 3.8 mm distal to the carina, which occupied the left main bronchus (Fig. 1). Given the stable clinical situation, we decided to perform a rigid bronchoscopy to extract this foreign body under general anesthesia, using HFJV ($FiO_2 = 1$; pressure of 2 atm; $f = 150$ cycles/min, and I-time of 50%), once the patient had accomplished the fasting time of 6 h.

Follow-up and result

After an uneventful procedure, the patient did not present any complication, was transferred to the post anesthetic care unit, and discharged to the ward 2 h after.



Fig. 1 – Computer tomography showing a foreign body at 3.8 mm from the carina, a round image of about 5 mm occupying the left main bronchus that conditions atelectasis of the left upper lobe. Minimum right pneumothorax was observed. Source: authors.

Discussion of the medical literature

Tracheobronchial FBA is uncommon in adults, compared with children. The more common aspirated materials are dental debris, appliances, or prosthesis even though food debris are also described, especially in elderly patients and patients with neurological disorders.²

Rigid bronchoscopy is more invasive than flexible bronchoscope, but it gives more operability to the pneumologist or the thoracic surgeon and remains the gold standard when consistent mass of material is planned to be removed, since the working channels are significantly larger.³

HFJV has demonstrated its usefulness especially in neonates and pediatric patients, providing adequate gas exchange and minimizing risk of chronic respiratory diseases.

In adults, it is used for anesthesia in the upper airway surgery, ensuring the ventilation of the patient through a cannula inserted into the trachea, below the surgical field, while it improves the surgeon's comfort, suppressing the movement of the surgical field. In surgical resection of the descending thoracic aortic aneurysms, with selective lung ventilation, HFJV demonstrated its interest to ensuring the oxygenation in the non-ventilated lung. Also it has been used for lung resection, left atrial ablation by percutaneous radiofrequency, lithotripsy and laparoscopic surgery.⁴⁻⁶

It has been described as well for acute respiratory failure in patients with shock, applying small tidal volumes during high frequency ventilation: the lower intrathoracic pressure minimizes the interferences with venous return to the heart. In the presence of high flow bronchopleural fistula or pneumothorax, HFJV facilitates alveolar recruitment and increases the mean pressure in the airway, but decreases peak pressure and the leak. In severe respiratory failure, HFJV can maintain higher end- expiratory volumes, increasing functional residual capacity without increasing the maximum alveolar pressure. A recent publication showed that the association of HFJV with PEEP in rabbits suffering from acute lung injury improved oxygenation.⁷

Literature concerning the interest of HFJV in the management of patients with pneumothorax is scarce. HFJV is a known cause of pneumothorax due to barotrauma, and this is the reason why monitoring airway pressures during its use is so important. However, when there is a preexistent pneumothorax, the adequate use of HFJV probably helps to reduce the air leak through the bronchopleural fistula because a lower elevation in the airway pressure, associated with a good compromise on the oxygenation.^{8,9}

Lessons

In the case described, rigid bronchoscopy minimized the risk of aspiration of blood and detritus due to continuous flow of gas upward, and avoided the increase of the air leak through the pneumothorax thanks to limited airway pressures, making it a clear indication and utility.

Ethical disclosures

Protection of human and animal subjects. The authors declare that no experiments were performed on humans or animals for this study.

Confidentiality of data. The authors declare that they have followed the protocols of their work center on the publication of patient data.

Right to privacy and informed consent. The authors have obtained the written informed consent of the patients or

subjects mentioned in the article. The corresponding author is in possession of this document.

Funding

None.

Conflict of interests

The authors have no conflict of interests to declare.

REFERENCES

1. Gonzalez MCE, Ordoñez Fernández V. Reporte de caso: quemadura de la vía aérea. *Rev Colomb Anestesiología*. 2013;41:226-8.
2. Rafanan AL, Mehta AC. Adult airway foreign body removal. What's new? *Clin Chest Med*. 2001;22:319-30.
3. Liberman M. Bronchoscopic evaluation of the trachea and dilation of the trachea. *Semin Thorac Cardiovasc Surg*. 2009;21:255-62.
4. Chandradeva K, Palin C, Ghosh SM, Pinches SC. Percutaneous transtracheal jet ventilation as a guide to tracheal intubation in severe upper airway obstruction from supraglottic oedema. *Br J Anaesth*. 2005;94:683-6.
5. García Araque HF, Valencia Orgaz O, López Vicente R, Gutiérrez Vidal SE. Anatomía de la vía aérea para el broncoscopista. Una aproximación a la anestesia. *Rev Colomb Anestesiología*. 2014;42:192-8.
6. Ihra G, Gockner G, Kashanipour A, Aloy A. High-frequency jet ventilation in European and North American institutions: developments and clinical practice. *Eur J Anaesthesiol*. 2000;17:418-30.
7. Bang JO, Ha SI, Choi IC. The effect of positive-end expiratory pressure on oxygenation during high frequency jet ventilation and conventional mechanical ventilation in the rabbit model of acute lung injury. *Korean J Anesthesiol*. 2012;63:346-52.
8. Ellsbury DL, Klein JM, Segar JL. Optimization of high-frequency oscillatory ventilation for the treatment of experimental pneumothorax. *Crit Care Med*. 2002;30:1131-5.
9. Wippermann CF, Schranz D, Baum V, Huth R. Independent right lung high frequency and left lung conventional ventilation in the management of severe air leak during ARDS. *Paediatr Anaesth*. 1995;5:189-92.