



Revista Colombiana de Anestesiología

Colombian Journal of Anesthesiology

www.revcolanest.com.co



Review

Considerations in pediatric patients with obstructive sleep apnea/hypopnea syndrome (OSAHS): From physiopathology to the perioperative period[☆]



Álvaro Amézquita-Trujillo, Juan Federico Garzón*

Fundación Hospital de la Misericordia – HOMI, Bogotá D.C., Colombia

ARTICLE INFO

Article history:

Received 18 May 2016
Accepted 15 March 2017
Available online 20 June 2017

Keywords:

Sleep apnea syndromes
Perioperative period
Child
Anesthesia
Physiopathology

ABSTRACT

Introduction: OSAHS is a respiratory disorder with perioperative complications, which increase morbidity and mortality. The characteristics of the pediatric population require a careful perioperative assessment.

Objective: This article addresses relevant physiopathology, clinical and paraclinical topics, perioperative treatment and care of pediatric patients with OSAHS.

Materials and methods: A literature search was conducted in PubMed and Embase databases using the MeSH terms: sleep apnea syndromes, obstructive sleep apnea, physiopathology, perioperative period, and perioperative care. The titles and/or the abstracts of each of the obtained results were reviewed and 60 articles were finally selected.

Results and conclusion: The knowledge of the physiopathology, the adequate preanesthetic assessment and the pertinent request of paraclinical care, allows for the adjustment of the perioperative care and to improve the results of the patient.

© 2017 Sociedad Colombiana de Anestesiología y Reanimación. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Consideraciones en el paciente pediátrico con síndrome de apnea/hipopnea obstructiva del sueño (SAHOS): desde la fisiopatología al perioperatorio

RESUMEN

Introducción: El SAHOS, es un trastorno respiratorio con complicaciones perioperatorias, que aumentan la morbimortalidad; las características de la población pediátrica hacen necesaria una cuidadosa valoración perioperatoria.

Palabras clave:

Síndrome de la apnea del sueño
Periodo perioperatorio

[☆] Please cite this article as: Amézquita-Trujillo Á, Garzón JF. Consideraciones en el paciente pediátrico con síndrome de apnea/hipopnea obstructiva del sueño (SAHOS): desde la fisiopatología al perioperatorio. Rev Colomb Anestesiolog. 2017;45:173-181.

* Corresponding author at: Av. Caracas No. 1-13, Fundación Hospital de la Misericordia, Salas de Cirugía, Oficina de Anestesia, Bogotá, Colombia.

E-mail address: andolinicor@gmail.com (J.F. Garzón).

2256-2087/© 2017 Sociedad Colombiana de Anestesiología y Reanimación. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Niño
Anestesia
Fisiopatología

Objetivo: Este artículo aborda tópicos relevantes de la fisiopatología, la clínica, paraclínicos, tratamientos y cuidados perioperatorios de los pacientes pediátricos con SAHOS.

Materiales y métodos: Se realizó una búsqueda de literatura, en base de datos de Pubmed y Embase, con los términos MESH: Syndromes sleep apnea, obstructive sleep apnea, Physiopathology, Perioperative Period, Perioperative Care. Se revisaron los títulos y/o resúmenes de cada uno de los resultados obtenidos y se escogieron finalmente 60 artículos.

Resultados y conclusión: El conocimiento de la fisiopatología, la adecuada valoración preanestésica, la pertinente solicitud de paraclínicos, permite establecer el ajuste de los cuidados perioperatorios y mejorar los resultados del paciente.

© 2017 Sociedad Colombiana de Anestesiología y Reanimación. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Obstructive sleep apnea/hypopnea syndrome (OSAHS) is part of a group of related diseases, known as sleep-disordered breathing (SDB), which includes: primary snoring (PS), upper airway resistance syndrome (UARS) and OSAHS. The latter is the most severe form and it has a higher risk of perioperative complications and systemic repercussions that increase morbidity.¹⁻⁴

The American Thoracic Society and the International Classification of Sleep Disorders¹ defines OSAHS as “a sleep-disordered breathing characterized by partial or complete, prolonged and intermittent obstruction of the upper airway that alters normal ventilation during sleep and normal sleep patterns”.¹

The episodes are characterized by snoring, desaturation, hypercapnia, respiratory effort, changes in intrathoracic pressure followed by cortical and subcortical activation with intermittent return to normal sleep patterns. Its prevalence ranges from 1% to 5% and occurs mainly between 2 and 8 years of age.⁵⁻⁸

The aim of the present review is to show the approach toward the pediatric patient with OSAHS from the pathophysiological perspective, considering relevant epidemiological data, addressing the preanesthetic assessment (clinical and paraclinical) and the recommendations about preoperative treatment.

Methodology

A search of the literature was performed with the MeSH terms: sleep apnea syndromes, obstructive sleep apnea, physiopathology, perioperative period and perioperative care in the Pubmed (207 articles) and Embase (33 articles) databases, finding a total of 240 articles of which titles and abstracts were reviewed. Finally, 60 articles were selected according to the inclusion of relevant topics in the pediatric population (see Fig. 1).

The results of the reading of the selected articles are presented as a summary in subtopics such as: pathophysiology, morbidity, epidemiology, preanesthetic assessment and preoperative treatment. Also, conclusions were drawn from the authors' interpretation of the articles.




Fig. 1 – Literature search.

Source: Authors.

Physiopatology

Permeability of pharyngeal airway

There are two described models for understanding the phenomena that determine the airway permeability: The neural model and the anatomical model.

Neural model

The size of the pharyngeal airway is determined by the balance between two opposing forces. The ones that collapse the pharynx are produced by the contraction of the inspiratory muscles (diaphragm and external intercostals) and generate a negative pressure gradient within the pharynx, which promotes the collapse of the extrathoracic airway.⁹⁻¹¹ This collapse is prevented by forces of the pharynx produced by the contraction of the dilator muscles. The activity of these two forces is regulated by neural mechanisms (chemical stimuli, state of consciousness, airway reflexes) that interact to maintain the airway permeable. Neural mechanisms work best during waking states. When the level of consciousness is reduced, the activity of the pharyngeal dilator muscles decreases rather than the activity of the inspiratory muscles, resulting in a loss of balance of the neural balance favoring the narrowing of the pharyngeal airway.^{10,11}

Anatomical model

Proposed by Isono,¹² it describes that the upper airway passes through a compartment within which is the pharyngeal soft tissue. This, at the same time, is limited by several bone structures such as the base of skull, cervical spine and mandible.^{10,11,13} During sleep, airway caliber is determined by soft tissue, bone structures, and by the tone of the pharyngeal dilator muscles (genioglossus, geniohyoid, and velopalatine muscles).^{10,11,13} During the contraction of the inspiratory muscles forces are generated that collapse the airway by the intraluminal negative pressure. Usually, the pharyngeal airway does not collapse until sub-atmospheric pressures are reached. In patients with OSAHS, collapse of the pharyngeal airway occurs at a positive pressure.¹²

The balance between the two forces (closure tendency, opening tendency) determines the anatomical balance of the pharyngeal airway. Its size is conditioned by the interaction between neural and anatomical mechanisms.^{10,14-16}

Increased upper airway resistance during sleep is an essential feature of OSAHS. In children, it is usually a multifactorial disorder in which various factors converge, including: stenosis of the upper respiratory tract (e.g., tonsillar hypertrophy), abnormal airway muscle tone, and genetic predisposition.¹⁷

Physiopathological factors involved can be divided into:

Craniofacial anatomical factors: Small mandible, large or retroposed tongue, enlarged pharyngeal fat deposit, hypertrophic lymphoid tissues in the upper respiratory tract (particularly adenoids and tonsils), inferior nasal turbinate hypertrophy, deviated nasal septum.

Factors that promote greater collapsibility: Inflammation of the upper airway and alteration of the neurological reflexes that control upper respiratory tract muscles.^{10,18}

Obstructive sleep events in children with OSAHS have an inverse pattern in relation to the sleep phase, compared with that observed in adults with OSAHS. 80% of obstructive events occur during REM sleep, while in adults 80% of obstructive events occur during non-REM sleep. Non-REM obstructive events occur less frequently in children, but they tend to increase in older children, African-American children, those in a lateral position, and those with low levels of oxyhemoglobin saturation. Non-REM obstructive events leading to wakefulness occur at a higher oxygen saturation than events during REM sleep.^{16,19}

OSAHS can be divided into two types: Type I associated with marked lymphadenoid hypertrophy in the absence of obesity and Type II associated with obesity and with mild hyperplasia of lymphadenoid tissue.⁶

The cross-sectional area of the pharynx in children with OSAHS is 5%–10% smaller than in healthy children. Breathing in an awake child with this pathology is normal. During general anesthesia and neuromuscular block, the upper airway decreases in caliber. The relative increase in the soft pharyngeal tissue is caused by inhibition of the dilator muscles of the airway and decrease in lung volume, which results in a cephalic displacement of the mediastinum and a decrease in longitudinal tension on the upper airway, increasing collapsibility of the pharyngeal airway.^{12,20}

Morbidity

OSAHS has systemic repercussions that increase the morbidity of patients who suffer from it.^{3,6} Alterations in behavior have been observed in these patients (hyperactivity, attention deficit, cognitive deficit and poor school performance), along with daytime sleepiness, cardiovascular alterations, metabolic alterations, alterations in somatic growth, metabolic syndrome.²¹

OSAHS and obesity are considered inflammatory diseases. An increase in C-reactive protein has been observed that could constitute an important marker in the diagnosis of OSAHS. The current evidence suggests the existence of a genetic predisposition demonstrated by apolipoprotein epsilon 4.^{22,23} Children with OSAHS, particularly the obese ones, tend to develop metabolic syndrome and a fatty liver.^{6,24,25}

Cardiovascular morbidity may be attributable to a complex interaction between intermittent hypoxia, episodic hypercapnia, recurrent fluctuations of intrathoracic pressure and sleep interruption. OSAHS can affect endothelial function causing hypertension due to increased release of endothelin and decreased nitric oxide production; In addition, sympathetic activation may exert direct action on the endothelium causing vasoconstriction. Inflammation may contribute to endothelial dysfunction by increasing the adhesion of inflammatory mediators and generating hypercoagulability. Some studies have shown that these alterations can be reversed 4 to 6 weeks after tonsillectomy was performed.^{22,26,27}

In some patients with OSAHS, delayed growth and development have been observed, which may be secondary to an alteration in the functioning and/or release of insulin-like growth factor 1 and growth hormone.⁶

Epidemiology

In 2008, Lumeng and Chervin carried out a systematic review of studies on the epidemiology of sleep-disordered breathing (SDB), finding the following prevalences: SDBs reported by parents 4%–11%; habitual snoring 1.5%–12%; sleep apnea events 0.2%–4%. OSAHS diagnosed by different criteria in diagnostic studies varies widely between 0.1 and 13%, but most studies report a figure between 1% and 4%.^{2,13,25} In general, SDB are more common in males, overweight children and African-American children. There are no data that support clear differences in age-based prevalence. However, some data shows a peak prevalence between 2 and 8 years of age.^{11,18,28}

In children younger than 2 years, OSAHS is frequently associated with facial/skull malformations, neurological abnormalities and obesity. In these cases, OSAHS is usually more severe and with a greater recurrence or incomplete recovery after surgery.²⁹

The incidence of postoperative respiratory complications in patients with OSAHS reaches 27% (6.4–27%), which varies according to age, severity of OSAHS, diagnostic criteria used and comorbidities. Children <3 years and an AHI (Apnea/Hypopnea Index) >10 have twice as much risk as children between 3 and 6 years of age.

These complications include oxygen desaturation <90%, increased respiratory effort, radiographic changes (edema, atelectasis, infiltrates, pneumothorax, pneumomediastinum, pleural effusion), laryngospasm, apnea, crisis of pulmonary hypertension, pneumonia and perioperative death, and negative pressure pulmonary edema.¹⁷ Respiratory complications may be related to an alteration in the response to hypoxemia and hypercapnia and overexpression of opioid receptors.^{22,30-32}

Preanesthetic assessment

In all patients, a complete medical history including a review of systems, emphasizing clinical risk factors, and a specific physical examination should be performed. The child with OSAHS is at risk for potentially fatal perioperative respiratory complications, so the perioperative approach must be multidisciplinary (pediatrics, surgery, anesthesia).^{14,33,34}

Most of the complications are respiratory. Children with OSAHS have a 20% greater chance of complications compared to healthy children undergoing adenotonsillectomy. In the general population, 21% of complications occur in the intraoperative period, 33% in the postanesthetic care unit and 46% in the patient's room. These complications can lead to severe neurological damage or death.³⁵⁻³⁷ Other less frequent complications such as negative pressure pulmonary edema and bleeding have been observed.²²

The majority of the literature focuses on the perioperative management of the child with OSAHS who will be taken to adenotonsillectomy, from which the management principles for other surgical interventions have been extrapolated.¹⁴ General principles of management include:

- Preoperative evaluation of snoring.
- Strict observation and respiratory monitoring.
- Analgesic and anesthetic opioid sparing techniques.³⁰
- Regional anesthesia techniques when possible.¹¹

The clinical presentation varies depending on the age. In childhood, there should be high suspicion of craniofacial dysmorphism, growth retardation, susceptibility to infections or delayed speech development. At school age it can manifest as agitation, attention deficit, poor school performance or secondary enuresis. In adolescence the clinical presentation becomes similar to the adult.^{14,28,33,38}

In the absence of characteristic symptoms of OSAHS in the pediatric age, the situation can be clarified with the systematic application of screening questions, emphasizing the presentation of nocturnal symptoms (snoring, excessive sweating, restless sleep, mouth breathing, apneas, wheezing, paradoxical or labored breathing, hyperextension of the neck during sleep). Also with the presence of diurnal symptoms such as difficulty concentrating, behavioral and mood disorders, morning headaches, excessive daytime sleepiness, and delayed growth.³⁹ It should be questioned directly by the presence of snoring (most children with OSAHS snore, and the absence of snoring makes a diagnosis of OSAHS much less likely, with a sensitivity of 91% and a specificity of 75%).^{17,27,33}

Table 1 – OSAHS severity in children and adults defined by The American Society's Working Group on Perioperative Management of Patients with Obstructive Sleep Apnea.

OSAHS severity	AIH children	AIH adults	SaO ₂ nadir
None	0	0-5	>92%
Mild	1-5	6-20	
Moderate	6-10	21-40	
Severe	>10	>40	<80%

Source: Modified and adapted from Schwengel et al.¹⁷ and Gross et al.⁴⁵

On physical examination, the findings suggestive of OSAHS include: mouth breathing, elongated facies, triangular and/or small chin, retroposition of the mandible, high palate, chest retractions, extreme body habits (developmental delay in infants, and obesity in children and adolescents), developmental delay. The magnitude of the tonsillar hypertrophy is not correlated with the severity of OSAHS.^{17,33}

Questionnaires and clinical history are useful but not adequate in clinical practice to differentiate primary snoring in children from pediatric OSAHS (normal polysomnographic studies); therefore, additional diagnostic testing is required. The accuracy of the clinical evaluation of pediatric OSAHS in predicting the positivity of sleep studies varies between 30% and 85%.¹⁴

Polysomnography performed in a sleep laboratory at night is considered the gold standard for the diagnosis and evaluation of OSAHS in children. Ideally, this test should include cardiorespiratory, electroencephalographic, electrooculographic and electromyographic monitoring.^{1,17,40} Polysomnography is indicated in patients at high risk and in the evaluation of the following clinical conditions: growth failure, unexplained polycythemia (particularly if it is associated with snoring), sickle cell disease associated with a medical history suggestive of OSAHS or frequent veno-occlusive crisis, morbid obesity, craniofacial anomalies, neuromuscular disorders, cor pulmonale, systemic arterial hypertension.^{14,41}

In polysomnography the following measurements are defined.^{17,18,33}

Apnea-Hypopnea Index (AHI)

Mean number of obstructive events (apnea-hypopnea) per hour of sleep. Diagnostic criteria for OSAHS in adults are a product of expert consensus and include an AHI of 5 or more and evidence of restless or disturbed sleep, daytime sleepiness, or other diurnal symptoms. IAH cut-off points of 5, 15 and 30 events/hour have been suggested to indicate mild, moderate and severe levels.^{14,42} There are few data available in the literature to establish diagnostic criteria for specific OSAHS in children from AHI. Currently, an AHI of 1-5 events per hour is used in research to identify children with OSAHS; however, widely varying definitions have been used^{1,33} (see Table 1).

Oxygen saturation nadir (see Table 1)

Respiratory Disturbance Index (RDI): Number of apnea/hypopnea events per hour of sleep. It is not specific for OSAHS, since it includes episodes of central and obstructive apnea.¹⁸

Due to the cost and the limited availability in many communities of polysomnography performed in the sleep laboratory, significant efforts have been made to explore other forms of diagnosis. Recording of audio at home at night, video recording, pulse oximetry, cardiorespiratory studies, and home polysomnography have been used. However, the validity of each of these methods has not been compared with the gold standard in the pediatric population.⁴³ It cannot be ruled out that any of these modalities may have some diagnostic value in specific populations. Brouillette et al.⁴⁴ concluded that in children with suspected OSAHS, severe isolated oxygen desaturation (SaO₂ <80%) or desaturation clusters (more than three episodes of SaO₂ <90%) are considered abnormal and have a positive predictive value of 97%, but a negative result does not rule out the diagnosis of OSAHS.

During the evaluation of the child with OSAHS, the airway obstruction site should be identified. The study of dynamic airway collapse patterns during sleep identifies the anatomical causes of the obstruction and facilitates the planning of interventions to alleviate such obstruction. To study these, differential pharyngeal pressures through catheters placed at different levels in the upper airway have been used; cine-fluoroscopy; video endoscopy; computed tomography; fiberoptic bronchoscopy.^{14,33}

Although most children scheduled for adenotonsillectomy do not require cardiac evaluation, patients with evidence of systemic hypertension and/or chronic hypercapnia (multiple episodes of severe hypoxemia <70%, compensatory metabolic alkalosis), which usually have increased pulmonary pressures and therefore are at risk of cor pulmonale, should have at least one evaluation with echocardiogram before proceeding with elective surgery. These cardiopulmonary abnormalities are seen less frequently in children than in adults.^{24,26,46}

Children at high risk for postoperative respiratory complications should be identified from preanesthetic assessment to plan their postoperative stay. Although there is no clear evidence of relevant observation time, there are some recommendations drawn from the literature that may contribute to the adequate planning of the perioperative treatment of the patient and to reduce the incidence of complications through appropriate postoperative monitoring (see Table 2).^{14,17,26,47}

Mild or moderate OSAHS without any risk factors: 3-7 h observation

Moderate/mild OSAHS without risk factors but with desaturation, oxygen requirement, difficult intraoperative airway management or postoperative obstruction: intermediate care for 24 h.

Mild/moderate OSAHS without risk factors but with severe pain requiring opioid management: intermediate care for 24 h.

Mild or moderate OSAHS with risk factors: intermediate care for 24 h.

Severe OSAHS: intensive care for 24 h (see Fig. 2).

Table 2 – Risk factors for pulmonary postoperative complications after adenotonsillectomy.

Children < 3 years old
Severe OASHS documented by polysomnography
Failure to thrive (FTT)
Obesity
Cardiac compromise (cor pulmonale, RVH, high blood pressure)
Down's syndrome
History of prematurity
Craniofacial abnormalities
Neuromuscular diseases
Chronic lung disease
Sickle cell disease
Nasal problems (septal deviation, turbinate hypertrophy)
Adenoid hypertrophy
Mallampati 3-4
Upper respiratory infection in the previous 4 weeks
Marked obstruction with inhalation induction
Difficulty breathing in the recovery room

Source: Adapted and modified from Brown,³ Patino et al.,¹⁴ Schwen-
gel et al.,¹⁷ Den Herder et al.,⁵⁹ Brown and Brouillette.⁶⁰

Perioperative treatment

Adenotonsillectomy is the recommended initial treatment for pediatric OSAHS.^{34,48} Positive-pressure ventilation (BiPAP/CPAP) in the preoperative period has been used to decrease the rate of postoperative complications and to increase airway permeability in adults. It may be useful in pediatric patients with severe OSAHS and secondary cardiovascular complications,^{9,17,26,49} since improvement in pulmonary hypertension and reduction of surgical risk have been observed. The pressures adjusted by the pediatric pulmonologist in the sleep laboratory are those used in the postoperative period.^{17,50,51}

Several studies compare early adenotonsillectomy with expectant management in different pediatric ages of patients with OSAHS and other lung diseases, finding that it does not significantly improve care or executive function measured by neuropsychological tests, but reduces symptoms and improves secondary outcomes of behavior, quality of life and polysomnography.⁵¹⁻⁵⁶

The following are evidence-based practice recommendations for the diagnosis and treatment of OSAHS^{57,58}:

Clinical history and physical examination are not sufficient to differentiate between primary snoring and OSAHS. Polysomnography is the gold standard for the diagnosis of OSAHS.

Children with Down's syndrome, craniofacial anomalies, obesity, or Prader-Willi syndrome (PWS) are at increased risk for OSAHS and often require polysomnography and specific therapies.

The pillars of therapy for OSAHS in children are adenotonsillectomy and the use of positive airway pressure devices.⁹ The use of continuous positive airway pressure therapy (CPAP) is an effective treatment in children. However, like what occurs in adults, adherence is a major problem and there may be side effects in children with long-term use.⁵⁰

Non-surgical therapies, including anti-inflammatory drugs, dental appliances and weight loss may be helpful in




Fig. 2 – Algorithm for risk assessment and postoperative disposition of the pediatric patient with OSAHS.

Source: Adapted and modified from Patino et al.¹⁴ and Schwengel et al.¹⁷

certain circumstances, such as mild OSAHS, either primary or persistent after adenotonsillectomy, although more data is needed.⁹

Airway surgery has been shown to be effective in children with craniofacial abnormalities, but not in the general pediatric population. Bariatric surgery is an option for extremely obese adolescents.

As for premedication, caution should be exercised with sedatives in children with severe OSAHS. If they have been received, they should be continuously monitored by clinical observation and, at least, continuous pulse oximetry until the transfer to surgical rooms. The residual effect of these drugs can be observed even in the recovery stage, especially in short-term procedures.¹⁴ Some data suggest that children with OSAHS may receive sedatives, but require monitoring until full recovery can be demonstrated.¹⁷

The strategies to keep the pharyngeal airway permeable consist of: Expansion of the bone limits, reduction of pharyngeal soft tissue volume and activation of the pharyngeal dilator muscles.¹² The expansion of the bone limit can be obtained with the triple airway maneuver. The reduction of the volume of the pharyngeal soft tissue can be obtained with the use of continuous positive airway pressure (CPAP) that acts like a pneumatic splint preventing the soft tissue from invading the upper airway. It also increases lung volume, resulting in a caudal displacement of the mediastinum that applies longitudinal tension over the pharyngeal airway, decreasing its collapse.^{12,50} Recruitment of the upper airway dilator muscles is not easy due to the selective inhibition of many anesthetics. Propofol apparently preserves genioglossal muscle activity.¹²

The position of the body influences the volume of the soft pharyngeal tissue. In the supine position, airway occlusion in children with OSAHS occurs in the upper 2/3 of the pharyngeal airway in the area between the adenoid tissue and the tonsils. The lateral position displaces the soft tissue outside the airway.^{12,26}

Conclusions

OSAHS is a disease whose prevalence in the pediatric population requires an in-depth knowledge of its physiopathology and the comorbidities that it involves at the pulmonary, cardiovascular, neurological, metabolic and hematological levels. Similarly, the approach in preanesthetic assessment should focus on a broad clinical evaluation, considering polysomnography and derived information as a fundamental diagnostic tool, that allows to determine the pertinence of a broader paraclinical evaluation to define the parameters of preoperative treatment and postoperative care in order to reduce complications.

Funding

The authors did not receive sponsorship to carry out this article.

Conflicts of interest

The authors have no conflicts of interest to declare.

REFERENCES

1. American Academy of Sleep Medicine. International classification of sleep disorders: diagnostic and coding manual. 2nd ed. American Academy of Sleep Medicine; 2005.
2. Lumeng JC, Chervin RD. Epidemiology of pediatric obstructive sleep apnea. *Proc Am Thorac Soc.* 2008;5:242-52.
3. Brown KA. Outcome, risk, and error and the child with obstructive sleep apnea. *Paediatr Anaesth.* 2011;21:771-80.
4. Tait AR, Voepel-Lewis T, O'Brien LM. Postsurgical behaviors in children with and without symptoms of sleep-disordered breathing. *Perioper Med.* 2014;3:8.
5. Lerman J. A disquisition on sleep-disordered breathing in children. *Paediatr Anaesth.* 2009;19 Suppl. 1:100-8.
6. Capdevila OS, Kheirandish-Gozal L, Dayyat E, Gozal D. Pediatric obstructive sleep apnea: complications, management, and long-term outcomes. *Proc Am Thorac Soc.* 2008;5:274-82.
7. Farag HM, Bondok RS, Elkader HAA. *Anesth Protocol Liver Transplant.* 2012;5:5-7.
8. American Society of Anesthesiologists. Practice guidelines for the perioperative management of patients with obstructive sleep apnea. *Anesthesiology.* 2014;120:268-86.
9. Kaditis A, Kheirandish-Gozal L, Gozal D. Algorithm for the diagnosis and treatment of pediatric OSA: a proposal of two pediatric sleep centers. *Sleep Med.* 2012;13:217-27.
10. Katz ES, D'Ambrosio CM. Pathophysiology of pediatric obstructive sleep apnea. *Proc Am Thorac Soc.* 2008;5:253-62.
11. Schwengel D, Dalesio NM, Stierer TL. Pediatric obstructive sleep apnea. *Anesthesiol Clin.* 2014;32:237-61.
12. Isono S. Developmental changes of pharyngeal airway patency: implications for pediatric anesthesia. *Paediatr Anaesth.* 2006;16:109-22.
13. Karlson KH. What's new in pediatric obstructive sleep apnea? *Clin Pulm Med.* 2008;15:226-31.
14. Patino M, Sadhasivam S, Mahmoud M, Hemmings HC. Obstructive sleep apnoea in children: perioperative considerations. *Br J Anaesth.* 2013;111 Suppl. 1:83-95.
15. MacHata AM, Kabon B, Willschke H, Prayer D, Marhofer P. Upper airway size and configuration during propofol-based sedation for magnetic resonance imaging: an analysis of 138 infants and children. *Paediatr Anaesth.* 2010;20:994-1000.
16. Vasu TS, Grewal R, Doghramji K. Obstructive sleep apnea syndrome and perioperative complications: a systematic review of the literature. *J Clin Sleep Med.* 2012;8:199-207.
17. Schwengel D, Sterni LM, Tunkel DE, Heitmiller ES. Perioperative management of children with obstructive sleep apnea. *Anesth Analg.* 2009;109:60-75.
18. Kheirandish-Gozal L, Gozal D. Sleep disordered breathing in children. A comprehensive clinical guide to evaluation and treatment. Springer Science; 2012. p. 13-24.
19. Spruyt K, Gozal D. REM and NREM sleep-state distribution of respiratory events in habitually snoring school-aged community children. *Sleep Med.* 2012;13:178-84.
20. Eikermann M, Martina Grosse-Sundrup SZ, Henry ME, Bittner EA, Hoffmann U, Chamberlin NL. Ketamine activates breathing and abolishes the coupling between loss of consciousness and upper airway dilator muscle dysfunction. *Anesthesiology.* 2012;116:35-46.

21. Bonsignore MR, McNicholas WT, Montserrat JM, Eckel J. Adipose tissue in obesity and obstructive sleep apnoea. *Eur Respir J*. 2012;39:746-67.
22. Kwok K, Ng DK, Chan C. Cardiovascular changes in children with snoring and obstructive sleep apnoea. *Ann Acad Med Singapore*. 2008;37:715-21.
23. Chou CH, Kang KT, Weng WC, Lee PL, Hsu WC. Central sleep apnea in obese children with sleep-disordered breathing. *Int J Obes (Lond)*. 2014;38:27-31.
24. Mitchell RB, Boss EF. Pediatric obstructive sleep apnea in obese and normal-weight children: impact of adenotonsillectomy on quality-of-life and behavior. *Dev Neuropsychol*. 2009;34:650-61.
25. Suárez C, Figueroa J, Len F, Mansilla E. Monitoreo grabado de oximetría de pulso en niños con hipertrofia adenoidea/amigdalina: su utilidad en el diagnóstico del síndrome de apnea obstructiva del sueño. *Arch Argent Pediatr*. 2013;111:196-201.
26. Robb P, Bew S, Kubba H, Murphy N, Primhak R, Rollin A, et al. Tonsillectomy and adenoidectomy in children with sleep-related breathing disorders: consensus statement of a UK multidisciplinary working party. *Ann R Coll Surg Engl*. 2009;91:371-3.
27. Brown KA. Evaluation of the child with sleep-disordered breathing scheduled for adenotonsillectomy. *Sleep Med Clin*. 2013;8:149-55.
28. Abdel-Aziz M, Ibrahim N, Ahmed A, El-Hamamsy M, Abdel-Khalik MI, El-Hoshy H. Lingual tonsils hypertrophy; a cause of obstructive sleep apnea in children after adenotonsillectomy: operative problems and management. *Int J Pediatr Otorhinolaryngol*. 2011;75:1127-31.
29. Madureira N. Obstructive sleep apnea syndrome in children younger than 2 years of age. *Sleep Med*. 2013;14:e193.
30. Raghavendran S, Bagry H, Detheux G, Zhang X, Brouillette RT, Brown Ka. An anesthetic management protocol to decrease respiratory complications after adenotonsillectomy in children with severe sleep apnea. *Anesth Analg*. 2010;110:1093-101.
31. Brown Ka, Laferrrière A, Moss IR. Recurrent hypoxemia in young children with obstructive sleep apnea is associated with reduced opioid requirement for analgesia. *Anesthesiology*. 2004;100:806-10, discussion 5A.
32. Lam KK, Kunder S, Wong J, Doufas AG, Chung F. Obstructive sleep apnea, pain, and opioids: is the riddle solved? *Curr Opin Anaesthesiol*. 2016;29:134-40.
33. Brooks LJ. Diagnosis and evaluation of obstructive sleep apnoea in children. *Ann Acad Med Singapore*. 2008;37:701-5.
34. Isaiah A, Pereira KD. Outcomes after adenotonsillectomy using a fixed anesthesia protocol in children with obstructive sleep apnea. *Int J Pediatr Otorhinolaryngol*. 2015;79:638-43.
35. Coté CJ, Posner KL, Domino KB. Death or neurologic injury after tonsillectomy in children with a focus on obstructive sleep apnea: Houston, we have a problem! *Anesth Analg*. 2014;118:1276-83.
36. Fouladpour N, Jesudoss R, Bolden N, Shaman Z, Auckley D. Perioperative complications in obstructive sleep apnea patients undergoing surgery: a review of the legal literature. *Anesth Analg*. 2016;122:145-51.
37. Benumof JL. Mismanagement of obstructive sleep apnea may result in finding these patients dead in bed. *Can J Anesth*. 2016;63:3-7.
38. Scholz J, Yaksh TL. Preclinical research on persistent postsurgical pain: what we don't know, but should start studying. *Anesthesiology*. 2010;112:511-3.
39. Tan H-L, Gozal D, Kheirandish-Gozal L. Obstructive sleep apnea in children: a critical update. *Nat Sci Sleep*. 2013;5:109-23.
40. Nolan JA. Principles of paediatric anaesthesia. *Anaesth Intensive Care Med*. 2016;17:265-9.
41. Hsu W-C, Kang K-T, Weng W-C, Lee P-L. Impacts of body weight after surgery for obstructive sleep apnea in children. *Int J Obes (Lond)*. 2013;37:527-31.
42. American Academy of Sleep Medicine Task Force. Sleep-related breathing disorders in adults: recommendations for syndrome definition and measurement techniques in clinical research. *Sleep*. 1999;22:667-89.
43. Pediatric Pulmonology Subcommittee on Obstructive Sleep Apnea Syndrome. American Academy of Pediatrics, clinical practice guideline: diagnosis and management of childhood obstructive sleep apnea syndrome. *Pediatrics*. 2002;109:704-71.
44. Brouillette R, Morielli A, Leimani A, Waters K, Luciano R, Ducharme FM. Nocturnal pulse oximetry as an abbreviated testing modality for pediatric obstructive sleep apnea. *Pediatrics*. 2000;105:405-12.
45. Gross J, Bachenberg K, Benumof JL, Al E. Practice guidelines for the perioperative management of patients with obstructive sleep apnea: a report by the American Society of Anesthesiologists Task Force on perioperative management of patients with obstructive sleep apnea. *Anesthesiology*. 2006;104:1081-93.
46. Huang Y-S, Guilleminault C, Lee L-A, Lin C-H, Hwang F-M. Treatment outcomes of adenotonsillectomy for children with obstructive sleep apnea: a prospective longitudinal study. *Sleep*. 2014;37:71-7.
47. Whippey A, Kostandoff G, Ma HK, Cheng J, Thabane L, Paul J, et al. Predictors of unanticipated admission following ambulatory surgery in the pediatric population: a retrospective case-control study. *Paediatr Anaesth*. 2016;26:831-7.
48. Setabutr D, Adil Ea, Chaikhoutdinov I, Carr MM. Impact of the pediatric tonsillectomy and polysomnography clinical practice guidelines. *Int J Pediatr Otorhinolaryngol*. 2014;78:517-21.
49. Cox RG. Anesthetic management of pediatric adenotonsillectomy. *Can J Anesth*. 2007;54:1021-3.
50. Sawyer AM, Gooneratne N, Marcus CL, Ofer D, Richards KC, Weaver T. A systematic review of CPAP adherence across age groups: clinical and empiric insights for developing CPAP adherence interventions. *Sleep Med Rev*. 2012;29:997-1003.
51. Ng DK, Wong JC, Chan C-H, Leung LCK, Leung S-Y. Ambulatory blood pressure before and after adenotonsillectomy in children with obstructive sleep apnea. *Sleep Med*. 2010;11:721-5.
52. Marcus C, Moore R, Rosen CL. A randomized trial of adenotonsillectomy for childhood sleep apnea. *N Engl J Med*. 2013;368:2366.
53. Busino RS, Quraishi H, Aguila H, Montalvo E, Connelly P. The impact of adenotonsillectomy on asthma in children. *Laryngoscope*. 2010;120 Suppl. 4:2007.
54. Levin J, Gagnon L, He X, Baum ED, Karas DE, Chupp GL. Improvement in asthma control and inflammation in children undergoing adenotonsillectomy. *Pediatr Res*. 2014;75:403-8.
55. Saito H, Asakura K, Hata M, Kataura AMK. Does adenotonsillectomy affect the course of bronchial asthma and nasal allergy? *Acta Otolaryngol Suppl*. 1996;523:212-5.
56. Ikeda FH, Horta P a DC, Bruscatto WL, Dolci JEL. Intellectual and school performance evaluation of children submitted to

-
- tonsillectomy and adenotonsillectomy before and after surgery. *Braz J Otorhinolaryngol.* 2012;78:17-23.
57. Ishman SL. Evidence-based practice. Pediatric obstructive sleep apnea. *Otolaryngol Clin North Am.* 2012;45:1055-69.
58. Cielo C, Brooks LJ. Therapies for children with obstructive sleep apnea. *Sleep Med Clin.* 2013;8:483-93.
59. den Herder C, Schmeck J, Appelboom DJK, de Vries N. Risks of general anaesthesia in people with obstructive sleep apnoea. *BMJ.* 2004;329:955-9.
60. Brown KA, Brouillette RT. The elephant in the room: lethal apnea at home after adenotonsillectomy. *Anesth Analg.* 2014;118:1157-9.