



Essay

The importance of infection control and prevention in anesthesiology[☆]



Andrés Zorrilla-Vaca ^{*₁}, Kevin Escandón-Vargas ¹

Program of Medicine and Surgery, School of Medicine, Universidad del Valle, Cali, Colombia

ARTICLE INFO

Article history:

Received 21 May 2016

Accepted 1 September 2017

Available online 23 November 2017

Keywords:

Patient safety

Anesthesiology

Communicable diseases

Anesthesia

Colombia

ABSTRACT

The key role of the field of infectious diseases in other medical specialties, including anesthesiology, is currently well known. The anesthesiologist faces a potential risk of contributing to the development of healthcare associated infections in the operating rooms; however, the infectious complications derived from anesthesia have been underestimated. It is important to acknowledge that there are some deficiencies in research, notification, and publication of reports on anesthesia-associated infectious events in developing countries, particularly in Colombia, which is the focus of our attention in this article. As far as we know, only five countries – most of them developed – have carried out studies on the practices and knowledge of the anesthesiology personnel with regards to universal recommendations for the prevention and control of anesthesia-associated infections. This document discusses the importance of infections in the area of anesthesiology and at present in Colombia. Furthermore, the need to comply with basic infection prevention and control precautions and of creating awareness of safe injection practices is recognized.

© 2017 Published by Elsevier España, S.L.U. on behalf of Sociedad Colombiana de Anestesiología y Reanimación. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

La importancia del control y prevención de enfermedades infecciosas en anestesiología

RESUMEN

Actualmente es bien conocido el protagonismo que el campo de las enfermedades infecciosas desempeña en otras especialidades médicas, incluyendo la anestesiología. El anestesiólogo tiene un riesgo potencial de contribuir al desarrollo de infecciones asociadas a la atención en salud en los quirófanos; sin embargo, las complicaciones infecciosas derivadas de la anestesia han sido subestimadas. Es importante reconocer que probablemente existen deficiencias en la investigación, notificación y publicación de reportes de

Palabras clave:

Seguridad del paciente

Anestesiología

Enfermedades transmisibles

Anestesia

Colombia

[☆] Please cite this article as: Zorrilla-Vaca A, Escandón-Vargas K. La importancia del control y prevención de enfermedades infecciosas en anestesiología. Rev Colomb Anestesiol. 2017;45:69-77.

* Corresponding author at: Facultad de Salud, Universidad del Valle, Calle 4B # 36-00, 760043 Cali, Colombia.

E-mail address: andres.zorrilla@correounivalle.edu.co (A. Zorrilla-Vaca).

¹ Both authors provided a similar contribution.

2256-2087/© 2017 Published by Elsevier España, S.L.U. on behalf of Sociedad Colombiana de Anestesiología y Reanimación. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

eventos infecciosos asociados a la anestesia en países en vías de desarrollo, particularmente en Colombia en el cual nos hemos enfocado en el presente artículo. Hasta donde se sabe, únicamente cinco países, la mayoría de ellos desarrollados, han realizado estudios sobre prácticas y conocimientos del personal de anestesiología respecto a las recomendaciones universales para la prevención y el control de infecciones asociadas a la anestesia. En el presente documento se discute la importancia de las infecciones en el campo de la anestesiología y el panorama actual de su situación en Colombia. Además, se resalta la necesidad de adherencia a precauciones básicas de prevención y control de infecciones y de concientización sobre las prácticas seguras de inyección.

© 2017 Publicado por Elsevier España, S.L.U. en nombre de Sociedad Colombiana de Anestesiología y Reanimación. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

The leading role of the field of infectious disease in other medical specialties is currently widely known. There is a growing need each year to do research on infectious diseases, in view of the increasing implications of infectious diseases for the world population, making it an exciting public health area. Infectious complications are relevant, both in developed as in developing countries, with impact at the community and hospital level.

In particular, healthcare-associated infections (HCAI) are relevant because they present in multiple healthcare environments and cause considerable morbidity, mortality, hospitalization, and costs.^{1,2} In developed countries, 3.5–12% of hospitalized patients develop at least one HCAI, while in the developing countries this percentage varies from 5.7% to 19.1%.²

Today, the prevention and control of HCAs in the surgical environment, including the anesthesiology staff, are mostly based on the understanding and practice of hand hygiene, hospital cleaning and disinfection (Ex., asepsis of the surgical site, sterilization of medical equipment), adequate implementation of invasive procedures, safe practices in the parenteral administration of medications, and use of sterile or new supplies^{3–5}; however, these recommendations and standards are frequently forgotten, ignored or infringed by the healthcare staff. Preventing nosocomial infections is essential nowadays, considering the increase in the number of multiple drug resistant organisms.^{2,5,6}

For a long time the HCAs in the operating room were associated to a large extent with general asepsis failures in the surgical environment, the type of surgical procedure, and surgeon practices, forgetting the potential risk for the anesthesiology staff during surgery.^{3,6} The relevance of HCAs in the area of anesthesiology has been evidenced for decades.^{7–9} This document discusses the importance of infections in anesthesiology and the outlook of this situation in Colombia, based on the available scientific evidence. Furthermore, emphasis shall be placed on the need to

adhere to the basic recommendations regarding infection prevention and control, and to raise awareness about safe injection practices.

Risk factors associated with pre-surgical preparation

Just as other branches in medicine, Anesthesiology is considered a dynamic medical specialty in terms of research and its professional practice focuses on the comprehensive care and safety of the patient. The prevention and control of HCAs is one of the key pillars of an ideal anesthetic practice.

Anesthesiologists often invade the body's physiological and mechanical barriers when doing respiratory tract invasive procedures (tracheal intubation) and the cardiovascular system (venous or arterial accesses), or when performing local or neuraxial blocks. These procedures are potential sources for the transmission of microorganisms to patients and may lead to infection in the presence of gaps in the universal precautions of infection control or non-compliance with the recommended healthcare practices by the staff.^{3,10–13}

Historically, a large number of common medical practices have been reported as risky, though they were originally intended to do good. These previously established practices have been reassessed in the light of new knowledge and so they have been banned or advised against because the current evidence suggests considerable risk for patients. In the area of anesthesiology it was for instance acceptable to reuse syringes for administering drugs to multiple patients, as long as the needle was changed, or to prepare intravenous injections using the same bag and administration channels for patients receiving the same therapy in one day.^{14–18} The risk of pathogen transmission among patients due to these practices is high, but we know it is totally preventable when these practices are abandoned.¹⁹ Unfortunately, it has been shown that anesthesiologists often are unaware of the importance of being compliant with all hygiene precautions and

of the fact that they are actually mechanical vectors of pathogens everyday, since they are frequently in contact with infected patients.^{8,20}

Although the cause–effect relationships in the practice of anesthesia and post-surgical infections are difficult to establish,¹¹ multiple publications since 1972 have associated anesthetic practices and equipment or contaminated anesthetic products with bacterial, viral, and fungal infections.^{10,21–33} Unfortunately, in some of these cases there have been hospital outbreaks, resulting in considerable morbidity or mortality.^{10,22,23,26–28,30,32,33} However, it is thought that outbreaks are just a proportion of the true burden of disease derived from unsafe drug injection practices and hospital gaps in infection control.³⁴ Specifically, infections from Hepatitis B Virus (HBV), Hepatitis C Virus (HCV) and Human Immunodeficiency Virus (HIV) may go undetected since these are initially asymptomatic, mildly symptomatic or unspecific.⁴

Risky behaviors in the operating room

Some of the mechanisms of transmission of frequently indiscriminate microorganisms in cases of anesthesia-associated infection include: reuse of single-use vials,^{23,26} reuse of vials or multidose vials in multiple patients,^{24,25,27,30,32,33} reuse of syringes or needles in multiple patients.^{22,23}

Propofol in particular is the anesthetic drug more often involved in infectious events due to its lipid emulsion that promotes bacterial growth and due to its frequent wrongful manipulation (for example, reusing syringes or vials in multiple patients, poor external disinfection of the vial,³⁵ and continuous propofol infusion) that facilitates its contamination.³⁶ As we documented in a recent systematic review of the risk of infections with the use of propofol,³⁶ all propofol-associated HCAIs have been reported in developed countries. This difference in scientific evidence does not seem to be different with other anesthetic products and practices studied with regards to microbial contamination or infection. Moreover, a recent systematic review on septic meningitis associated with regional anesthesia, showed the absence of clinical evidence in developing countries (Zorrilla-Vaca A., et al. Clinical occurrence of septic meningitis after spinal and epidural anesthesia, 1900–2015: a systematic review and peer reviewed). Although both anesthetic techniques show risks of infectious complications, according to a recent meta-analysis general anesthesia seems to be more closely associated.³⁷ In general, our literature search suggests that developing countries lack scientific evidence of anesthesia-associated complications or specific anesthetic practices.

One of the most researched topics is the compliance of the anesthesiologist with the recommended hygiene, prevention, and control measures, including safe injection practices involving the set of measures implemented in order to administer injections in a safe manner for the patient, the healthcare staff and third parties. **Table 1** illustrates the main findings of world cross-sectional trials aimed at identifying the level of

compliance with the practice and the knowledge of hygiene and biosafety of the anesthesiology staff.^{3,6,11,12,19,38–46} Several of these practices have been involved in the perioperative transmission of microorganisms and anesthesia-associated risk of post-surgical infection. It is concerning to find high percentage of reuse of single-use vials (50%–59.6%), reuse of multidose in multiple patients (41.3%), reuse of syringes in the same patient (46.4–82%) and reuse of syringes in multiple patients (39.8–59.2%). While there is evidence that the reuse of syringes in different patients has decreased in the most recent studies, other unsafe injection practices have remained constant or increased. Hand hygiene and the use of gloves and facemask, show a higher percentage of compliance in the most recent trials.

Woodbury et al. determined that among the 89 professionals involved in anesthesia, the primary factors preventing them from using brand new materials under certain circumstances in their clinical practice was costs (71%), convenience/efficiency (36%), environmental impact (16%), time (12%) or negligence (4%).⁶ Gounder et al. also determined among 522 anesthesiologists that the major barriers to use vials of new medicines for multiple patients were the shortage of drugs (44%), the intention to reduce wastage (44%), in addition to high costs (27%).⁴⁵ None of these factors, including cost savings, justifies reuse practices against the current recommendations; for instance, the cost of a iatrogenic infection case is certainly higher than the cost of using new materials, not to mention the ethical implications involving human distress and harmful consequences.^{6,42}

Few studies have been completed on epidemiological surveillance of anesthesia-associated infections. A Japanese trial reported that 8.3% of 6,437 patients undergoing surgery in a two-year period were infected with a virus (HBV, HCV or HIV) prior to surgery.²⁰ Another multicenter trial in France determined a 3.4 per every 1000 patients of anesthesia-associated hospital acquired infections.⁴⁷

Overall, the findings herein presented emphasize the need for further education and the importance of adhering to the universal recommendations and basic infection control guidelines. However, some professionals, despite their knowledge of the information fail to change their practice habits if they are not fully convinced about the impact and the reasons behind the recommendations. Consequently, stressing the negative outcomes associated with unsafe injection practices may have a positive effect.⁶ A poor hygiene practice by one single anesthesiologist, may result in catastrophic infectious events for multiple patients in a healthcare institution, not only placing patients at risk, but also jeopardizing the professional's job performance.⁴² Anesthesia providers have the responsibility and the opportunity to give patients good healthcare, so they must ensure the safe administration of medications and suitable performance of anesthetic procedures.⁶ The difficulties in developing countries – still unknown – should be considerably more severe as compared to the developed countries. It is then absolutely necessary to undertake national trials to validate the injection and manipulation practices of anesthetic agents by the anesthesiology staff. **Fig. 1** illustrates the hierarchy

Table 1 – Findings of studies done in anesthesiology staff on hygiene practices and biosafety knowledge for the prevention of transmission of microorganisms.

Practice	Implications	Data	Country	Year	Ref.
Hand hygiene in between procedures	Hand hygiene prevents the mechanical transmission of microorganisms in the hospital environment.	73–82.3%	USA	2015	46
		95.1%	Brazil	2011	3
		52%	France	2006	43
		93.7%	New Zealand	2006	42
		83.9%	United Kingdom	1999	12
		58–97.5% ^a	USA	1995	11
Use of gloves	Gloves protect the healthcare staff exposed to fluids and other potentially contaminated media	96.3%	Brazil	2011	3
		23%	France	2006	43
		84.2%	New Zealand	2006	42
		29%	United Kingdom	1999	41
		54.5%	United Kingdom	1999	12
		86.3%	USA	1995	11
		36.8–66.6% ^b	USA	1995 ^c	40
		23.2–55.6% ^b	USA	1995 ^d	40
Use of facemask	The facemask is a mechanism to prevent the transmission of microorganisms via droplets and aerosols from the environment to the respiratory tract of the healthcare staff	16%	United Kingdom	1992	39
		8–89% ^e	United Kingdom	1990	38
		95.2%	Brazil	2011	3
		59.5%	New Zealand	2006	42
		7.5%	United Kingdom	1999	41
Eye protection	Goggles provide protection to the healthcare staff from fluids and other potentially contaminated media	68.3%	United Kingdom	1999	12
		94.8%	USA	1995	11
		26.2%	Brazil	2011	3
		37%	New Zealand	2006	42
Disinfection of outer surfaces of vials before use	Outer surfaces (for example protective rubber) may be a source of microorganisms for the syringe needle	23.9%	USA	1995	11
		7% ^f	USA	1995 ^c	40
		29.2%	USA	2014	6
		19.8–30.2%	Brazil	2011	3
Reuse of single-use vials (access more than once)	Single use vials are potential reservoirs for microorganisms in case of several punctures (whether for one or several patients) since usually these do not contain antimicrobial additives	45.6%	New Zealand	2006	42
		51.8%	United Kingdom	1999	12
		65.6%	USA	1995	11
		59.6%	USA	2014	6
	31% ^g	31% ^g	USA	2013	45
		49%	USA	2013	19 h
		50% ⁱ	USA	2012	44

Table 1 (Continued)

Practice	Implications	Data	Country	Year	Ref.
Reuse of multidose vials in multiple patients	Although multidose vials usually contain antimicrobial additives, these may be a source of contamination and transmission of microorganisms when reused among several patients	5% 41.3%	USA New Zealand	2013 2006	45 42
Reuse of syringes in one same patient	Extended exposure of used syringes to the environment increases the risk of contamination	82% 46.4%	USA Brazil	2013 2011	19 h 3
Reuse of syringes in multiple patients	Syringes may be a source of contamination and transmission of microorganisms if reused among different patients, despite the absence of visible blood or other body fluids inside	4% 4% 1.2% ⁱ 2% 0% 6.9% 20.2% 39.8% 59.2%	USA USA Brazil France New Zealand United Kingdom USA USA USA	2013 2013 2011 2006 2006 1999 1995 1995 ^c 1995 ^d	45 19 h 3 43 42 12 11 40 40
Needle capping	Capping needles is a practice that leads to an unnecessary risk of biological accidents and encourages reuse, even if with the same product or in the same patient. Needles are intended for single use	26% 85.6% 87.1% 35%	United Kingdom USA USA United Kingdom	1999 1995 ^e 1995 ^d 1992	41 40 40 39

^a Percentage of hand hygiene depending on whether the contact was with low or high-risk patients.

^b Percentage use of gloves depending on type of practice (venous catheterization, arterial catheterization).

^c Survey performed in 1991.

^d Survey performed in 1990.

^e Percentage use of gloves depending on the type of practice (tracheal intubation, peripheral cannulation, central cannulation).

^f Percentage use of eye protection among the group of anesthesiologists that do not use prescription glasses.

^g Percentage reuse of propofol vials in multiple patients.

^h The survey was administered to students in the anesthesia nursing program, with at least 3 months of clinical training.

ⁱ Percentage with regards to propofol administration.

Source: Authors.




Fig. 1 – Pyramid illustrating the hierarchy of risk factors for anesthesia-associated infectious complications (from top to bottom: surgical asepsis, biosecurity (gloves and facemask), reuse of vials in multiple patients, reuse of syringes in multiple patients).

Source: Authors.



Fig. 2 – Advertising campaign image “One and Only Campaign”.

Source: <http://www.oneandonlycampaign.org/>.

of risk factors for anesthesia-associated infectious complications.

Surveillance and control in Colombia

At the end of our literature search, no studies were found on infections in anesthesia in Colombia, which suggests big gaps in the knowledge and awareness about the issue. In Colombia, probably the only paper on the topic was a study we recently published, showing that 6.1% of propofol vials used in the operating rooms in a third level hospital were contaminated after being used for anesthesia.⁴⁸ Although the clinical impact of this finding was not measured, surprisingly enough we found that only 26.1% of the vials used were punctured just once. A recent cross-sectional research at the national level

established that the level of reuse of vials and syringes was 37.9% and 6.2%, respectively.⁴⁹

Apparently Colombia has no specific published recommendations on anesthetic handling or guidelines for the prevention and control of anesthesia-related infections. In contrast, countries such as the United States, the United Kingdom, Canada, Australia, New Zealand, Spain, Hong Kong, and South Africa do have protocols and hygiene guidelines for the prevention and control of anesthesia-associated infections.⁵⁰⁻⁵⁶

Despite the low research interest of anesthesiology in infectious disease at the national level, it is important to highlight some of the activities sponsored by the Colombian Society of Anesthesiology and Resuscitation (S.C.A.R.E.) such as the “Revised-Comprehensive Surgical Risk Measurement” program (MIRQ-R) implemented by the promotion and prevention area, the XXXI Colombian Congress of Anesthesiology that focused on “Patient Safety” and a recent publication in the Colombian Journal of Anesthesiology entitled “Evidence-based Clinical Practice Manual: patient preparation for surgery and transfer to the operating room”.⁵⁷

Various strategies and educational advances have been implemented in other countries. In June 2008, the Coalition of Safe Injection Practices was founded with the aim of stopping unsafe injection practices in the United States.⁵⁸ This coalition, together with the US Centers for Disease Control and Prevention (CDC), launched the “One and Only Campaign” with a view to raise awareness among the healthcare professionals and the general public about safe injection practices (Fig. 2). The name of the campaign addresses the use of a needle and a syringe for a single injection in one patient. Table 2 illustrates the key recommendations for safe injection practices.⁵⁹

The interest of anesthesiology in infectious complications

There are several explanations for the lack of interest of anesthesiology in infectious diseases, particularly in developing countries such as Colombia: (1) limited knowledge about prevention and control guidelines in Health Care Associated Infections, (2) the lack of awareness regarding patient safety in surgical hospital environments, (3) minimal epidemiological follow-up and surveillance of HCAs in the operating room, (4) little economic support for implementing research activities and doing a local analysis of healthcare centers relevant data, (5) research preferences favoring mechanical applications and physiological principles of anesthesia, instead of researching adverse events associated with anesthetics contamination and basic cleansing and disinfection procedures, and (6) the perception that infectious diseases are beyond the scope or unfamiliar to anesthesiology.

We believe that the following aspects may generate increased interest: (1) the severe clinical consequences of a practice of anesthesia that fails to comply with the universal recommendations on hygiene and infections prevention and

Table 2 – Key recommendations for safe injection practices.

Recommendation	Recommended practices	Level of evidence
Use of the aseptic technique	<ul style="list-style-type: none"> Avoid contaminating the sterile injection equipment Use alcohol to disinfect the outer surface (plug, rubber, glass) of vials or ampoules (in particular propofol ampoules) before use. 	I-A _a,b
Single use of sterile syringes, needles and cannulas	<ul style="list-style-type: none"> Syringes, needles and cannulas are sterile and single use. Should not be reused in the same or in a different patient. Do not administer medications from a syringe to multiple patients, even if changing the needle or cannula. Prefilled propofol syringes are single use and the infusion shall be administered within the first 6 h of opening the syringe. 	I-A _b
Use of vials indicated for single use of parenteral medications – whenever possible	<ul style="list-style-type: none"> Do not use single use vials in multiple patients, or mix the residual contents of used vials. Propofol vials shall be used in one single patient and the infusion shall be administered within the first 12 h after opening of puncturing the vial. 	I-A _b
Use of multidose vials in one single patient – whenever possible	<ul style="list-style-type: none"> In case of using multidose vials, syringes, needles and cannulas shall be sterile. Do not store multidose vials in treatment areas; store according to the manufacturer's recommendations; dispose of any contaminated or under suspicion of being contaminated vials. 	I-A
Use of infusion lines and equipment for one single patient	<ul style="list-style-type: none"> Do not use IV solution bags or bottles as a source of anesthesia for more than one patient. Consider syringes, needles or cannulas as contaminated upon contact with a patient or after use to connect the infusion equipment. 	I-B I-B

Level I-A: strongly recommended practice for implementation and with strong evidence based on well designed experimental, clinical or epidemiological studies.

Level I-B: strongly recommended practice for implementation and evidence-based on some experimental, clinical or epidemiological studies and a strong theoretical foundation.

^a Harrison et al.³⁸

^b Hajjar and Girard.⁴⁷

Source: Authors. Based on Siegel et al.⁵⁹ and adapted from King and Ogg,⁴⁴ Hemingway et al.³⁵

control, (2) the growing generalized interest of epidemiological surveillance to learn about the outcomes associated with anesthetic procedures, (3) the actions of professional anesthesiologists avoiding legal issues due to malpractice or medical negligence, and (4) the need to fight and prevent antimicrobial resistance.

Conclusions

The anesthesiologist faces the potential risk of contributing to the development of HCAIs in the OR; however, infectious complications from anesthesia have been underestimated. In our country, infectious diseases are not as significant in anesthesiology as in other medical specialties, but this is not due to lack of merits. Despite the strong bond between the anesthesiologist and patient safety, there are few studies that clearly depict the epidemiology of anesthesia-associated infections as one of the potential healthcare complications. It is important to acknowledge that there are some flaws in

research, notification and reporting of anesthesia-associated infectious events in Colombia. The knowledge and practices of the anesthesiology staff in our country with regards to the universal recommendations for the prevention and control of anesthesia-associated infections are also unknown. Research in the area of anesthesia-associated infectious diseases would be a novelty for the healthcare staff.

Funding

Andres Zorrilla-Vaca received a research grant from S.C.A.R.E.

Conflicts of interest

The authors have no conflicts of interest to declare.

REFERENCES

1. Villalobos A, Barrero L, Rivera S, Ovalle M, Valera D. Vigilancia de infecciones asociadas a la atención en salud, resistencia bacteriana y consumo de antibióticos en hospitales de alta complejidad, Colombia, 2011. *Biomédica Rev del Inst Nac Salud.* 2014;34:67-80.
2. World Health Organization. Report on the Burden of Endemic Health Care-Associated Infection Worldwide; 2011. Available from: http://apps.who.int/iris/bitstream/10665/80135/1/9789241501507_eng.pdf WHO Libr Cat Data. 2011; 40 [cited in: October 20/2015].
3. Kishi D, Videira R. Description of nosocomial infection prevention practices by anesthesiologists in a university hospital. *Rev Bras Anestesiol.* 2011;61:177-81.
4. Perz JF, Thompson ND, Schaefer MK, Patel PR. US outbreak investigations highlight the need for safe injection practices and basic infection control. *Clin Liver Dis.* 2010;14: 137-51.
5. Munoz-Price LS, Birnbach DJ. Hand hygiene and anesthesiology. *Int Anesthesiol Clin.* 2013;51:79-92.
6. Woodbury A, Knight K, Fry L, Margolias G, Lynde GC. A survey of anesthesiologist and anesthetist attitudes toward single-use vials in an academic medical center. *J Clin Anesth.* 2014;26:125-30.
7. Du Moulin GC, Hedley-Whyte J. Hospital-associated viral infection and the anesthesiologist. *Anesthesiology.* 1983;59:51-65.
8. Browne R, Chernesky M. Infectious diseases and the anaesthetist. *Can J Anaesth.* 1988;35:655-65.
9. Zorrilla-Vaca A, Escandón-Vargas K. Bacteriostatic effect of mixtures of 1% propofol with 4% lidocaine versus 4% lidocaine alone: regards on microbiologic studies in the field of anesthesiology. *Infect Control Hosp Epidemiol.* 2014;44:747-9.
10. Ross RS, Viazov S, Gross T, Hofmann F, Seipp H-M, Roggendorf M. Transmission of hepatitis C virus from a patient to an anesthesiology assistant to five patients. *N Engl J Med.* 2000;343:1851-4.
11. Tait A, Tuttle D. Preventing perioperative transmission of infection: a survey of anesthesiology practice. *Anesth Analg.* 1995;80:764-9.
12. El Mikatti N, Dillon P, Healy TE. Hygienic practices of consultant anaesthetists: a survey in the north-west region of the UK. *Anaesthesia.* 1999;54:13-8.
13. Loftus RW, Koff MD, Birnbach DJ. The dynamics and implications of bacterial transmission events arising from the anesthesia work area. *Anesth Analg.* 2015;120:853-60.
14. Parlow J. Blood contamination of drug syringes used in anaesthesia. *Can J Anaesth.* 1989;36:S61-2.
15. Trépanier CA, Lessard MR, Brochu JG, Denault PH. Risk of cross-infection related to the multiple use of disposable syringes. *Can J Anaesth.* 1990;37:156-9.
16. Blogg CE, Ramsay MA, Jarvis JD. Infection hazard from syringes. *Br J Anaesth.* 1974;46:260-2.
17. Koepke JW, Reller LB, Masters HA, Selner JC. Viral contamination of intradermal skin test syringes. *Ann Allergy.* 1985;55:776-8.
18. Lutz CT, Bell CE, Wedner HJ, Krogstad DJ. Allergy testing of multiple patients should no longer be performed with a common syringe. *N Engl J Med.* 1984;310:1335-7.
19. Ford K. Survey of syringe and needle safety among student registered nurse anesthetists: are we making any progress? *AANA J.* 2013;81:37-42.
20. Asai T, Matsumoto S, Shingu K, Harmer M. Incidence of blood-borne infectious micro-organisms: would you still not wear gloves? *Anaesthesia.* 2000;55:591-2.
21. Olds JW, Kisch AL, Eberle BJ, Wilson JN. *Pseudomonas aeruginosa* respiratory tract infection acquired from a contaminated anesthesia machine. *Am Rev Respir Dis.* 1972;105:628-32.
22. Postsurgical infections associated with an extrinsically contaminated intravenous anesthetic agent—California, Illinois, Maine, and Michigan, 1990. *MMWR Morb Mortal Wkly Rep.* 1990;39:426-7.
23. Bennett SN, McNeil MM, Bland LA, Arduino MJ, Villarino ME, Perrotta DM, et al. Postoperative infections traced to contamination of an intravenous anesthetic, propofol. *N Engl J Med.* 1995;333:147-54.
24. Kidd-Ljunggren K, Broman E, Ekwall H, Gustavsson O. Nosocomial transmission of hepatitis B virus infection through multiple-dose vials. *J Hosp Infect.* 1999;43:57-62.
25. Tallis G, Ryan G, Lambert S, Bowden D, McCaw R, Birch C, et al. Evidence of patient-to-patient transmission of hepatitis C virus through contaminated intravenous anaesthetic ampoules. *J Viral Hepat.* 2003;10:234-9.
26. Centers for Disease Control and Prevention (CDC). Invasive *Staphylococcus aureus* infections associated with pain injections and reuse of single-dose vials – Arizona and Delaware, 2012. *Morb Mortal Wkly Rep.* 2012;61:501-4.
27. Alter MJ, Ahtone J, Maynard JE. Hepatitis B virus transmission associated with a multiple-dose vial in a hemodialysis unit. *Ann Intern Med.* 1983;99:330-3.
28. Comstock RD, Mallonee S, Fox JL, Moolenaar RL, Vogt TM, Perz JF, et al. A large nosocomial outbreak of hepatitis C and hepatitis B among patients receiving pain remediation treatments. *Infect Control Hosp Epidemiol.* 2004;25: 576-83.
29. Chant K, Kociuba K, Munro R, Crone S, Kerridge R, Quin J, et al. Investigation of possible patient-to-patient transmission of hepatitis C in a hospital. *NSW Public Heal Bull.* 1994;5:47-51.
30. Fischer G, Schaefer M, Labus B, Sands L, Rowley P, Azzam I, et al. Hepatitis C virus infections from unsafe injection practices at an endoscopy clinic in Las Vegas, Nevada, 2007-2008. *Clin Infect Dis.* 2010;51:267-73.
31. Yu H, Tang G, Liaw W, Yien H, Lee T. *Pseudomonas cepacia* induced septic shock after propofol – a case report. *Acta Anaesthesiol Sin.* 2000;38:53-6.
32. Massari M, Petrosillo N, Ippolito G, Solforosi L, Bonazzi L, Clementi M. Transmission of hepatitis C virus in a gynecological surgery setting. *J Clin Microbiol.* 2001;39:2860-3.
33. Centers for Disease Control and Prevention. Transmission of hepatitis B and C viruses in outpatient settings – New York, Oklahoma, and Nebraska, 2000–2002. *MMWR Morb Mortal Wkly Rep.* 2003;52:901-6.
34. Pugliese G, Gosnell C, Bartley JM, Robinson S. Injection practices among clinicians in United States health care settings. *Am J Infect Control.* 2010;38:789-98.
35. Hemingway CJ, Malhotra S, Almeida M, Azadian B, Yentis SM. The effect of alcohol swabs and filter straws on reducing contamination of glass ampoules used for neuroaxial injections. *Anaesthesia.* 2007;62:286-8.
36. Zorrilla-Vaca A, Arevalo J, Escandón-Vargas K, Soltanifar D, Mirski M. Infectious diseases risk and propofol anesthesia, 1989–2015. *Emerg Infect Dis.* 2015;22:981-92.
37. Zorrilla-Vaca A, Grant M, Mathur V, Li J, Wu C. The impact of neuraxial versus general anesthesia on the incidence of postoperative surgical site infections following knee or hip arthroplasty: a meta-analysis. *Reg Anesth Pain Med.* 2016;41:555-63.
38. Harrison CA, Rogers DW, Rosen M. Blood contamination of anaesthetic and related staff [see comments]. *Anaesthesia.* 1990;45:831-3.
39. O'Donnell NG, Asbury AJ. The occupational hazard of human immunodeficiency virus and hepatitis B virus infection. I.

- Perceived risks and preventive measures adopted by anaesthetists: a postal survey. *Anaesthesia*. 1992;47:923-8.
40. Rosenberg AD, Bernstein DB, Bernstein RL, Skovron ML, Ramanathan S, Turndorf H. Accidental needlesticks: do anesthesiologists practice proper infection control precautions? *Am J Anesthesiol*. 1995;22:125-32.
 41. McNamara JT, Stacey SG, McCluskey A. Poor anaesthetist hygienic practices – a problem across all grades of anaesthetist. *Anaesthesia*. 1999;54:718-9.
 42. Ryan AJ, Webster CS, Merry AF, Grieve DJ. A national survey of infection control practice by New Zealand anaesthetists. *Anaesth Intensive Care*. 2006;34:68-74.
 43. Carbone A, Veber B, Hajjar J, Zaro-Goni D, Maugat S, Seguier JC, et al. Evaluation of practices involving a cross infection risk in anaesthesia. *Ann Fr Anesth Reanim*. 2006;25: 1158-64.
 44. King C, Ogg M. Safe injection practices for administration of propofol. *AORN*. 2012;95:365-72.
 45. Gounder P, Beers R, Bornschlegel, Katherine Hinterland K, Balter S. Medication injection safety knowledge and practices among anesthesiologists: New York State, 2011. *J Clin Anesth*. 2013;25:521-8.
 46. Fernandez PG, Loftus RW, Dodds TM, Koff MD, Reddy S, Heard SO, et al. Hand hygiene knowledge and perceptions among anesthesia providers. *Anesth Analg*. 2015;120:837-43.
 47. Hajjar J, Girard R. Surveillance of nosocomial infections related to anesthesia. A multicenter study. *Ann Fr Anesth Reanim*. 2000;19:47-53.
 48. Zorrilla-Vaca A, Escandón-Vargas K, Brand-Giraldo V, León T, Herrera M, Payán A. Bacterial contamination of propofol vials used in operating rooms of a third-level hospital. *Am J Infect Control*. 2016;44:e1-3.
 49. Zorrilla-Vaca A, Ariza F, León T. Practicas de manipulación del propofol: resultados de un estudio colombiano de corte transversal. *Rev Colomb Anestesiol*. 2017;45:300-9.
 50. ASA Committee On Occupational Health Task Force On Infection Control. Recommendations for Infection Control for the Practice of Anesthesiology. Available from: [https://asahq.org/~media/sites/asahq/files/public/resources/asa%20committees/recommendations-for-infection-control-for-the-practice-of-anesthesiology-\(1\).pdf?la=en](https://asahq.org/~media/sites/asahq/files/public/resources/asa%20committees/recommendations-for-infection-control-for-the-practice-of-anesthesiology-(1).pdf?la=en) [cited in: October 20/2015].
 51. Association of Anaesthetists of Great Britain and Ireland. Infection control in anaesthesia. *Anaesthesia*. 2008;63:1027-36.
 52. Merchant R, Chartrand D, Dain S, Dobson G, Kurrek MM, Lagacé A, et al. Guidelines to the practice of anesthesia – revised edition 2015. *Can J Anaesth*. 2015;62: 54-67.
 53. Australian, New Zealand College of Anaesthetists (ANZCA), Australian and New Zealand College of Anaesthetists (ANZCA) guidelines on infection control in anaesthesia. Available from: <http://www.anzca.edu.au/documents/ps28-2015-guidelines-on-infection-control-in-anaes.pdf> [cited in: October 20/2015].
 54. De Miguel Guijarro A. Recomendaciones para el control de la infección en la práctica de la anestesia. *Rev Esp Anestesiol Reanim*. 2013;60 Suppl. 1:86-93.
 55. The Hong Kong College of Anaesthesiologists. Guidelines on infection control in anaesthesia; 2015. Available from: https://www.hkca.edu.hk/ANS/standard_publications/guidep15.pdf [cited in: October 20/2015].
 56. SASA guidelines for infection control in anaesthesia in South Africa 2014. *South Afr Anaesth Analg*. 2014;20:S1-39.
 57. Rincón Valenzuela DA, Escobar B. Manual de práctica clínica basado en la evidencia: preparación del paciente para el acto quirúrgico y traslado al quirófano. *Rev Colomb Anestesiol*. 2015;43:32-50.
 58. One needle, one syringe, only ONE time. Healthcare coalition launches new training video. *AANA NewsBull*. 2010;64:17.
 59. Siegel JD, Rhinehart E, Jackson M, Chiarello L. 2007 Guideline for isolation precautions: preventing transmission of infectious agents in health care settings. *Am J Infect Control*. 2007;35:S65-164.