



Revista Colombiana de Anestesiología

Colombian Journal of Anesthesiology

www.revcolanest.com.co



Case report

Anaesthetic management of a patient diagnosed with amyotrophic lateral sclerosis taken to caesarean section: Case report[☆]



Renato Moreno-Gonzales^{a,b,*}, Gunther Vásquez-Rojas^a, Meiling Rojas Fun^a

^a Instituto Nacional Materno Perinatal, Lima, Peru

^b Instructor, Latinoamericano Curso SAFE: Anestesia Obstétrica, Peru

ARTICLE INFO

Article history:

Received 17 April 2017

Accepted 4 August 2017

Available online 22 November 2017

Keywords:

Amyotrophic lateral sclerosis

Motor neuron disease

Cesarean section

Anesthesia

Case report

ABSTRACT

Introduction: In the obstetric population with amyotrophic lateral sclerosis, caesarean delivery may be indicated in order to avoid greater maternal respiratory distress induced by the physiologic changes at the end of gestation and the progressive muscle weakness caused by the disease. Optimal anaesthetic management is required in order to guarantee the ideal conditions for surgery and to preserve patient respiratory function.

Objective: To describe the anaesthetic management of a patient diagnosed with amyotrophic lateral sclerosis scheduled for caesarean section.

Methods and results: We describe the case of a 38-week pregnant woman diagnosed with amyotrophic lateral sclerosis who underwent caesarean section under sequential combined spinal-epidural anaesthesia. In addition, a review of the literature was conducted.

Conclusions: Combined spinal-epidural anaesthesia was a good option as anaesthetic technique for caesarean section in this patient with amyotrophic lateral sclerosis.

© 2017 Published by Elsevier España, S.L.U. on behalf of Sociedad Colombiana de Anestesiología y Reanimación. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Anestesia para cesarea en paciente con esclerosis lateral amiotrófica: reporte de caso

RESUMEN

Introducción: En las gestantes con diagnóstico de esclerosis lateral amiotrófica puede estar indicado parto por cesárea para evitar una mayor dificultad respiratoria materna debido a los cambios al final de la gestación, y a la debilidad muscular progresiva propia de la enfermedad, motivo por el cual es necesario un manejo anestésico que garantice las condiciones idóneas para la cirugía y preserve la función respiratoria.

Palabras clave:

Esclerosis lateral amiotrófica

Enfermedad de la neurona motora

Cesárea

Anestesia

Informe de Caso

[☆] Please cite this article as: Moreno-Gonzales R, Vásquez-Rojas G, Rojas Fun M. Anestesia para cesarea en paciente con esclerosis lateral amiotrófica: reporte de caso. Rev Colomb Anesthesiol. 2017;45:86-89.

* Corresponding author at: Instituto Nacional Materno Perinatal, Avenida los Ingenieros 175, Departamento 402, Lima, Peru.

E-mail address: drrenatomoreno@yahoo.es (R. Moreno-Gonzales).

2256-2087/© 2017 Published by Elsevier España, S.L.U. on behalf of Sociedad Colombiana de Anestesiología y Reanimación. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Objetivo: Describir el manejo anestésico de una paciente con diagnóstico de Esclerosis lateral amiotrófica programada para cesárea

Métodos y resultados: Mostramos el caso de una gestante de 38 semanas con diagnóstico de Esclerosis lateral amiotrófica a quien se le realizó una cesárea, y cuya técnica anestésica empleada fue CSE (combinada espinal-epidural) secuencial. Además, se realizó una revisión de la literatura al respecto.

Conclusiones: La anestesia regional CSE (combinada espinal-epidural) secuencial fue una buena opción como técnica anestésica en esta paciente con esclerosis lateral amiotrófica para cesárea.

© 2017 Publicado por Elsevier España, S.L.U. en nombre de Sociedad Colombiana de Anestesiología y Reanimación. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Amyotrophic lateral sclerosis (ALS) is a progressive neurological disorder that mainly affects male patients between the fifth and sixth decades of life, with an incidence of 1:50 000.¹ It is characterised by degeneration of higher and lower motor neurons because of degenerative changes in lateral corticospinal tracts of the spinal cord and, consequently, muscle denervation.²

ALS, or Lou-Gehrig disease, is extremely rare in pregnant women. Only some 12 cases have been reported of pregnant women with ALS since 1993. It is important to highlight that respiratory demands during pregnancy may result in acute respiratory failure in patients with ALS. For this reason, it is recommended to establish baseline respiratory function early on in pregnancy and ensure regular follow-up. In some cases, vaginal delivery may be expected because the perineum is not affected and uterine dynamics is not altered.³ However, physiological changes that become more intense towards the end of pregnancy, such as reduced residual functional capacity resulting from lower residual volume due to diaphragm displacement by the pregnant uterus, may worsen the mother's respiratory status secondary to progressive muscle weakness inherent to the disease. This could lead to the indication of early caesarean section.

There are few recommendations in the literature regarding anaesthetic management of women in childbirth with ALS. Epidural analgesia and anaesthesia have been used in these patients with no evidence of worsening of their neurological condition following the procedure. Progressive loss of innervation leads to muscle atrophy and hypersensitivity of acetylcholine nicotinic receptors. These patients are more sensitive to the use of non-depolarising neuromuscular relaxants⁴ and, furthermore, the use of depolarising muscle relaxants is associated with neuromyotonia, rhabdomyolysis and severe hyperkalemia.⁵ Because of all these considerations, a pregnant woman with ALS is a true challenge when it comes to peri-operative management and critical care.

Case description

A 38-year-old primigravida 38 weeks pregnant diagnosed with ALS some 10 years before was scheduled for category 4

caesarean section by obstetric indication. Upon neurological assessment, the patient presented with progressive muscle weakness, severe dysarthria, dysphagia, preserved superior function, fasciculations in the anterior region of the chest and the tongue, and is emotionally labile.

In terms of muscle function, the patient presented with quadriparesis (muscle strength in upper limbs 2/5 and 2/5 in the lower limbs), hypotonia, generalised laxity, hyper-reflexia and positive bilateral Babinsky sign. In terms of pulmonary function, the chest X-ray showed evidence of moderate fibroemphysema and there was severe restriction on spirometry, leading to a diagnosis of mild respiratory failure. There was absence of cardiovascular symptoms, grade 2 cardiological risk (Goldman score), no functional capacity assessment was made, and there was no evidence of murmurs or cyanosis. Anaesthetic assessment: ASA III, body weight 50 kg, height 155 cm, BMI 20.83, mouth opening 3 cm, thyromental distance greater than 6 cm, modified Mallampati grade 3, preserved cervical mobility and mandatory decubitus. Vital signs: heart rate 86 bpm, arterial blood pressure 134/92 mmHg, respiratory rate 20 rpm, oxygen saturation 95% in room air, and no fever.

The patient and husband were explained the anaesthetic technique, and after discussing the risks and benefits, a sequential combined spinal-epidural (CSE) was performed.

Before anaesthesia administration, the patient received ranitidine 50 mg IV, metoclopramide 10 mg IV and dexamethasone 4 mg IV. Basic ASA monitoring was instituted and advanced airway equipment and aspiration tube were prepared. With the patient in lateral decubitus and following asepsis and antisepsis, the epidural space was identified at the level of L3-L4, 4 cm from the skin; using a N° 18 Tuohy needle and the technique of intermittent loss of resistance, the epidural space was reached with no complications. A N° 27 Whitacre spinal needle was then inserted through the Tuohy needle until CSF flow was ascertained, and hyperbaric bupivacaine 6 mg plus fentanyl 10 µg were injected. Finally, a N° 20 multi-orifice epidural catheter was put in place and fixed 9 cm from the skin.

The catheter aspiration test was negative and the decision was made not to use a test dose; sensory anaesthetic level was obtained up to T7. With the patient in supine decubitus and 15° uterine lateralisation, 0.25% isobaric bupivacaine in 10 ml was administered through the epidural catheter 5 min after the intrathecal injection, achieving sensory block of T5 after 5 min.

The goal was to achieve adequate anaesthetic level while minimising the haemodynamic changes associated with a single-dose spinal technique. Oxygen was delivered through a reservoir mask in order to reach a level of oxygen saturation greater than 96%, and surgery was initiated. There were no heart rate or blood pressure variations greater than 10% from baseline, and there was no need for vasopressor use.

Childbirth occurred 15 min after the establishment of the neuroaxial block, with the outcome of a healthy male neonate who weighed 3520 g, with a size of 49 cm, and an APGAR score of 8 and 9 at 1 and 5 min, respectively. Oxytocin 3 IU IV was administered and the same dose was repeated 3 min later, and an infusion of 3 IU over 1 h was prepared. The surgical procedure continued uneventfully and lasted 21 min, with intermittent aspiration of saliva from the oral cavity. Total blood loss was 400 ml and the patient was administered 1000 ml of saline solution with the result of clear urine diuresis of 100 ml. The patient was given multimodal analgesia with metamizole 2 g IV and epidural morphine 2 mg.

The patient was transferred to the post-anaesthesia care unit (PACU) where she remained for 3 h with no complications. She was prescribed thromboprophylaxis with enoxaparin 40 mg SQ every 24 h, starting 12 h after discharge from the operating room.

Discussion

Pregnancy in patients with ALS is a true challenge for the medical team involved. There are certainly several considerations regarding anaesthetic management, of which respiratory function optimisation is perhaps the most important.

Pre-operative preparation is crucial and must focus on assessment of family and personal history of problems with anaesthesia, assessment of the airway and concurrent medical issues, prophylaxis to prevent aspiration as well as adequate hydration, and basic peri-operative monitoring.

Measures for preventing aspiration and post-operative nausea and vomiting (fasting of more than 8 h, H₂ blockers, metoclopramide 10 mg and ondansetron 4–8 mg) must be instituted together with antibiotic prophylaxis in order to diminish the risk of surgical site infection.

There are no clear recommendations for the selection of the anaesthetic technique. However, if general anaesthesia is chosen, it is important to consider potential complications with the airway, and the use of mechanical ventilation. Tracheal intubation is a critical moment because of gestation, some difficult airway predictors and the inability to use muscle relaxants due to the patient's clinical condition. Likewise, motor compromise, dysphagia and respiratory insufficiency could prolong mechanical ventilation which, in our situation, is provided in a maternal intensive care unit, with the associated risk of exposure to infection and prolonged length of stay among other things. However, intravenous general anaesthesia without the use of muscle relaxants may be an option for anaesthetic management.

Although there is little evidence in the literature regarding regional anaesthesia, it has been described that its use is not associated with neurological decline. In this patient, we decided to use a sequential combined spinal-epidural

technique with low-dose spinal anaesthesia in order to induce little haemodynamic changes without exacerbating respiratory function, and ensuring adequate block level by means of the epidural catheter.

The regional technique also offers the possibility of administering epidural morphine for adequate post-operative analgesia and few adverse effects when compared to systemic administration.

Whatever the anaesthetic technique, it is important to consider the high risk of development of deep vein thrombosis in these patients, with the ensuing complication of pulmonary thromboembolism due to factors such as prolonged immobilisation, gestation, and the surgical procedure itself. Consequently, consideration must be given to the use of both pharmacological as well as non-pharmacological thromboprophylaxis measures.

Conclusions

Sequential combined spinal-epidural anaesthesia (CSE) was shown to be a safe anaesthetic technique for caesarean section in this patient with ALS. It provided adequate sensory and motor block for the surgery with minimum haemodynamic changes and no exacerbation of the respiratory insufficiency present in this patient. It allowed for rapid recovery and adequate post-operative pain management.

Ethical disclosures

Protection of human and animal subjects. The authors declare that no experiments were conducted in humans or animals for this research.

Confidentiality of data. The authors declare having followed the protocols of their institution regarding disclosure of patient information.

Right to privacy and informed consent. The authors obtained the informed consent from the patient and/or subjects mentioned in this article. The informed consent form is available from the corresponding author.

Funding

The authors did not receive any funding for their work on this article.

Conflict of interest

The authors declare having no conflict of interest.

REFERENCES

1. Soriani MH, Desnuelle C. [Epidemiology of amyotrophic lateral sclerosis]. *Rev Neurol (Paris)*. 2009;165:627–40.
2. Thampi SM, David D, Thomson Chandy T, Nandhakumar A. Anaesthetic management of a patient with amyotrophic lateral

- sclerosis for transurethral resection of bladder tumor. *Indian J Anesth.* 2013;57:197-9.
3. Roanne P. Espinal cord disorders. In: *Obstetric anesthesia and uncommon disorders*. 2nd ed. New York: Cambridge University Press; 2008. p. 198-9.
 4. Bader AM. Neurologic and neuromuscular disease. In: Chestnut DH, et al., editors. *Chestnut's obstetric anesthesia: principles and practice*. 5th ed. Philadelphia: Elsevier; 2014. p. 1135.
 5. Gattenlohner S, Schneider C, Thamer C, Klein R, Roggendorf W, Gohlke, et al. Expression of foetal type acetylcholine receptor is restricted to type 1 muscle fibers in human neuromuscular disorders. *Brain.* 2002;125:130-219.