



Colombian Journal of Anesthesiology

Revista Colombiana de Anestesiología

www.revcolanest.com.co

OPEN

Wolters Kluwer

Focused critical care echocardiography: training for pediatricians and pediatric intensivists in an intensive care unit

Ecocardiografía dirigida al paciente crítico: un entrenamiento para pediatras e intensivistas pediatras en una unidad de cuidado intensivo pediátrico

Carolina Puente-Reyes^{a,b}, Hyalmar Plazas-Castro^{a,b},
María Claudia Guzmán^{a,b}, Oscar Javier León^{a,b},
Ledys María Izquierdo^{a,b,d}, Carlos Rodríguez^{a,c}

^a Department of Paediatrics and Paediatric Critical-Intensive Care, Universidad el Bosque, Bogotá, Colombia

^b Paediatric Intensive Care Unit, Hospital Cardiovascular del Niño de Cundinamarca, Bogotá, Colombia

^c School of Medicine, Universidad Nacional de Colombia, Bogotá, Colombia

^d Pediatric Intensive Care Unit, Hospital Militar Central, Bogotá, Colombia.

Keywords: Echocardiography, Critical Illness, Pediatricians, Intensive Care Units, Physicians

Palabras clave: Ecocardiografía, enfermedad crítica, pediatras, unidades de cuidados intensivos, médicos

Abstract

Introduction: Focused critical care echocardiography (FCCE) has become a necessary competency for physicians working in pediatric intensive care units.

Objective: To assess a theoretical and practical training program designed to explore skills acquisition for obtaining views and some echocardiographic measurements suggested for FCCE.

Materials and methods: A 26-hour long theoretical and practical training for pediatricians and pediatric intensivists under the guidance of a pediatric cardiologist. The program

included qualitative analysis of the variables pertaining to basic echocardiographic windows, and quantitative analysis of FCCE.

Results: There were significant differences between having prior echocardiography knowledge, associated with a higher score in the 4-chamber apical window (mean: 9.0; standard deviation [SD]: 1.02; $P=0.021$), and better correlation with the pediatric cardiologist regarding left ventricular function measurements (mean: 92.02; SD: 6.3; $P=0.036$).

Conclusion: The program was useful for basic level training in FCCE with an optimal level of acquisition of the main echocardiographic windows and some echocardiographic measurements.

How to cite this article: Puente-Reyes C, Plazas-Castro H, Guzmán MC, León OJ, Izquierdo LM, Rodríguez C. Focused critical care echocardiography: training for pediatricians and pediatric intensivists in an intensive care unit. Colombian Journal of Anesthesiology. 2019;47:23–31.

Read the Spanish version of this article at: <http://links.lww.com/RCA/A838>.

Copyright © 2018 Sociedad Colombiana de Anestesiología y Reanimación (S.C.A.R.E.). Published by Wolters Kluwer. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Correspondence: Paediatric Intensive Care Unit, Hospital Cardiovascular del Niño de Cundinamarca, Carrera 1 Este No. 31-58, Soacha, Bogotá, Colombia. E-mail: ledysmai@gmail.com

Colombian Journal of Anesthesiology (2019) 47:1

<http://dx.doi.org/10.1097/CJ9.0000000000000087>

Resumen

Introducción: La ecocardiografía, enfocada al cuidado crítico, se ha convertido hoy en día en una competencia necesaria del médico que labora en las unidades de cuidado intensivo pediátrico.

Objetivo: Evaluar un programa de entrenamiento teórico y práctico, diseñado para explorar la adquisición de habilidades en la obtención de imágenes y algunas medidas ecocardiográficas sugeridas para la ecocardiografía enfocada al cuidado crítico.

Materiales y métodos: Se realizó un entrenamiento teórico-práctico, de 26 horas de duración, por médicos pediatras y pediatras intensivistas, bajo la tutoría de un cardiólogo pediatra. El programa incluyó análisis cualitativos de las variables pertenecientes a las ventanas básicas y cuantitativos de la ecocardiografía enfocada al cuidado crítico.

Resultados: Las diferencias significativas se presentaron entre tener conocimientos previos en ecocardiografía, que se asoció a un mejor puntaje en la ventana apical cuatro cámaras (media: 9,0 DE: 1,02 $P=0,021$), y mejor correlación con el cardiólogo pediatra en la medición de la funcionalidad del ventrículo izquierdo (media: 92,2 DE: 6,3 $P=0,036$).

Conclusiones: Este programa de entrenamiento fue útil para el entrenamiento en nivel básico de la ecocardiografía enfocada al cuidado crítico, con un nivel óptimo en la adquisición de las principales ventanas ecocardiográficas, y para la toma de algunas medidas ecocardiográficas.

Introduction

The purpose of focused critical care echocardiography (FCCE)^{1,2} performed by expert cardiologists, as compared with conventional echocardiography, is early identification of hemodynamic changes in order to help with diagnostic, clinical, and therapeutic decision-making.^{3,4} This crucial information has been documented over the past 15 years by means of multiple studies in adults and children.^{1,3-6} According to the recommendations from the American Society of Echocardiography (ASE), quantitative measurements in FCCE are not necessary⁷ and their usefulness for research within the intensive care setting has not been demonstrated to date.

In Colombia, there are no formal training programs in FCCE addressed to pediatricians working in pediatric intensive care units (PICU) or pediatric intensivists, just as there are no studies documenting their performance. The objective of this study was to assess a 26-hour theory and practice FCCE training program addressed to pediatricians and pediatric intensivists working in a PICU. Moreover, it sought to achieve and document skills acquisition by correlating the performance of the trainees with that of the pediatric cardiologist acting as the gold standard, not only in terms of basic level image acquisition but also the ability to obtain some quantitative measurements that can be used in the clinical and research settings of pediatric critical care.

Methodology

Design

Cross-sectional analytical study.

Population

Practical training sessions were carried out in patients under 18 years of age admitted to the ICU of *Hospital Cardiovascular del Niño de Cundinamarca* (Children's Cardiovascular Hospital of Cundinamarca [Soacha, Colombia]), except in children with cardiac malformations or who were hemodynamically unstable. Healthy children of hospital employees were also included. Five pediatricians and 6 pediatric intensivists participated under the guidance of a pediatric cardiologist.

The parents of all the children subjected to echocardiography gave their informed consent. The study was approved by the institutional ethics and research committee of *Hospital Cardiovascular del Niño de Cundinamarca* as stated in Minutes No. 024 of December 23, 2017.

Training model and curricular content

Phase 1. Opening: Reading bibliographic material on basic echocardiography, FCCE and guidelines^{4,8-10} was provided 15 days before the start of the course. In addition, trainees received a document prepared by the authors of the training program containing principles of cardiovascular ultrasound, common imaging formats, basic echocardiographic planes (parasternal, apical, subcostal, and suprasternal), and a table with basic echocardiography measurements focused on critical care with their equation and windows. Five hours of self-learning were required.

Phase 2. Theoretical-practical course conducted by a pediatric intensive care cardiologist in 2 classroom sessions, as follows:

Six hours of theory (Table 1), based on the recommendations of the echocardiography and critical care societies.^{7,11,12}

Six hours of practice under the supervision of a pediatric cardiologist expert in echocardiography, using 3 portable machines: Siemens Acuson CV70 (United States of America, © Copyright 2018 Let Medical Systems, Corp. Store), GE Vivid 7 Ultrasound System (General Electric, Healthcare), and Phillips Sonos 5500 (Philips Electronics North America Corporation). The trainees completed a total of 3 FCCEs and the session ended with the expert pediatric cardiologist performing a complete echocardiogram (Table 2).

Phase 3. Performance of at least 20 echocardiography scans by the trainees (unsupervised) in patients admitted to the PICU; this number has been established in prior studies.^{13,14} At the end of this phase, trainees had to submit a log of practice echocardiograms which they received at the start of the training program (Annex 1). Scans could be

Table 1. Theoretical content of the training program.

Basic ultrasound and echocardiography principles
General management of portable echocardiography machines
Basic cardiac anatomy
Echocardiographic images, windows, and views
General concepts of color and Doppler modalities of echocardiography machines
Subjective and objective assessment of left ventricular function
Subjective and objective assessment of right ventricular function
Intravascular volume assessment with calculation of central venous pressure and inferior vena cava distensibility/collapsibility
General assessment of the pericardium and cardiac valves, and signs of cardiac tamponade
Indirect signs of pulmonary hypertension

Source: Authors.

repeated in the same patient at different points in time. A total of 4 weeks were devoted to this phase of the program.

Phase 4. The trainees went through a final practical examination under the supervision and evaluation of an expert pediatric cardiologist. Eligibility criteria for the final examination were completion of the 3 phases previously described, comprising the training syllabus. For this test, the trainees had to perform a complete FCCE with the 4 main echocardiographic windows, including the following measurements: inferred central venous pressure (CVP) based on inferior vena cava (IVC) diameter and its variability; IVC distensibility or collapsibility index (dIVC),

taking into consideration whether the patient was on ventilation or not; tricuspid annular plane systolic excursion (TAPSE); and left ventricular ejection fraction (LVEF) on M mode, which is the most widely used in clinical practice, especially in pediatric patients¹⁵ (Table 3).

The scale developed by Gaudet et al¹⁴ was used for assessing the quality of the acquired images. The scale consists of 2 parts:

For the first part, each FCCE window is assessed against a checklist of the main structures that needed to be visualized and scored from 0 (failure to acquire the image or inadequate image for assessment) to 2 (optimum-quality image). At the end of the checklist, the quality of

Table 2. Practical training program.

M mode LV systolic and diastolic diameter: long-axis parasternal window LV shortening fraction calculation: long-axis parasternal window or short axis in papillary muscle view LVEF calculation: long-axis parasternal window or short axis in papillary muscle view, widely used in pediatrics ¹⁵ IVC distensibility/collapsibility index calculation: subcostal window, IVC view TAPSE calculation: 4-chamber apical view
2D mode Qualitative assessment of LV function: normal or mild, moderate or severe dysfunction Qualitative assessment of RV function: normal or mild, moderate or severe dysfunction Assessment of indirect signs of pulmonary hypertension, particularly in short-axis parasternal window and 4-chamber view Assessment of the presence and severity of pericardial effusion and signs of cardiac tamponade Assessment of the normal cardiac anatomy and detection of abnormalities, including congenital heart diseases in general Situs solitus
Color mode Presence and severity of mitral or tricuspid regurgitation: mild, moderate, or severe Presence and permeability of patent ductus arteriosus and systemic-pulmonary fistula

IVC=inferior vena cava, LV=left ventricle, LVEF=left ventricular ejection fraction, RV=right ventricle, TAPSE=tricuspid annular plane systolic excursion.
Source: Authors.

Table 3. Description of echocardiographic variable measurements: CVP, TAPSE, dIVC, LVEF in M mode.

Name	Definition	Indicator
LVEF	LVEF was done using the M mode or bidimensional mode. ¹⁵ The calculation is derived from the measurement of the SF. It must be measured based on end-systolic (ESD) and end-diastolic LV diameter (EDD) just under the mitral valve leaflets in the short or long axis parasternal windows. It is calculated using the formula $SF = EDD - ESD / EDD \times 100$	Normal (EF $\geq 55\%$) Slightly reduced (EF 41%–55%) Moderately reduced (EF 31%–40%) Markedly reduced (EF $\leq 30\%$)
dIVC	$dIVC = (D_{max} - D_{min}) / D_{min} \times 100$, for patients on invasive mechanical ventilation IVC collapsibility index in extubated patients $(D_{max} - D_{min}) / D_{max} \times 100$	(dIVC $> 18\%$) for volume responders and non-responders (dIVC $< 18\%$) (dIVC $> 36\%$) for volume responders and non-responders (dIVC $< 36\%$)
TAPSE	4-chamber apical view to assess the distance of RV annular systolic excursion along the longitudinal plane	Indicates RV systolic dysfunction when it is smaller than 16 mm in adults and smaller than 12 in children ¹⁵
CVP	The primary way to assess preload and the ability to respond to fluids by echocardiography is by analyzing IVC diameter and its variability, with CVP inference	IVC diameter of less than 1.5 and more than 50% collapse: CVP between 3 and 5 Diameter between 1.5 and 2.1 with more than 50% collapse: CVP between 5 and 10 Diameter between 1.5 and 2.1 with less than 50% collapse: CVP between 10 and 15 Diameter greater than 2.1 and less than 50% collapse: CVP between 15 and 20 Dilated IVC (more than 2.1) with no collapse: CVP greater than 20

CVP=central venous pressure, dIVC distensibility index inferior vena cava, D_{max} =maximum diameter, D_{min} =minimum diameter, EDD=end-diastolic LV diameter, EF=ejection fraction, ESD=end systolic LV diameter, LV=left ventricle, LVEF=left ventricular ejection fraction, RV=right ventricle, SF=shortening fraction, TAPSE=tricuspid annular plane systolic excursion.

Source: Authors.

the echocardiographic window is assessed in terms of gain, depth, and image centering.

For the second part, the diagnostic quality of the images is assessed in terms of the ability to review LV size and systolic function, right ventricular (RV) size and systolic function, intravascular volume, and presence of pericardial effusion. Each item allows a score of 0 for inadequate images for diagnostic assessment, 1 for borderline quality images, and 2 for adequate images for diagnostic assessment.¹⁴ On the Gaudet et al image acquisition assessment scale, trainee performance is calculated dividing the score obtained for each item over the maximum score possible, and expressed as a percentage.

In each of the patients in whom the trainees did their final practical examination, the expert pediatric cardiologist conducted a complete echocardiogram and measured the required qualitative variables, 15 minutes before the trainee. The quantitative measurements calculated by the trainees were compared to those of the expert pediatric cardiologist and the correlation was expressed

as a percentage. Values above 80% were considered as a good level of correlation.

Data collection

Data regarding trainee characteristics were collected in an Excel spreadsheet (Microsoft Corporation, Redmond, WA). These included age, gender, level of education, prior training in echocardiography, number of echocardiograms performed before the practical examination, degree of satisfaction with the course following their FCCE training in terms of usefulness and applicability for their professional practice. The latter item was rated as “not useful”, “somewhat useful”, or “very useful”.

Echocardiographic variables¹⁴ were entered as follows: score and performance in long-axis parasternal window (minimum 0 points, maximum 12 points); short-axis parasternal window (minimum 0 points, maximum 18 points); score and performance in apical 4-chamber window (minimum 0 points, maximum 10 points); score

and performance in long-axis subcostal window (minimum 0 points, maximum 8 points); score and performance in subcostal echocardiographic window for inferior vena cava (minimum 0 points, maximum 8 points); score and image quality diagnostic yield (minimum 0 points, maximum 12 points); score and total performance on the Gaudet et al diagnostic image acquisition assessment scale (minimum 0 points, maximum 68 points); percentage of total performance on the Gaudet image assessment scale (minimum 0%, maximum, 100%). Performance was expressed as a percentage of trainee score \times 100/maximum score.

The correlation between measurements by the trainee versus the pediatric cardiologist was recorded as follows: TAPSE; LVEF; CVP calculation inferred on the basis of caval diameter and its variation during the respiratory cycle; dIVC (minimum 0%, maximum 100%).

Statistical analysis

The Shapiro–Wilk normality test was performed, observing a symmetrical distribution of the data. Descriptive statistics were applied using proportion estimations and relative frequencies for categorical variables; and central trend measurements were used for continuous variables. The Student t test was used to compare continuous variables for independent samples. The following were used as predictive variables: having prior knowledge of echocardiography, level of education, and number of training echocardiograms performed before the final practical examination. The control variables selected were the scores obtained for each echocardiographic window (long-axis and short-axis parasternal window, apical 4-chamber window, and long-axis and IVC subcostal windows) and the correlation percentage obtained for each echocardiographic measurement in relation to the expert pediatric cardiologist (inferred CVP, dIVC, TAPSE, and LVEF). The Stata 8.0 (Stata Corporation, College Station, TX) and SPSS Statistics 23 software packages were used for all the analyses.

Results

A total of 11 trainees participated in the FCCE training, 5 (45.5%) pediatricians and 6 (54.5%) pediatric intensivists. Mean age was 39 years, with a range between 28 and 50. Of the study population, 45.5% or the total number of pediatric intensivists that went through the training, had prior knowledge of echocardiography (general knowledge and performance of pediatric echocardiograms) (Table 4).

The course lasted 26 hours. During the training process, FCCEs were performed in a total of 198 PICU inpatients, with a mean of 18 echocardiographic scans per trainee. Not all students were able to perform all 20 echocardiograms, because of the limited time period. Upon comple-

Table 4. Characteristics of the participants in the training program.

	n=11 n (%)
Mean age \times (min–max)	39 (28–50)
Male gender (n%)	6 (54.5)
Level of education (n%)	
Pediatrician	5 (45.5)
Pediatric intensivists	6 (54.5)
Prior knowledge of echocardiography (n%)	
Yes	5 (45.5)
Number of echocardiograms performed before the test \times mean (SD)	18 (3.5)
Degree of trainee satisfaction (n%)	
Very useful	11 (100)

max = maximum, min = minimum, SD = standard deviation.

Source: Authors.

tion of the program, 100% of the participants rated it as very useful for their professional practice (Table 4). Overall, the training program resulted in high performance on the Gaudet et al image assessment scale, with a mean of 94.5%, and also a high average level of correlation for the measurements performed: 77.25% in relation to the expert pediatric cardiologist (Table 5).

The echocardiographic windows for which the trainees obtained the best average score were the subcostal long-axis view and the subcostal window for the IVC, with performances of 100% and 98.3% on the scale, respectively. The window with the best average score and performance (81.8%) was the apical 4-chamber view (Table 5).

The echocardiographic measurement with the highest average correlation between the trainees and the expert pediatric cardiologist, and a good correlation (higher than 80%) was inferred CVP (100% of the trainees), followed by dIVC (81.8% of the trainees), TAPSE (72.7% of the trainees) and LVEF (54.5% of the trainees) (Table 6).

When continuous variables were compared using the Student t test, significant differences were found only between having prior knowledge of echocardiography and obtaining a higher score for the apical 4-chamber view (mean, 9.0; standard deviation [SD], 1.02 $P=0.021$). This prior knowledge was associated also with improved correlation with the pediatric cardiologist for LVEF measurement (mean, 92.2; SD, 6.3; $P=0.036$). It was also found that a greater number of training echocardiograms (before taking the practical examination at the end of the course, with a mean of 18 echocardiograms per trainee)

Table 5. Trainee scores and performances in echocardiographic image acquisitions.

Echocardiographic window	Average score on the image acquisition assessment scale mean (SD)	Performance on the image acquisition assessment scale (%)
Average score parasternal long axis × score 0–12 points	11.4 (0.735)	95
Average score parasternal long axis × score 0–18 points	17.18 (0.815)	95.4
Average score apical 4-chamber view × score 0–10 points	8.18 (1.0)	81.8
Average score subcostal long axis × score 0–8 points	8 (0.0)	100
Average score subcostal IVC axis × score 0–8 points	7.9 (0.3)	98.3
Average image quality score × score 0–12 points	11.8 (0.4)	98.3
Average score for all views × score 0–68 points	64.5 (2.2)	94.5

IVC=inferior vena cava, SD=standard deviation.

Source: Authors.

was associated with a higher score for the sum of all the views (mean, 65.3; SD, 1.9; $P=0.038$). Likewise, it was sufficient for obtaining some measurements for the assessment of intravascular volume and cardiac function, with more than an 80% correlation with the expert pediatric cardiologist (Table 5).

Discussion

In general terms, the 26-hour FCCE training program resulted in a very good average performance on the Gaudet et al¹⁴ image assessment scale, and very high correlation level for the various measurements with the expert pediatric cardiologist.

Studies published in recent years, including a few in pediatrics like the one by Gaspar et al,¹⁵ have documented that an average of 16 to 30 echocardiograms in training programs may be sufficient to achieve adequate and

optimal FCCE performance.^{7,13,15,16} However, in this level proposed by the ASE, which is equivalent to the basic level of the American College of Chest Physicians/La Société de Réanimation de Langue Française,⁷ the only goal is to train in the acquisition of echocardiographic images and qualitative assessment of cardiac function⁷; but a few studies,^{15,17,18} this included, have found that this number of echocardiograms has also been sufficient for obtaining some measurements to assess intravascular volume and cardiac function with a correlation level greater than 70% with the cardiologist and/or expert echocardiographer.

The observation is of great value for clinical and research work in intensive care. Hence, the proposal of expanding the objectives of the basic level of the American College of Chest Physicians/La Société de Réanimation de Langue Française or of ASE level I, to include the calculation of these echocardiographic measurements. This will help optimize and obtain more objective answers and diagnoses with the bedside use of FCCE.

For people with prior experience, these types of programs may provide more expertise and help maintain their level of competency. In contrast, this study documented that 9 hours of real-time practice for pediatricians and pediatric intensivists doing FCCE resulted in a very good level of basic FCCE training and acquisition of some measurements for global cardiac function assessment.

The best performance was obtained with the subcostal long-axis window (98.75%), while the window with performances under 80% (81.8%) was the apical 4-chamber view (Table 5). It may be concluded that subcostal and parasternal views are the easiest to obtain and that the learning curve for acquiring an adequate 4-chamber view requires additional training and, consequently, a greater number of practical echocardiograms and longer training than used in this study (18 scans and 26 hours, respectively). The apical view has been recognized by the ASE as one

Table 6. Correlation for each echocardiographic measurement between the trainees and the pediatric cardiologist.

Echocardiographic measurement	Yes n (%) n=11
Inferred CVP	11 (100)
IVC distensibility index	9 (81.8)
TAPSE	8 (72.7)
LVEF	6 (54.5)

Note: Positive correlation defined as a correlation level higher than 80% for each echocardiographic measurement as compared to the intensive care pediatric cardiologist. %=percentage, CVP=central venous pressure, IVC=inferior vena cava, LVEF=left ventricular ejection fraction, n=number, TAPSE=tricuspid annular plane systolic excursion.

Source: Authors.

of the most difficult echocardiographic windows to acquire, even for cardiologists in training.⁷

The echocardiographic measurements for which trainees achieved a correlation of more than 80% with the pediatric cardiologist, considered the easiest to learn, were inferred CVP, dIVC and TAPSE. The most difficult to learn, requiring more training, was LVEF. The above shows that FCCE learning curve and skill improve with a greater number of echocardiographic scans and that more experience is required to improve performance in the 4-chamber view and the level of correlation for LVEF with the expert pediatric cardiologist.

RV dysfunction and failure has been recognized and studied more and more in the critical care setting; it is found to be an important morbidity and mortality factor in critically ill patients, affected not only by cardiovascular conditions but by the most frequent conditions seen in the ICU: acute respiratory distress syndrome, sepsis, and pulmonary embolism.^{18,19}

The gold standard for measuring RV function in view of the complex geometry is nuclear magnetic resonance imaging, which is hardly of any use in the critical care setting, even less so if bedside real-time monitoring of the RV is required for target-guided therapy designed to optimize and improve RV function.²⁰

TAPSE, together with RV fractional area change appeared to have validated these characteristics in recent years.^{20,21} The ASE^{7,8} considers that TAPSE may be obtained easily and is a measurement of longitudinal RV function. Although it only measures longitudinal function, it has been shown to have good correlation with techniques that assess global systolic RV function and must be part of the systematic assessment of the right cavities.⁸

The study explored, for the first time, the possibility of training for TAPSE measurement as part of the basic level FCCE curriculum, in order to assess the RV in the critical care setting. In average, it was the second best correlated measurement with the expert cardiologist, and the third for which a larger number of trainees achieved a level of correlation above 80% (8 out of 11 trainees). These findings suggest that TAPSE may be introduced as part of FCCE training as a useful and power tool for the quantitative assessment of RV systolic function.

Strengths

The study explored, for the first time, the possibility of teaching TAPSE measurement as part of the basic level FCCE curriculum to assess RV function in the critical care setting. This study is the first in Colombia to establish the learning curve during the entire training of pediatricians and pediatric intensivists. It shows that they are capable of performing FCCE in critically ill children and emphasizes that every time there are doubts about anatomic or functional abnormalities, the case must always be discussed and reassessed by a pediatric cardiologist.

Limitations

One of the main limitations was the small number of participants which did not allow a different type of analysis. The majority of trainees work in a PICU, with cardiovascular emphasis, and although they did not have prior knowledge in echocardiography, they have been exposed to it.

Conclusion

The training program was useful to achieve the basic FCCE level, with optimum performance in the acquisition of the main windows and some echocardiographic measurements to assess volemia, response to fluids and LV and RV function, following a 26-hour course and the performance of 18 echocardiographic scans during the practical component of the program.

An important contribution was the finding that, as part of the basic FCCE level of training, a very good level of correlation may be achieved for TAPSE, thus resulting in a more objective measurement of RV function and bedside monitoring of the RV. This type of training is the initial exploration of pediatric FCCE in Colombia for point-of-care academic, clinical and research purposes.

Ethical responsibilities

Human and animal protection. The authors declare that no experiments were conducted in humans or animals for this research.

Data confidentiality. The authors declare having followed the protocols of their institution regarding patient data disclosure.

Right to privacy and informed consent. The authors declare that not patient data appear in this article.

Funding

The authors did not receive any sponsorship for this article.

Conflict of interest

The authors declare having no conflict of interest.

References

1. Oren-Grinberg A, Talmor D, Brown SM. Concise definitive review: focused critical care echocardiography in the ICU. *Crit Care Med* 2013;41:2618–2626.
2. Beaulieu Y. Specific skill set and goals of focused echocardiography for critical care clinicians. *Crit Care Med* 2007;35 (5 suppl): S144–S149.
3. Singh Y. Echocardiographic evaluation of hemodynamics in neonates and children. *Front Pediatr* 2017;15:201.
4. Amaral Gaspar H, Saady Morhy S. The role of focused echocardiography in pediatric intensive care: a critical appraisal. *Bio Med Res Int* 2015;2015:1–7.

5. Manasia AR, Nagaraj HM, Kodali RB, et al. Feasibility and potential clinical utility of goal-directed transthoracic echocardiography performed by noncardiologist intensivists using a small hand-carried device (SonoHeart) in critically ill patients. *J Cardiothorac Vasc Anesth* 2005;19:155–159.
6. Noritomi DT, Vieira MLC, Mohovic T, et al. Echocardiography for hemodynamic evaluation in the intensive care unit. *Shock* 2010;34:59–62.
7. Spencer KT, Kimura BJ, Korcarz CE, et al. Focused cardiac ultrasound: recommendations from the American Society of Echocardiography. *J Am Soc Echocardiogr* 2013;26:567–581.
8. Rudski LG, Lai WW, Afilalo J, Hua L, Handschumacher MD, Chandrasekaran K, et al. Guidelines and standards. Guidelines for the Echocardiographic Assessment of the Right Heart in Adults: A Report from the American Society of Echocardiography. *J Am Soc Echocardiogr* 2010;23:685–713.
9. Harjola V-P, Mebazaa A, Øelutkienė J, et al. Contemporary management of acute right ventricular failure: a statement from the Heart Failure Association and the Working Group on Pulmonary Circulation and Right Ventricular Function of the European Society of Cardiology. *Eur J Heart Fail* 2016;18:226–241.
10. Lai WW, Geva T, Shirali GS, et al. Guidelines and standards for performance of a pediatric echocardiogram: a report from the Task Force of the Pediatric Council of the American Society of Echocardiography. *J Am Soc Echocardiogr* 2006;19:1413–1430.
11. Mayo PH, Beaulieu Y, Doelken P, et al. American College of Chest Physicians/La Société de Réanimation de Langue Française Statement on Competence in Critical Care Ultrasonography. *Chest* 2009;135:1050–1060.
12. Price S, Via G, Sloth E, et al. Echocardiography practice, training and accreditation in the intensive care: document for the World Interactive Network Focused on Critical Ultrasound (WINFOCUS). *Cardiovasc Ultrasound* 2008;6:49.
13. Millington SJ, Hewak M, Arntfield RT, et al. Outcomes from extensive training in critical care echocardiography: identifying the optimal number of practice studies required to achieve competency. *J Crit Care* 2017;40:99–102.
14. Gaudet J, Waechter J, McLaughlin K, et al. Focused critical care echocardiography: development and evaluation of an image acquisition assessment tool. *Crit Care Med* 2016;44:e329–e335.
15. Gaspar HA, Morhy SS, Lianza AC, et al. Focused cardiac ultrasound: a training course for pediatric intensivists and emergency physicians. *BMC Med Educ* 2014;14:25.
16. See KC, Ong V, Ng J, et al. Basic critical care echocardiography by pulmonary fellows: learning trajectory and prognostic impact using a minimally resourced training model. *Crit Care Med* 2014;42:2169–2177.
17. Croft LB, Duvall WL, Goldman ME. A pilot study of the clinical impact of handcarried cardiac ultrasound in the medical clinic. *Echocardiography* 2006;23:439–446.
18. Longjohn M, Wan J, Joshi V, et al. Point-of-care echocardiography by pediatric emergency physicians. *Pediatr Emerg Care* 2011; 27:693–696.
19. Guzmán MC, Izquierdo LM, Carvajal D, et al. Clinical outcomes of pulmonary hypertension in children with pneumonia and respiratory failure. *J Respir Disord* 2017;2:1–6.
20. Krishnan S, Schmidt GA. Acute right ventricular dysfunction: real-time management with echocardiography. *Chest* 2015;147: 835–846.
21. Ventetuolo CE, Klinger JR. Management of acute right ventricular failure in the intensive care unit. *Ann Am Thorac Soc* 2014;11: 811–822.

Source: Authors.

SCANS PERFORMED BEFORE THE WRITTEN TEST																											
TRAINEE NAME:																											
DIAGNOSIS: Age(m):																											
Subcostal view				I	A	MB	Long parasternal				I	A	MB	Short parasternal			I	A	MB								
4 chamber view				I	A	MB	5 Chamber View				I	A	MB	Suprasternal axis			I	A	MB								
LVEF				%	Y	N	PASP				mmHg	Y	N	TAPSE				cm	Y	N	IVC				%	Y	N
Area fraccional						%	Y	N	Other																		
DIAGNOSIS: Age(m):																											
Subcostal view				I	A	MB	Long parasternal				I	A	MB	Short parasternal			I	A	MB								
Vista 4 camaras				I	A	MB	5 Chamber View				I	A	MB	Suprasternal axis			I	A	MB								
LVEF				%	Y	N	PASP				mmHg	Y	N	TAPSE				cm	Y	N	IVC				%	Y	N
Area fraccional						%	Y	N	Other																		
DIAGNOSIS: Age(m):																											
Subcostal view				I	A	MB	Long parasternal				I	A	MB	Short parasternal			I	A	MB								
Vista 4 camaras				I	A	MB	5 Chamber View				I	A	MB	Suprasternal axis			I	A	MB								
LVEF				%	Y	N	PASP				mmHg	Y	N	TAPSE				cm	Y	N	IVC				%	Y	N
Area fraccional						%	Y	N	Other																		
DIAGNOSIS: Age(m):																											
Subcostal view				I	A	MB	Long parasternal				I	A	MB	Short parasternal			I	A	MB								
Vista 4 camaras				I	A	MB	5 Chamber View				I	A	MB	Suprasternal axis			I	A	MB								
LVEF				%	Y	N	PASP				mmHg	Y	N	TAPSE				cm	Y	N	IVC				%	Y	N
Area fraccional						%	Y	N	Other																		
DIAGNOSIS: Age(m):																											
Subcostal view				I	A	MB	Long parasternal				I	A	MB	Short parasternal			I	A	MB								
Vista 4 camaras				I	A	MB	5 chamber view				I	A	MB	Suprasternal axis			I	A	MB								
LVEF				%	Y	N	PASP				mmHg	Y	N	TAPSE				cm	Y	N	IVC				%	Y	N
Area fraccional						%	Y	N	Other																		