

## Comparison of non-linear models to describe growth of Iranian Guilan sheep<sup>□</sup>

*Comparación de modelos no lineales para describir el crecimiento de ovejas iraní Guilan*

*Comparaçãõ de modelos não lineares para descrever o crescimento de ovinos iraniano Guilan*

Navid Ghavi Hossein-Zadeh<sup>1\*</sup>, PhD; Mohammad Golshani<sup>2</sup>, MSc.

<sup>1</sup>*Department of Animal Science, Faculty of Agricultural Sciences, University of Guilan, Rasht, Iran.*

<sup>2</sup>*Agricultural Organization of Guilan Province, Rasht, Iran.*

*(Received: September 15, 2015; accepted: April 14, 2016)*

*doi: 10.17533/udea.rccp.v29n3a05*

### Summary

**Background:** non-linear mathematical models, empirically developed by plotting body weight against age, are used to describe the growth curve in different animals. **Objective:** to describe the growth pattern in Guilan sheep using non-linear models. **Methods:** six non-linear mathematical equations (Brody, Negative exponential, Logistic, Gompertz, von Bertalanffy and Richards) were used to describe the growth curves in Guilan sheep. The Agricultural Organization of Guilan province (Rasht, Iran) provided the dataset used in this study. The dataset included 42,257 weight records of lambs from birth to 240 days of age during years 1994 to 2014. Each model was separately fitted to body weight records of all lambs, males and females, using the NLIN and MODEL procedures of SAS. The models were tested for goodness of fit using adjusted coefficient of determination, root means square error (RMSE), Durbin-Watson statistic, akaike's information criterion (AIC) and bayesian information criterion (BIC). **Results:** the Richards model provided the best fit to the growth curve in females and all lambs, with the lowest RMSE, AIC, and BIC values compared to the other models. The Brody model provided the best fit of growth in male lambs due to the lower values of AIC and BIC compared to the other models. The negative exponential model provided the worst fit of growth for males, females and all lambs. **Conclusion:** the evaluation of the growth equations used in this study indicates the potential of non-linear functions for fitting body weight records of Guilan sheep.

**Keywords:** *body weight, fat-tailed sheep, growth function, growth pattern.*

□ To cite this article: Ghavi Hossein-Zadeh N, Golshani M. Comparison of non-linear models to describe growth of Iranian Guilan sheep. Rev Colomb Cienc Pecu 2016; 29:199-209.

\* Corresponding author: Navid Ghavi Hossein-Zadeh. Department of Animal Science, Faculty of Agricultural Sciences, University of Guilan, Rasht, 41635-1314, Iran. Tel: + 98 13 33690274. Fax: + 98 13 33690281. E-mail: nhosseinzadeh@guilan.ac.ir

## Resumen

**Antecedentes:** los modelos matemáticos no lineales desarrollados empíricamente mediante la comparación del peso corporal contra la edad se han usado para describir la curva de crecimiento en diferentes animales. **Objetivo:** describir el patrón de crecimiento en ovejas Guilan usando modelos no lineales. **Métodos:** seis ecuaciones matemáticas no lineales (Brody, Exponencial negativo, Logístico, Gompertz, von Bertalanffy y Richards) se utilizaron para describir las curvas de crecimiento en ovejas Guilan. El conjunto de datos utilizados en este estudio se obtuvo de la Organización para la Agricultura de la provincia Guilan (Rasht, Irán), y comprendió 42,257 registros de peso de corderos, recogidos desde el nacimiento hasta los 240 días de edad durante los años 1994 a 2014. Cada modelo fue ajustado por separado a los registros de peso corporal de todos los corderos, machos y hembras, utilizando los procedimientos NLIN y MODEL de SAS. Los modelos fueron probados para la bondad del ajuste mediante el coeficiente de determinación ajustado, la raíz cuadrada del error cuadrático medio (RMSE), el estadístico de Durbin-Watson, el criterio de información de akaike (AIC) y el criterio de información bayesiano (BIC). **Resultados:** el modelo de Richards proporcionó el mejor ajuste de la curva de crecimiento en las hembras y todos los corderos debido a los valores más bajos de RMSE, AIC y BIC con respecto a los otros modelos. El modelo Brody proporcionó el mejor ajuste de la curva de crecimiento de los corderos machos debido a los valores más bajos de AIC y BIC con respecto a los otros modelos. El modelo Exponencial negativo proporcionó el peor ajuste de la curva de crecimiento para machos, hembras y todos los corderos. **Conclusión:** la evaluación de diferentes ecuaciones de crecimiento utilizadas en este estudio indica el potencial de las funciones no lineales para el ajuste de los registros de peso corporal en ovejas de raza Guilan.

**Palabras clave:** *función de crecimiento, ovejas de cola gorda, patrón de crecimiento, peso corporal.*

## Resumo

**Antecedentes:** os modelos matemáticos não lineares desenvolvidos empiricamente através da representação gráfica do peso corporal com a idade têm sido adequados para descrever a curva de crescimento em diferentes animais. **Objetivo:** o objetivo deste estudo foi descrever o padrão de crescimento em ovinos Guilan usando modelos não lineares. **Métodos:** seis equações matemáticas não lineares (Brody, Negativo exponencial, Logístico, Gompertz, von Bertalanffy e Richards) foram utilizadas para descrever as curvas de crescimento em ovelhas Guilan. O conjunto de dados utilizado neste estudo foi obtido a partir da Organização Agrícola da província de Guilan (Rasht, Irã) e composta 42.257 registros de peso de cordeiros que foram recolhidos desde o nascimento até 240 dias de idade, durante 1994 a 2014. Cada modelo foi ajustado separadamente para os registros de peso corporal de todos os cordeiros, machos e fêmeas, utilizando os procedimentos NLIN e modelo em SAS. Os modelos foram testados para a qualidade do ajuste por meio do coeficiente de determinação ajustado, a raiz quadrada de erro quadrático médio (RMSE), o estatístico Durbin-Watson estatística, o critério de informação de Akaike (AIC) e critério de informação Bayesiano (BIC). **Resultados:** o modelo de Richards providencia o melhor ajuste da curva de crescimento nas fêmeas e todos os cordeiros, devido aos menores valores de RMSE, AIC e BIC do que outros modelos. O modelo Brody da o melhor ajuste da curva de crescimento em cordeiros devido aos menores valores de AIC e BIC do que outros modelos. O modelo exponencial negativo oferece o pior ajuste de curva de crescimento para machos, fêmeas e todos os cordeiros. **Conclusão:** avaliação diferentes equações de crescimento utilizados neste estudo indicaram o potencial das funções não lineares para a montagem de registros de peso corporal de Guilan.

**Palavra chave:** *função crescimento, ovinos de cauda gorda, padrão de crescimento, peso corporal.*

## Introduction

Guilan sheep is a fat-tailed breed in Iran, numbering some 400,000 animals in the north of the country, and distributed in the northern and western parts of Guilan Province in the mountains between Assalem, Khalkhal, Oshkourat, and Deilaman. This breed can also be found in some areas of the Guilan-Zanjan border. Mean adult live weight in this breed is

35 Kg (77 lbs) for rams and 31 Kg (67 lbs) for ewes. The coat is yellowish-white to pure white, with brown patches on the head, face, and the bottom of the legs. This breed is valued mainly due to its ability to live in mountainous areas with rain-fed foothills and foothill steppes with 1300 mm (51 inches) annual rainfall. Young ewes are randomly exposed to the rams for the first time at approximately 1.5 years of age. Ewes are kept in the flock up to 7 years of age. Ewes are

supplemented, depending upon requirements, for a few days after lambing. Rams are kept until a male offspring is available for replacement. During the breeding season, single-sire pens are used allocating 20-25 ewes per ram. Lambs remain with their dam until weaning. Lambs are ear-tagged and weighted immediately after lambing. During the suckling period, lambs are fed dry alfalfa after three weeks of age. Lambs are weaned at approximately 90 days of age. Animals are kept on natural pasture during spring, summer and autumn. Since environmental conditions are adverse, the animals are kept indoors during the winter months. The flock is mainly kept on pasture and fed cereal pasture, with supplemental feed, including alfalfa and wheat straw, provided especially around the mating season (Eteqadi *et al.*, 2014).

Growth is defined as increase in live weight or dimension against age. Changes in live weight or dimension for a period of time are explained by growth curves (Keskin *et al.*, 2010). Analysis of animal growth performance through the life span is helpful to establish appropriate feeding strategies and the best age for slaughtering. Studies focusing on growth curves have increased in recent years due to the development of new computational methods for faster and more accurate analyses as well as the availability of new models (Souza *et al.*, 2013). Slow growth rates resulting in low market weight has been identified to be one of the factors limiting profitability in production system (Noor *et al.*, 2001; Abegaz *et al.*, 2010). Growth rate is related to rate of maturing and mature weight and these latter traits have been suggested to be related with other lifetime productivity parameters in sheep (Bedier *et al.*, 1992; Abegaz *et al.*, 2010).

Non-linear mathematical models, empirically developed by plotting body weight against age, are suitable to describe the growth curve in different animal groups (Malhado *et al.*, 2009). The use of mathematical growth models provides a good way of condensing the information into a few parameters with biological meaning, to facilitate both the interpretation and the understanding of the phenomenon (Fitzhugh, 1976; Malhado *et al.*, 2009). Growth curve parameters provide potentially useful criteria for modifying the relationship between body weight and age through selection (Kachman and

Gianola, 1984) and an optimum growth curve can be obtained by selection for desired values of growth curve parameters (Bathaei and Leroy, 1998). Growth curves provide several applications to animal production, such as (1) evaluation of the response to treatments over time; (2) analysis of the interaction between subpopulations (or treatments) and time; and (3) identification of heavier animals at younger ages within a population (Bathaei and Leroy, 1996; Freitas, 2005; Malhado *et al.*, 2009).

No previous studies have been conducted on growth curve characteristics of the Guilan sheep. Therefore, the objective of this study was to describe the growth pattern in Guilan sheep using non-linear models. For this purpose, six mathematical models (Brody, Negative exponential, Logistic, Gompertz, von Bertalanffy, and Richards) currently used to fit the growth pattern of animals were examined to evaluate their efficiency in describing the growth curve of Guilan sheep.

## Materials and methods

The data set used in this study was obtained from the Agricultural Organization of Guilan Province (Rasht, Iran) and comprised 42,257 weight records, which were collected on 18,972 lambs from birth to 240 days of age from years 1994 to 2014. The data were screened several times and defective and out of range records were deleted. After editing the initial data set, 41,894 body weight records (made up of 19,879 for males and 22,015 for female lambs) were used for statistical analysis.

Non-linear growth functions used to describe the growth curves of Guilan sheep are presented in Table 1. The Brody, Negative exponential, Logistic, Gompertz, Von Bertalanffy and Richards functions were fit to the data to model the relationship between weight and age. Each model was fitted separately to body weight records of all lambs, male and female lambs using the NLIN and MODEL procedures in SAS (SAS Institute Inc., Cary, NC, USA; 2002) and the parameters were estimated. The NLIN procedure produces least squares or weighted least squares estimates of the parameters of a non-linear model. For each non-linear model to be analyzed, the model

(using a single dependent variable) and the names and starting values of the parameters to be estimated must be specified. When non-linear functions were fitted, the Gauss-Newton method was used as the iteration method. To begin this process the NLIN procedure first examines the starting value specifications of the parameters. If a grid of values is specified, NLIN procedure evaluates the residual sum of squares at each combination of parameter values to determine the set of parameter values producing the lowest residual sum of squares. These parameter values are used for the initial step of the iteration. The MODEL procedure analyzes models in which the relationships among the variables comprise a system of one or more nonlinear equations. Primary uses of the MODEL procedure are estimation, simulation, and forecasting of nonlinear simultaneous equation models (Ghavi Hossein-Zadeh, 2014).

**Table 1.** Functional forms of equations used to describe the growth curve of Guilan sheep.

Equation	Functional form
Brody	$y = a(1 - be^{-kt})$
Negative exponential	$y = a - (ae^{-kt})$
Logistic	$y = \frac{a}{1 + be^{-ct}}$
Gompertz	$y = ae^{-be^{-kt}}$
von Bertalanffy	$y = a(1 - be^{-kt})^3$
Richards	$y = a(1 - be^{-kt})^{-m}$

y = body weight at age t (day); a = asymptotic weight, which is interpreted as mature weight; b = is an integration constant related to initial animal weight. The value of b is defined by the initial values for y and t; k = is the maturation rate, which is interpreted as weight change in relation to mature weight to indicate how fast the animal approaches adult weight; m = is the parameter that gives shape to the curve by indicating where the inflection point occurs.

The models were tested for goodness of fit (quality of prediction) using adjusted coefficient of determination ( $R^2_{adj}$ ), residual standard deviation or root means square error (RMSE), Durbin-Watson statistic (DW), Akaike's information criterion (AIC) and Bayesian information criterion (BIC).

$R^2_{adj}$  was calculated using the following formula:

$$R^2_{adj} = 1 - \left[ \frac{(n-1)}{(n-p)} \right] (1 - R^2)$$

Where:

$R^2$  = is the multiple coefficient of determination

$$(R^2 = 1 - \frac{RSS}{TSS}).$$

TSS = is the total sum of squares.

RSS = is the residual sum of squares.

n = is the number of observations (data points).

p = is the number of parameters in the equation.

The  $R^2$  value is an indicator measuring the proportion of total variation about the mean of the trait explained by the growth curve model. The coefficient of determination lies always between 0 and 1, and the fit of a model is satisfactory if  $R^2$  is close to unity.

RMSE is a sort of generalized standard deviation and was calculated as follows:

$$RMSE = \sqrt{\frac{RSS}{n-p-1}}$$

Where.

RSS = is the residual sum of squares.

n = is the number of observations (data points).

p = is the number of parameters in the equation.

RMSE is one of the most important criteria to compare the suitability of growth models. Therefore, the best model is the one with the lowest RMSE.

DW was used to detect the presence of autocorrelation in the residuals from the regression analysis. In fact, the presence of autocorrelated residuals suggests that the function may be inappropriate for the data. The Durbin-Watson statistic ranges in value from 0 to 4. A value near two indicates no autocorrelation; a value toward 0 indicates positive autocorrelation; a value toward 4 indicates negative autocorrelation (Ghavi Hossein-Zadeh, 2014). DW was calculated using the following formula:

$$DW = \frac{\sum_{t=1}^n (e_t - e_{t-1})^2}{\sum_{t=1}^n e_t^2}$$

Where:

$e_t$  = is the residual at time t.

$e_{t-1}$  = is the residual at time t-1.

AIC was calculated as using the following equation (Burnham and Anderson, 2002):

$$AIC = n \times \ln(RSS) + 2p$$

AIC is a good statistic to compare models of different complexity because it adjusts the RSS for the number of parameters in the model. A smaller numerical value of AIC indicates a better fit when comparing models.

BIC combines maximum likelihood (data fitting) and choice of model by penalizing the (log) maximum likelihood with a term related to model complexity, as follows:

$$BIC = n \ln\left(\frac{RSS}{n}\right) + p \ln(n)$$

A smaller numerical value of BIC indicates a better fit when comparing models.

## Results

Estimated parameters of non-linear growth functions for Guilan sheep are presented in Table 2. Also, goodness of fit statistics for the six growth models fitted to body weight records are presented in Table 3.  $R_{adj}^2$  values had little differences among the models for all lambs, males and females, but Brody model and Richards equation provided the greatest  $R_{adj}^2$  values for all lambs, males and females, and Negative exponential provided the lowest values of  $R_{adj}^2$  for males, females and all lambs (Table 3). Also, Negative exponential function had the lowest values of DW for males, females and all lambs, but other functions had little differences in this regard. In general, DW values were relatively low for different models and the Negative exponential model provided

the lowest value for males, females and all lambs. The Richards equation provided the lowest values of AIC and BIC for females and all lambs, but the Brody model provided the lowest values of AIC and BIC for male lambs. Richards and Brody equations provided the lowest values of RMSE for females and all lambs. Also, Richards, Brody and von Bertalanffy equations provided the lowest values of RMSE for male lambs. But, Negative exponential model provided the greatest values of RMSE, AIC and BIC for all lambs, males and female lambs. Therefore, the Brody model was selected as the best model for fitting the growth curve in males, and the Richards model provided the best fit of growth curve in females and all lambs. The Negative exponential model provided the worst fit of growth curve for males, females and all lambs. The parameter  $a$  is considered as an estimate of asymptotic weight. For the data set studied here there were low asymptotic weight estimates for Brody and Richards equations as the best fitted growth models (Table 2).

The  $k$  parameter, which represents maturation rate, is another important trait to be considered, since it indicates the growth speed to reach the asymptotic weight. In this study, females generally showed higher values for this parameter than males. The negative correlation of -0.99 to -0.98 between  $a$  and  $k$  parameters obtained, was based on the best model for males, females and all lambs. Observed body weights of animals from 1 to 240 days of age are depicted in Figure 1. As shown, there was a sigmoidal trend for body weights along with increase in age. Also, predicted body weights (Kg) as a function of age (days) obtained with different growth models for all lambs, males and females, are presented in Figure 2. The estimated growth curves were typically sigmoid.

## Discussion

Once an appropriate model of the growth curve is selected, selection emphasis can then be directed exclusively to the level of the growth curve. It is important to develop an optimal strategy to achieve a desired growth shape through changing the growth model parameters. In previous studies, a broad range of growth models have been selected, depending on how accurately they fit the data. Tariq *et al.* (2013) selected the Morgan-Mercer-Flodin model as the best

**Table 2.** Parameter estimates for the different growth models in Guilan sheep (standard errors are in parentheses).

Item	Parameter	Model					
		Brody	Negative exponential	Logistic	Gompertz	von Bertalanffy	Richards
All lambs	a	27.42 (0.19)	24.60 (0.12)	21.68 (0.045)	23.13 (0.07)	24.03 (0.09)	31.68 (1.85)
	b	0.89 (0.001)	-	5.69 (0.05)	2.01 (0.008)	0.50 (0.001)	0.96 (0.01)
	k	0.0068 (0.00009)	0.0096 (0.0001)	0.0251 (0.0001)	0.0152 (0.00009)	0.0122 (0.00009)	0.0041 (0.0008)
	m	-	-	-	-	-	-0.73 (0.06)
Male lambs	a	29.07 (0.30)	25.80 (0.19)	22.46 (0.07)	24.08 (0.11)	25.10 (0.14)	30.73 (1.84)
	b	0.90 (0.002)	-	5.81 (0.07)	2.03 (0.01)	0.50 (0.002)	0.93 (0.03)
	k	0.0064 (0.0001)	0.0091 (0.0001)	0.0248 (0.0002)	0.0149 (0.0001)	0.0119 (0.0001)	0.0052 (0.0010)
	m	-	-	-	-	-	-0.87 (0.10)
Female lambs	a	26.21 (0.23)	23.69 (0.15)	21.04 (0.06)	22.38 (0.09)	23.20 (0.11)	33.07 (3.71)
	b	0.89 (0.002)	-	5.61 (0.07)	2.00 (0.01)	0.49 (0.002)	0.98 (0.01)
	k	0.0071 (0.0001)	0.0099 (0.0001)	0.0254 (0.0002)	0.0155 (0.0001)	0.0125 (0.0001)	0.0032 (0.0010)
	m	-	-	-	-	-	-0.65 (0.07)

fitted equation for growth curve in Mengali sheep breed of Balochistan. Similar to the results of this study, Gbangboche *et al.* (2008) selected the Brody model as the best function to fit the growth curve of the African Dwarf sheep. On the other hand, Freitas (2005) reported that Logistic, Von Bertalanffy and Brody functions were more versatile to fit growth curves in sheep. In the Bergamasca sheep in Brazil (McManus *et al.*, 2003), among the fitted growth models (Brody, Richards, and Logistic), the Logistic model showed the goodness of fit. Lambe *et al.* (2006) selected the Richards and Gompertz models for their accuracy of fit among four competing models (Gompertz, Logistic, Richards and the Exponential model). The Gompertz and von Bertalanffy models showed the best fit in Morkaraman and Awassi lambs (Topal *et al.*, 2004); Gompertz model in Suffolk sheep (Lewis *et al.*, 2002). Malhado *et al.* (2009) reported that both Gompertz and Logistic

functions presented the best fit of growth curve in Dorper sheep crossed with the local Brazilian breeds Morada Nova, Rabo Largo, and Santa Inês. Bathaei and Leroy (1996), evaluating growth in Mehraban Iranian fat-tailed sheep, selected the Brody function because of simplicity of interpretation and ease of estimation. Sarmento *et al.* (2006) observed that the Gompertz function presented the best adjustment when compared to the other models in studies of growth curves of Santa Inês sheep herds in the state of Paraíba, Brazil. Akbas *et al.* (1999) studied the fitting performance of Brody, Negative exponential, Gompertz, Logistic and von Bertalanffy models to data on weight-age of Kivircik and Daglic male lambs and found that the Brody model was the best equation for describing the growth of lambs. Similar to the current results, Goliomytis *et al.* (2006) and Ghavi Hossein-Zadeh (2015) reported the excellent fit of the Richards function to the weight-age data

**Table 3.** Goodness of fit for different growth curves in Guilan sheep.

Item	Statistics	Model						
		Brody	Negative exponential	Logistic	Gompertz	von Bertalanffy	Richards	
All lambs	$R_{adj}^2$	0.9530	0.9378	0.9520	0.9527	0.9528	0.9530	
	DW	0.70	0.53	0.71	0.71	0.71	0.70	
	RMSE	3.16	3.64	3.20	3.17	3.17	3.16	
	AIC	542406	554126	543311	542664	542536	542393	
	BIC	96559	108270	97463	96817	96688	96554	
	Male lambs	$R_{adj}^2$	0.9532	0.9372	0.9522	0.9529	0.9530	0.9532
Male lambs	DW	0.71	0.53	0.72	0.71	0.71	0.71	
	RMSE	3.20	3.70	3.23	3.21	3.20	3.20	
	AIC	242981	248794	243383	243084	243029	242982	
	BIC	46254	52059	46656	46357	46302	46263	
	Female lambs	$R_{adj}^2$	0.9540	0.9395	0.9529	0.9537	0.9538	0.9540
	Female lambs	DW	0.70	0.54	0.71	0.71	0.71	0.70
RMSE		3.09	3.55	3.13	3.10	3.10	3.09	
AIC		269871	275901	270378	270025	269951	269856	
BIC		49756	55779	50264	49910	49837	49750	


$R_{adj}^2$ : Adjusted coefficient of determination; RMSE: Root means square error; DW: Durbin–Watson; AIC: Akaike information criteria; BIC: Bayesian Information Criteria.

of Karagouniko and Shall sheep, respectively. On the other hand, da Silva *et al.* (2012) observed that Richards model was problematic during the process of convergence in Santa Inês sheep.

According to the low DW values obtained in the current study from fitting the non-linear models of the growth curve, it was concluded that there was positive autocorrelation between residuals for all models. The Negative exponential model provided the most positive autocorrelation among different models. It seems that residual autocorrelation presented no problem, or only a very slight one in the current study. Positive autocorrelation is a serial correlation in which a positive error for one observation increases the chances of a positive error for another observation (Ghavi Hossein-Zadeh, 2014). The lower estimates of  $a$  parameter, obtained from the best fitted models in this study, may indicate that animals are lighter as adults and may be considered fast-growing, as these sheep require less time to reach maturity compared to other breeds (da Silva *et al.*, 2012).

The definition of an optimum adult weight is controversial as it depends on the species, breed, selection method, management system and environmental conditions (Malhado *et al.*, 2009). It was reported that average adult live weight in Guilan sheep is 35 Kg (77 lbs) for rams and 31 Kg (67 lbs) for ewes (Eteqadi *et al.*, 2014). These values were generally consistent with the estimates of the  $a$  parameter obtained from the best fitted models in this study.

Contrary to the current results, da Silva *et al.* (2012) reported high asymptotic weight estimates in Santa Inês sheep, but their animals were between 120 and 774 days of age. Lôbo *et al.* (2006) evaluated the growth curve of Santa Inês sheep from birth to 550 days of age and obtained greater  $a$  value than the current study. Also, McManus *et al.* (2003) in the study of growth curve from birth to 2,000 days of age in Bergamácia sheep obtained a greater estimate of  $a$  parameter compared with this study. Generally, consistent with the current results, Malhado *et al.* (2008)




**Figure 1.** Observed body weights for all lambs (a), males (b), and female lambs (c).

studied the growth curve of Texel×Santa Inês crossed sheep from birth up to 365 days of age and obtained  $a$  values ranging from 29.14 to 32.16 Kg. Also, Malhado *et al.* (2009) reported  $a$  values of 29.35 to 32.41 in the fit of different growth functions for the Morada Nova, Rabo Largo, and Santa Inês Brazilian sheep breeds. These Brazilian breeds were weighted from birth up to

210 days of age in the study by Malhado *et al.* (2009). Toral (2008) warned that inferences about estimated parameters may often be subject to errors because the datasets that are used to fit non-linear growth functions do not include animal growth until maturity. Similarly, Garner *et al.* (2005) stated that when the available data cover only growth before reaching maturity,





**Figure 2.** Predicted body weights (Kg) as a function of age (days) obtained with different growth models for all lambs (a), male (b), and female (c) lambs.

these data can produce erroneous  $a$  and  $k$  estimates. Animals with high  $k$  values show precocious maturity in relation to those with lower  $k$  values and similar initial weight. The greater values of  $k$  for female lambs in this study indicates higher maturity rates (i.e., they reached mature weight earlier).

Because of the narrow deviation range in the weight at birth, the variation among the  $k$  values becomes a reliable predictor of the growth rate (Malhado *et al.*, 2009). Therefore, animals with higher  $k$  values reach asymptotic weight earlier. Bathaei and Leroy (1996) estimated much higher

$k$  values for body weight than the current study, for males and females of the Mehraban Iranian fat tailed sheep, when applying the Brody growth model over a 48-month period. This difference could be attributed to the time unit (month) and time period (four years) they used compared with the present study, and the different growth function used (Goliomytis *et al.*, 2006). Goonewardene *et al.* (1981), Rogers *et al.* (1987) and Perotto *et al.* (1992) obtained different  $k$  values when applying different growth functions to the same weight-age data, indicating that the model of choice affects parameter estimates.

The most important biological correlation for a growth curve is between  $a$  and  $k$  parameters. The negative correlation between these parameters in the current study implies that the earliest animals are less likely to exhibit high adult weight; i.e., animals that have higher adult weights generally present lower growth rates than animals with a lower adult weight (da Silva *et al.*, 2012). Also, similar to the sigmoidal patterns of growth curves in this study were obtained by other studies (Goliomytis *et al.*, 2006; Malhado *et al.*, 2009; da Silva *et al.*, 2012).

The six non-linear functions investigated in the present study were adequate in describing the growth pattern in Guilan sheep. However, the Brody model provided the best fit of growth curve in males, and the Richards model for females and all lambs due to the lower values of RMSE, AIC, and BIC and greater values of  $R_{adj}^2$  than the other models. The results of this study show that current growth curve models would be very helpful in genetic breeding programs, flock nutritional management, and decision making on culling of animals with slow growth rate. In this condition, selection could be made for animals with faster growth rate and or that are early-maturing after observing their individual growth curve. After selecting a desired model of the growth curve for this breed of sheep, it is worth noting to develop appropriate genetic selection and management strategies along with the provision of good environmental conditions in the flock to achieve a suitable shape of growth curve via modifying the model parameters.

## Acknowledgements

The corresponding author would like to acknowledge the University of Guilan for financial support of this research (Project No. 989).

## Conflicts of interest

The authors declare they have no conflicts of interest with regard to the work presented in this report.

## References

- Abegaz S, Van Wyk JB, Olivier JJ. Estimation of genetic and phenotypic parameters of growth curve and their relationship with early growth and productivity in Horro sheep. *Arch Tierz* 2010; 53(1):85-94.
- Akbas Y, Taskýn T, Demirören E. Comparison of several models to fit the growth curves of kivircik and daglic male lambs. *Turkish J Vet Anim Sci* 1999; 23 Suppl 3:537-544.
- Bathaei SS, Leroy PL. Growth and mature weight of Mehraban Iranian fat-tailed sheep. *Small Rumin Res* 1996; 22:155-162.
- Bathaei SS, Leroy PL. Genetic and phenotypic aspects of the growth curve characteristics in Mehraban Iranian fat-tailed sheep. *Small Rumin Res* 1998; 29:261-269.
- Bedier NZ, Younis AA, Galal ESE, Mokhta MM. Optimum ewe size in desert Barki sheep. *Small Rumin Res* 1992; 7:1-7.
- Burnham KP, Anderson DR. Model selection and multimodel inference: a practical-theoretic approach, 2nd edn. Berlin, Germany: Springer-Verlag; 2002.
- da Silva LSA, Fraga AB, da Silva FDL, Beelen PMG, Silva RMDO, Tonhati H, Barros CD C. Growth curve in Santa Inês sheep. *Small Rumin Res* 2012; 105:182-185.
- Eteqadi B, Ghavi Hossein-Zadeh N, Shadparvar AA. Population structure and inbreeding effects on body weight traits of Guilan sheep in Iran. *Small Rumin Res* 2014; 119:45-51.
- Fitzhugh Jr HA. Analysis of growth curves and strategies for altering their shape. *J Anim Sci* 1976; 42:1036-1051.
- Freitas AR. Curvas de crescimento na produção animal. *R Bras Zootec* 2005; 34:786795.
- Garnero AV, Marcondes CR, Bezerra LAF, Oliveira HN, Lôbo RB. Genetic parameters of maturation rate and asymptotic weight in Nelore females. *Braz Arch Vet Med Anim Sci* 2005; 57:652-662.
- Gbangboche AB, Gleke-Kalai R, Albuquerque LG, Leroy P. Comparison of non-linear growth models to describe the growth curve in West African Dwarf sheep. *Animal* 2008; 2:1003-1012.

- Ghavi Hossein-Zadeh N. Comparison of non-linear models to describe the lactation curves of milk yield and composition in Iranian Holsteins. *J Agr Sci* 2014; 152:309-324.
- Ghavi Hossein-Zadeh N. Modeling the growth curve of Iranian Shall sheep using non-linear growth models. *Small Rumin Res* 2015; 130:60-66.
- Goliomytis M, Orfanos S, Panopoulou E, Rogdakis E. Growth curves for body weight and carcass components, and carcass composition of the Karagouniko sheep, from birth to 720 d of age. *Small Rumin Res* 2006; 66:222-229.
- Goonewardene LA, Berg RT, Hardin RT. A growth study of beef cattle. *Can J Anim Sci* 1981; 61: 1041-1048.
- Kachman SD, Gianola D. A Bayesian estimator of variance and covariance components in nonlinear growth models. *J Anim Sci* 1984; 59 Suppl 1:176.
- Keskin I, Dag B, Sariyel V, Gokmen M. Estimation of growth curve parameters in Konya Merino sheep. *South Afr J Anim Sci* 2010; 39(2):163-168.
- Lambe NR, Navajas EA, Simm G, Bünge L. A genetic investigation of various growth models to describe growth of lambs of two contrasting breeds. *J Anim Sci* 2006; 84:2642-2654.
- Lewis RM, Emmans GC, Dingwall WS. A description of the growth of sheep and its genetic analysis. *Anim Sci* 2002; 74:51-62.
- Lôbo RNB, Villela LCV, Lôbo AMBO, Passos JRS, Oliveira AA. Genetic parameters of estimated characteristics of the growth curve of Santa Inês sheep. *R Bras Zootec* 2006; 35 (3):1012-1019.
- Malhado CHM, Carneiro PLS, Santos PF, Azevedo DMMR, Souza JC, Affonso PRAM. Curva de crescimento emovinos mestic, os Santa Inês×Texel criados no Sudoeste do Estado da Bahia. *Rev Bras Saúde Prod An* 2008; 9:210-218.
- Malhado CHM, Carneiro PLS, Affonso PRAM, Souza Jr AAO, Sarmiento JLR. Growth curves in Dorper sheep crossed with the local Brazilian breeds, Morada Nova, Rabo Largo, and Santa Inês. *Small Rumin Res* 2009; 84:16-21.
- McManus C, Evangelista C, Fernandes LAC, Miranda RM, Moreno-Bernal FE, Santos NR. Growth curves of Bergamácia Sheep raised in the Federal District. *R Bras Zootec* 2003; 32 (5):1207-1212.
- Noor RR, Djajanegara A, Schüler L. Selection to improvement birth and weaning weight of Javanese Fat Tailed sheep. *Arch Tierz* 2001; 44:649-659.
- Perotto D, Cue RI, Lee AJ. Comparison of nonlinear functions for describing the growth curve of three genotypes of dairy cattle. *Can J Anim Sci* 1992; 72:773-782.
- Rogers SR, Pesti GM, Marks HL. Comparison of three nonlinear regression models for describing broiler growth curves. *Growth* 1987; 51:229-239.
- Sarmiento JLR, Regazzi AJ, Souza WH, Torres RA, Breda FC, Menezes GRO. Study of growth curve of Santa Inês sheep. *R Bras Zootec* 2006; 35 (2):435-442.
- SAS. SAS User's guide v. 9.0: Statistics. SAS Institute, Inc, Cary, NC, USA; 2002.
- Souza LA, Carneiro PLS, Malhado CHM, Silva FF, Silveira FG. Traditional and alternative nonlinear models for estimating the growth of Morada Nova sheep. *R Bras Zootec* 2013; 42(9):651-655.
- Tariq M, Iqbal F, Eydurán E, Bajwa MA, Huma ZE, Waheed A. Comparison of non-linear functions to describe the growth in Mengali sheep breed of Balochistan. *Pak J Zool* 2013; 45(3):661-665.
- Topal M, Ozdemir M, Aksakal V, Yildiz N, Dogru U. Determination of the best nonlinear function in order to estimate growth in Morkaraman and Awassi lambs. *Small Rumin Res* 2004; 55:229-232.
- Toral FLB. Number and range of weighing for estimating parameters of growth curves in cattle. *R Bras Zootec* 2008; 37:2120-2128.