

Inbreeding depression on growth traits of Iranian Lori sheep[□]

Depresión endogámica en las características de crecimiento de la oveja Lori Iraní

Depressão endogâmica de características de crescimento em as ovelhas Lori Iraniana

Zahra Yeganehpur^{1*}, MSc; Hedayatollah Roshanfekr¹, PhD; Jamal Fayazi¹, PhD; Mir Hossein Beyranvand², MSc.

¹Department of Animal Science, Faculty of Animal Science and Food Industries, Khuzestan Ramin Agricultural and Natural Resources University, Ahwaz, Iran.

²Lorestan Agricultural and Natural Resources Research Centre, Khorramabad, Iran.

(Received: October 03, 2015; accepted: July 19, 2016)

doi: 10.17533/udea.rccp.v29n4a03

Summary

Background: Lori sheep is an Iranian heavy breed known for its superiority in terms of disease resistance, adaptability to the mountainous west of the country and meat production potential. **Objective:** to estimate and compare the inbreeding coefficient in Lori sheep, and its impact on growth traits. **Methods:** data and pedigree information were collected in Lorestan Agricultural and Natural Resources Research Center, west of Iran, during 2001-2010. Traits included were birth weight (BW), weaning weight (WW), 6-month weight (6MW) and 9-month weight (9MW). The inbreeding coefficient estimation was carried out through the CFC software, and quantification of individual inbreeding regression on the characteristics was conducted using The Wombat software. All animals were divided into four categories according to the inbreeding coefficients obtained from their pedigree: the first category included non-inbred animals ($F = 0\%$); and the second, third, and fourth categories included inbred animals as $0 < F \leq 0.10$, $0.10 < F \leq 0.20$, and $F > 0.20$, respectively. **Results:** inbreeding coefficients were 0.69% and 2.24% in the entire population and inbred population, respectively. Inbreeding regression for BW, WW, 6 MW and 9 MW were estimated as +4.5, -10.3, -76.3, and -77.4 g, respectively. The inbreeding trend was positive and significant for the whole population (0.215; $p < 0.001$), but not significant for the inbred population. **Conclusion:** these results confirm a low level of inbreeding and suggest that direct controlled mating could be an appropriate method to avoid inbreeding depression.

Keywords: *inbreeding categories, inbreeding effects, mating system, pedigree statistics.*

□ To cite this article: Yeganehpur Z, Roshanfekr H, Fayazi J, Hossein M. Inbreeding depression on growth traits of Iranian Lori sheep. Rev Colomb Cienc Pecu 2016; 29:264-273.

* Corresponding author: Zahra Yeganehpur. Department of Animal Science, Faculty of Animal Science and Food Industries, Khuzestan Ramin Agricultural and Natural Resources University, Ahwaz, Iran. Tel.: + 98 921 397 2191. E-mail: zahra_yeganeh67@mail.com

Resumen

Antecedentes: la oveja Lori es una de las razas pesadas Iraníes más conocidas por su superioridad en términos de resistencia a enfermedades, adaptabilidad a la zona montañosa del oeste del país y potencial para producción de carne. **Objetivo:** estimar y comparar el coeficiente de endogamia en la oveja Lori y su impacto sobre las características de crecimiento. **Métodos:** los datos y la información de pedigrí se recogieron en Lorestan Agricultural and Natural Resources Research Center, al oeste de Irán, desde los años 2001 al 2010. Los rasgos incluidos fueron peso al nacer (BW), peso al destete (WW), peso a los 6 meses (6 MW) y peso a los 9 meses (9 MW). La estimación del coeficiente de endogamia se llevó a cabo mediante el software CFC, y la cuantificación de la regresión de endogamia individual sobre las características se hizo con el software Wombat. Los animales fueron clasificados en cuatro categorías de acuerdo a los coeficientes de endogamia obtenidos de sus pedigrís; la primera categoría incluyó animales no endogámicos ($F = 0\%$); la segunda, tercera y cuarta categorías incluyeron animales endogámicos como $0 < F \leq 0,10$, $0,10 < F \leq 0,20$ y $F > 0,20$, respectivamente. **Resultados:** los coeficientes de endogamia fueron de 0,69 y 2,24% en toda la población y en la población endogámica, respectivamente. La regresión de endogamia para BW, WW, 6 MW, y 9 MW se estimó como +4,5, -10,3, -76,3 y -77,4 g, respectivamente. La tendencia endogámica fue positiva y significativa para toda la población (0,215; $p < 0,001$), pero no significativa para la población endogámica. **Conclusión:** estos resultados confirman un bajo nivel de endogamia y sugieren que el apareamiento directo controlado podría ser un método apropiado para evitar la depresión endogámica.

Palabras clave: *categorías endogámicas, efectos endogámicos, estadísticas de pedigrí, sistema de apareamiento*

Resumo

Antecedentes: as ovelhas Lori são uma das raças pesadas iranianas conhecidas por sua superioridade comparativa sobre outras raças em termos de resistência a doenças, adaptabilidade ao montanhoso oeste do país e potencial para alta produção de carne. **Objetivo:** estimar e comparar o coeficiente de endogamia em ovelhas Lori e seu impacto sobre características de crescimento. **Métodos:** foram coletados dados e informações do pedigree no Lorestan Agricultural and Natural Resources Research Center, na província de Lorestan, oeste do Irã, entre 2001 e 2010. As características incluídas foram o peso ao nascimento (BW), o peso ao desmame (WW), o peso aos 6 meses (6MW), e o peso de 9 meses (9MW). A estimativa do coeficiente de endogamia foi realizada através de software CFC, e a quantificação da regressão de endogamia individual sobre as características foi levada a cabo através do software Wombat. Todos os animais foram classificados em quatro categorias de acordo com os coeficientes de endogamia obtidos por seu pedigree: a primeira categoria incluiu os animais não-endogâmicos ($F = 0\%$); e a segunda, terceira e quarta categorias incluíram os animais endogâmicos como $0 < F \leq 0,10$, $0,10 < F \leq 0,20$ e $F > 0,20$, respectivamente. **Resultados:** os coeficientes de endogamia foram de 0,69% e 2,24%, em toda a população e população consanguínea, respectivamente. A regressão da endogamia para BW, WW, 6 MW e 9 MW foi estimada em +4,5, -10,3, -76,3 e -77,4 g, respectivamente. A tendência da endogamia foi positiva e significativa em toda a população (0,215; $p < 0,001$), mas não significativa na população consanguínea. **Conclusão:** os resultados confirmaram o baixo nível de endogamia e sugerem que a aplicação de um sistema de monta controlada direta poderia ser um método apropriado para evitar a depressão por endogamia.

Palavras chave: *categorias de endogamia, efeitos de endogamia, estatísticas do pedigree, sistema de acasalamento.*

Introduction

Inbreeding is defined as the mating of individuals whose relatedness between them is greater than the average degree of relationship existing in the population (Lush, 1945). Inbreeding seems to be a useful tool to improve the population by increasing the frequency of desirable genes, but the initial consequence of inbreeding is inbreeding depression. Inbreeding depression implies the reduction of

additive genetic variance and phenotypic values for efficiency traits in inbred animals. Nowadays, this is a serious concern in animal breeding in which selection responses are maximized using animal model and Best Linear Unbiased Predictors (BLUP) of breeding value. The inbreeding level needs to be limited to maintain diversity at an acceptable level so as to improve the livestock programs (Van Wyk *et al.*, 2009). Despite some positive inbreeding effect in selecting schemes, breeders are aware of

its deleterious effects and try to avoid it. Inbreeding response is not the same for all animals. There is a wide range of variation in the literature regarding the estimation of inbreeding depression of growth traits (Mirza *et al.*, 1999; Norberg and Sørensen, 2007; Van Wyk *et al.*, 2009; Pedrosa *et al.*, 2010; Ghavi Hossein-Zadeh, 2012; Eteqadi *et al.*, 2014; Mokhtari *et al.*, 2014).

Lori sheep is an important mutton fat-tailed Iranian breed, numbering approximately 2.5 million heads, mainly distributed in the Zagros Mountains of Lorestan province. It can also be found in some areas of Khuzestan, Kermanshah and Ilam provinces (Mohammadi *et al.*, 2015). This breed has an average litter size of 1.25 lambs and its daily gain rate is higher compared to other Iranian sheep breeds. Average mature live weight in this breed is 62 Kg for ewes and 67 Kg for rams; milk yield ranges from 80 to 85 Kg in a 3-month period, approximately, with annual greasy fleece weight between 1.5 to 2 Kg and 2 to 3 Kg in ewes and rams, respectively (Jalali Zonour, 2010).

There are no reports, to our knowledge, evaluating the inbreeding effects on growth traits in Lori lambs. This is the first study evaluating inbreeding depression and its trend on growth traits, to plan optimal designations to select programs. The aim of this study was to estimate and compare the inbreeding coefficient in Lori sheep and its effect on growth traits during a 10-year period.

Materials and methods

Data and management of animals

Data and pedigree information were collected at the Lorestan Agricultural and Natural Resources Research Center, Khorramabad, Iran, from 2001 to 2010. The traits were birth weight (BW), weaning weight (WW), 6-month weight (6MW) and 9-month weight (9MW). The number of available records, mean, standard deviation, coefficient of variation and number of sires and dams are shown in Table 1.

Iranian Lori sheep is a meat purpose breed. Maiden ewes were randomly mated to the rams for the first time at approximate 18 months of age. Ewes were

Table 1. Descriptive statistics for growth traits of Iranian Lori sheep.

	BW	WW	6MW	9MW
No. of records	6440	5646	5073	4757
No. of sires	265	263	258	258
No. of sires with records and progeny in data	175	107	64	112
No. of dams	1794	1801	1777	1684
No. of dams with records and progeny in data	1396	1403	1463	1421
Mean (Kg)	3.49	20.30	25.38	29.80
SD	0.56	1.18	3.65	7.39
Inbreeding mean (F, %)	0.749	0.302	0.204	0.250

BW: birth weight; WW: weaning weight; 6MW: 6-month weight; 9MW: 9-month weight; SD: standard deviation.

usually kept in the flock for six parities (8 years of age) and culling was based on health and low performance. Selected rams were typically 3 to 4 years old and separately kept from ewes. Mating season initiated in early October until mid-November during the whole study. Consequently, lambing was restricted to one season, from mid-February until late March. Single-sire pens were used to allocate 20-25 ewes per ram. All lambs were weighed and ear tagged within 24 h of birth and pedigree and birth information were recorded. After lambing, lambs were separately kept in stalls with their dams for three days. Lambs were fed their mother milk together with good quality alfalfa hay *ad libitum* from 3 weeks of age until weaning, i.e. approximately 90 days. They were raised separately from older animals until 1 year of age. During this period, lambs were sold after weaning, generally due to poor growth or health issues. Weaned lambs were reared by sex group on natural pasture during spring and summer. In the fall, both lambs and ewes grazed on stubble fields of wheat, and barley, then kept indoors during the winter months with a ration composed of wheat barley straw and dry alfalfa, three times per day. The flock was fed a concentrate mixture containing alfalfa, barley grain, and wheat straw during the mating season as a flushing ration.

Statistical analysis

Data and pedigree file were statistically analyzed to assess the inbreeding depression of growth traits in Lori lamb and inbreeding trend. All of the

phenotypic records exceeding 3.5 standard deviations above or below the mean were excluded. The GLM procedure of SAS (Version 9.2, SAS Institute Inc., Cary, NC, USA; 2009) was used to determine the fixed effects, which substantially affect the studied traits. Least square mean was estimated for each trait via the algorithm of Average Information Restricted Maximum Likelihood (AIREML) of the Wombat software (Meyer, University of New England Armidale, NSW 2351, Australia; 2007) by fitting six single trait animal models ignoring or including the additive direct, maternal genetic, and permanent environmental effects. The models included the fixed effects of sex (2 levels, male or female), litter size (2 levels, single or twin), age of ewe at lambing (6 levels, 2 to 7-year-old) and year of birth (10 levels, from 2001 to 2010) and also inbreeding categories and animal random effects. Age of lambs at 3, 6 and 9 months (in days) was considered as a linear covariate for WW, 6MW, and 9MW. Interactions amongst the mentioned fixed effects were not significant, so they were not considered in the final model.

The CFC (contribution, inbreeding and coancestry) software (Sargolzaei *et al.*, Niigata University, Niigata 950-2181, Japan; 2006) was used to calculate the pedigree statistics and analysis of genetic structure of the population.

All the animals were classified in four categories based on the inbreeding coefficient of individual: the first category included non-inbred animals ($F = 0$); second, third, and fourth categories included inbred animals as $0 < F \leq 0.10$, $0.10 < F \leq 0.20$ and $F > 0.20$, respectively. Inbreeding depression was estimated as the regression of growth traits (corrected for fixed

effects) on the inbreeding coefficients of individual lamb using an animal model. Inbreeding trend was also estimated using the regression of the linear individual inbreeding on the year of birth via the Reg procedure of SAS (Version 9.2 SAS Institute Inc., Cary, NC, USA; 2009).

Results

The number of animals (in total), inbred animals, sires, dams, founders, and non-founders in the pedigree were 7,097, 2,126, 273, 1,955, 657, and 6,440, respectively. There were also 942 full sib groups in total, with average family size of 2.06 in the pedigree file. Table 2 illustrates the inbreeding categories. The maximum and minimum number of animals was in the first ($F = 0$) and fourth category ($F > 0.20$). Inbreeding categories were only significant for BW and 6MW for all the traits.

The different inbreeding categories for growth traits by lamb sex and litter size are presented in Tables 3 and 4, respectively. The BW of male and female lambs in different inbreeding categories were significant ($p < 0.05$). Inbreeding categories for BW by litter size, for both male and female and 6MW were significantly different ($p < 0.05$).

As it is shown in Table 3, considerable differences were observed in neither male lambs nor female lambs in the WW. Notwithstanding, non-significant difference amongst the four categories for 6MW female lambs; the males in the third category were lighter than others ($p < 0.05$). On the other hand, the 9MW male lamb was non-significant amongst the four

Table 2. Inbreeding category distribution for growth traits in Lori lambs.

Inbreeding category	BW (Kg)	WW (Kg)	6MW (Kg)	9MW (Kg)
	% Animal (mean \pm SD)	% Animal (mean \pm SD)	% Animal (mean \pm SD)	% Animal (mean \pm SD)
F = 0	66.99 (3.44 \pm 0.01) ^b	93.52(20.30 \pm 0.02) ^a	96.98(25.40 \pm 0.05) ^a	90.24 (29.94 \pm 0.11) ^a
0<F \leq 0.10	31.27 (3.62 \pm 0.01) ^a	5.17 (20.39 \pm 0.07) ^a	2.09 (25.14 \pm 0.34) ^a	8.72 (28.41 \pm 0.36) ^a
0.10<F \leq 0.20	1.47 (3.41 \pm 0.06) ^b	0.99 (20.18 \pm 0.14) ^a	0.61 (23.35 \pm 0.52) ^b	0.84 (29.44 \pm 1.20) ^a
F \geq 0.20	0.27 (3.14 \pm 0.15) ^c	0.32 (19.99 \pm 0.30) ^a	0.32(24.71 \pm 0.97) ^{ab}	0.20 (26.57 \pm 1.90) ^a

BW: birth weight; WW: weaning weight; 6MW: 6-month weight; 9MW: 9-month weight; SD: standard deviation.

Means with the same superscript letter in each subclass within a column do not differ. Significant effect at $p < 0.05$.

inbreeding categories. However, the 9MW female lamb within the fourth category was lighter than the others ($p < 0.05$).

Although BW of single-born lambs within the second category was heavier than that of others, twins pertaining to the fourth category were lighter than those of others ($p < 0.01$; Table 4). However, no significant differences were observed in WW of single and twin-born lambs. The 6MW of single born lambs in the third category was lighter than that of others,

but twins within the fourth category were heavier than those of others ($p < 0.05$). Although 9MW of single born lambs indicated no significant differences in the categories, the twins in the fourth category were heavier than others ($p < 0.05$).

Regression coefficients of growth traits on inbreeding of Lori lambs for a change of 1% in inbreeding are presented in Table 5. By increasing the inbreeding level, 6MW and 9MW significantly decreased to -76.3 g and -77.4 g, respectively ($p < 0.05$).

Table 3. Inbreeding category distribution by sex of Lori lambs.

Birth type	Inbreeding category	BW (Kg)	WW (Kg)	6MW (Kg)	9MW (Kg)
		Mean \pm SD	Mean \pm SD	Mean \pm SD	Mean \pm SD
Male	F = 0	3.54 \pm 0.56 ^{ab}	20.49 \pm 1.30 ^a	26.46 \pm 4.35 ^a	29.61 \pm 9.26 ^a
	0.0 < F \leq 0.10	3.72 \pm 0.60 ^a	20.54 \pm 1.34 ^a	26.93 \pm 4.04 ^a	27.35 \pm 8.66 ^a
	0.10 < F \leq 0.20	3.45 \pm 0.53 ^b	20.23 \pm 1.08 ^a	23.28 \pm 3.78 ^b	32.91 \pm 10.01 ^a
	F \geq 0.20	3.41 \pm 0.39 ^b	20.23 \pm 1.38 ^a	25.32 \pm 4.17 ^{ab}	28.92 \pm 6.45 ^a
Female	F = 0	3.33 \pm 0.50 ^a	20.15 \pm 1.08 ^a	24.78 \pm 3.00 ^a	30.18 \pm 5.53 ^a
	0.0 < F \leq 0.10	3.50 \pm 0.56 ^a	20.30 \pm 1.24 ^a	24.06 \pm 2.74 ^a	29.31 \pm 5.98 ^a
	0.10 < F \leq 0.20	3.56 \pm 0.62 ^a	20.13 \pm 0.99 ^a	23.39 \pm 2.33 ^a	27.57 \pm 5.22 ^a
	F \geq 0.20	2.95 \pm 0.69 ^b	19.74 \pm 0.67 ^a	23.68 \pm 3.48 ^a	23.62 \pm 3.25 ^b

BW: birth weight; WW: weaning weight; 6MW: 6-month weight; 9MW: 9-month weight; SD: standard deviation.

Means with the same superscript letter in each subclass within a column do not differ. Significant effect at $p < 0.05$.

Table 4. Inbreeding category distribution for growth traits by litter size of Lori lambs.

Birth type	Inbreeding category	BW (Kg)	WW (Kg)	6MW (Kg)	9MW (Kg)
		Mean \pm SD	Mean \pm SD	Mean \pm SD	Mean \pm SD
Single	F = 0	3.63 \pm 0.47 ^b	20.46 \pm 1.30 ^a	25.64 \pm 3.66 ^a	29.88 \pm 7.50 ^a
	0 < F \leq 10	3.80 \pm 0.50 ^a	20.54 \pm 1.34 ^a	25.35 \pm 3.83 ^a	28.38 \pm 7.42 ^a
	10 < F \leq 20	3.55 \pm 0.52 ^b	20.25 \pm 1.08 ^a	23.34 \pm 2.81 ^b	29.14 \pm 7.83 ^a
	F \geq 20	3.31 \pm 0.44 ^c	19.97 \pm 1.38 ^a	24.18 \pm 3.83 ^{ab}	25.95 \pm 5.77 ^a
Twin	F = 0	3.01 \pm 0.43 ^a	20.00 \pm 0.80 ^a	24.79 \pm 3.54 ^b	24.79 \pm 3.54 ^b
	0 < F \leq 10	3.04 \pm 0.48 ^a	19.98 \pm 0.92 ^a	24.70 \pm 2.89 ^b	24.70 \pm 2.89 ^b
	10 < F \leq 20	2.93 \pm 0.48 ^a	20.03 \pm 0.94 ^a	23.36 \pm 3.23 ^b	23.36 \pm 3.23 ^b
	F \geq 20	1.9 \pm 0.00 ^b	20.00 \pm 0.00 ^a	28.35 \pm 2.33 ^a	28.35 \pm 2.33 ^a

BW: birth weight; WW: weaning weight; 6MW: 6-month weight; 9MW: 9-month weight; SD: standard deviation.

Means with the same superscript letter in each subclass within a column do not differ. Significant effect at $p < 0.05$.

Table 5. Regression coefficients for growth traits (in grams) on Lori lamb inbreeding for 1% change in inbreeding.

Lambs	BW \pm SE	WW \pm SE	6MW \pm SE	9MW \pm SE
Male	-1.7 \pm 4.6 ^{ns}	-12.5 \pm 12.1 ^{ns}	-92.9 \pm 27.3*	114.6 \pm 119.9 ^{ns}
Female	7.0 \pm 4.0 ^{ns}	-11.0 \pm 10.5 ^{ns}	-83.9 \pm 17.4*	-233.8 \pm 35.2***
Single	0.5 \pm 3.0 ^{ns}	-16.2 \pm 10.1 ^{ns}	-95.7 \pm 18.7**	-109.5 \pm 25.2*
Twin	-6.3 \pm 5.4 ^{ns}	-1.7 \pm 11.3 ^{ns}	-8.1 \pm 13.5 ^{ns}	145.4 \pm 116.7 ^{ns}
All	4.5 \pm 3.1 ^{ns}	-10.3 \pm 8.0 ^{ns}	-76.3 \pm 13.2**	-77.4 \pm 11.3**

BW: birth weight; WW: weaning weight; 6MW: 6-month weight; 9MW: 9-month weight; SD: standard deviation. ^{ns}non-significant, *Significant at $p < 0.05$, **Significant at $p < 0.01$, ***Significant at $p < 0.001$.

Annual mean of inbreeding in whole and inbred populations of Lori lambs are illustrated in Figures 1 and 2, respectively.



**Figure 1.** Annual mean of inbreeding in whole population of Lori lambs.

Figure 1 shows an overall increase in the inbreeding magnitude, irrespective of fluctuations in different years. In Figure 1, the inbreeding trend of growth traits shows an ascending change with mild fluctuations over a 10-year period. The mean of inbreeding decreased in year 2008 due to prevention of closed mating and selection of non-related animals to mate by the breeder. The maximum of inbreeding was observed in 2010 since inbred rams and ewes were mated with each other. The mean of inbreeding for twin lambs was 0 until 2003. In 2010, the mean of inbreeding was 2.19% for all of the lambs, showing an increased trend compared to the base year (2001). In total, there was an overall increase in the magnitude of inbreeding, irrespective of the fluctuations in different years (Figure 1).

Figure 2 shows the non-significant inbreeding trends in inbred population of Lori lambs. In Figure 2, considerable and irregular fluctuations were observed in annual inbreeding mean for male and single-born lambs; there were abrupt increases in the annual

**Figure 2.** Annual mean of inbreeding in inbred population of Lori lambs.

inbreeding mean for some years. Generally, relatively severe fluctuations were observed, especially for male and single-born lambs over a 10-year period, but inbreeding trends were insignificant.

Inbreeding trends (% per year) in whole and inbred populations of Lori lambs are demonstrated in Table 6. In the current study, positive and significant inbreeding trends were only observed in whole population for male, female, single-born lambs, twin-born lambs, and all lambs. Inbreeding trend for all of the animals was 0.215 per year during the 10 years.

Discussion

According to Table 2, BW of lambs within the second category ($0 < F \leq 0.10$) was heavier than other categories ($p < 0.01$); however, the differences were not remarkable between the first and third ($0.10 < F \leq 0.20$) categories. Consistent with the current results, Ghavi Hosseini-Zadeh (2012) reported that BW of Moghani lambs pertaining to $0 < F \leq 0.10$ category was heavier than that of lambs belonging to others. Nevertheless, there were no significant differences between seven

Table 6. Inbreeding trends (% per year) in Lori sheep population.

Lambs	Whole population		Inbred population	
	Inbreeding trend \pm SE	R ² (%)	Inbreeding trend \pm SE	R ² (%)
Male	0.212 \pm 0.04***	78.9	-0.010 \pm 0.006 ^{ns}	23.8
Female	0.223 \pm 0.04***	80.4	-0.001 \pm 0.003 ^{ns}	5.3
Single	0.223 \pm 0.04***	81.1	0.022 \pm 0.01 ^{ns}	38.7
Twin	0.202 \pm 0.04***	76.5	0.001 \pm 0.004 ^{ns}	5.8
All	0.215 \pm 0.04***	79.6	0.003 \pm 0.003 ^{ns}	9.7

R²: regression fit, ^{ns}non-significant (p>0.05), ***Significant effect at p<0.001.

inbreeding categories of BW in the study of Dorostkar *et al.* (2012). Concerning WW, it was not significant amongst four inbreeding categories. This finding was different from the results reported by Ghavi Hossein-Zadeh (2012), indicating that the WW of lambs within F>0.10 inbreeding was heavier than those of other categories (p<0.05). Moreover, heavier WW related to category F \geq 0.25 reported by Dorostkar *et al.* (2012). The 6MW of lambs in the third category of inbreeding was lighter than those of the first and second categories (p<0.05); however, the difference was not significant between the third and fourth categories. Contrary to our findings, Dorostkar *et al.* (2012) reported heavier 6MW relating to F \geq 0.25 in Moghani sheep. Also, there were no considerable differences amongst the four inbreeding categories on 9MW (p>0.05).

Inbreeding depression in males might be further reduced by sex chromosome hemizygoty, affecting the dominance of some mutations, as well as by the absence of sexual selection (Ebel and Phillips, 2016). The inbreeding coefficient is strongly determined by the two main factors, including depth and completeness of pedigree and intensity of selection. Intensity of selection is often increased by reproductive technologies focused on few superior animals, especially rams, and the application of advanced genetic evaluation methods.

Analyzing a total of 8,494 lamb records, Ghavi Hossein-Zadeh (2012) showed no significant differences in BW of single and twin born lambs, WW of single born lambs, 6MW, and 9MW of twins. The author found heavier WW of twin-born lambs in the F>0.10 compared to twins within the first and second categories (p<0.05). Moreover, the same researcher

reported remarkable differences for 6MW and 9MW of single-born lambs in the F>0.10 with the lambs belonging to 0<F \leq 0.10.

The regression coefficient of BW on inbreeding of lambs was positive and negligible (4.5 g, p>0.05; Table 5). Similarly, Dorostkar *et al.* (2012) found non-significant inbreeding depression per 1% increase in inbreeding for BW. Conversely, Van Wyk *et al.* (1993) observed a significant reduction of BW for Elsenburg Dormer lambs as 8 g per 1% increase in inbreeding. Khan *et al.* (1995) and Mirza *et al.* (1999) reported regression coefficients of -8 and -7 g, respectively. Norberg and Sørensen (2007) obtained significant reductions for BW, being 112, 82, and 88 g per 10% increase in Texel, Shropshire, and Oxford Down lamb inbreeding, respectively. Pedrosa *et al.* (2010) found a significant inbreeding depression for BW as 3.4 g per 10% increase in Santa Inês lamb inbreeding. In addition, the findings by Ghavi Hossein-Zadeh (2012), Mokhtari *et al.* (2014), and Eteqadi *et al.* (2014) were not similar to the findings of the present study. Ghavi Hossein-Zadeh (2012) reported that BW was affected by inbreeding of lambs being -9 g (p<0.001). The inbreeding depression of BW obtained as -7 g (p<0.05) using 6,082 Iran-Black sheep records analyzed by Mokhtari *et al.* (2014). Eteqadi *et al.* (2014) obtained inbreeding depression of BW as -5.5 g using 14,549 Guilan lamb records over an 18-year period.

The current results indicate that by increasing the inbreeding level, WW decreased, but not reached significance (-10.3 g, p>0.05). This finding was similar to the study by Mokhtari *et al.* (2014) and incompatible with the study by Dorostkar *et al.* (2012), being -37 and -291 g, respectively. Also, the WW of male, female, single and twin-born lambs was negative and non-significant decreased by 12.5 -11.0, -16.2, and -1.7 g owing to 1% increase in inbreeding, respectively. The studies by Khan *et al.* (1995), Mirza *et al.* (1999), and Akhtar *et al.* (2000) are in agreement with the present results. The WW of Rambouillet lamb was 21.11 \pm 0.15 Kg; it decreased 10 g per 1% increase in inbreeding (Khan *et al.*, 1995). A negligible reduction was found for WW of Lohi lambs as 6.9 g per 1% increase due to inbreeding (Mirza *et al.*, 1999). Conversely, the inbreeding regression of WW was 99 g (p<0.01) in Elsenburg

Dormer lambs (Van Wyk *et al.*, 1993). In contrast, a non-significant decrease was observed as 6 g in WW of Horro lambs (Negussie *et al.*, 2002). However, by increasing the inbreeding level, WW of Hissardale lambs insignificantly increased (Akhtar *et al.*, 2000). Also, considering the sex of lamb and litter size, Eteqadi *et al.* (2014) reported a significant decline of 30.1, 28.9, 27.1, and 75.7 g in WW of male, female, single, and twin-born Guilan lambs per 1% increase in lamb inbreeding, respectively.

The reduction due to inbreeding for 6MW as 76.3 g found in the present study was higher than that observed by Eteqadi *et al.* (2014). Considering lamb sex, there were significant regression coefficients on male and female lamb inbreeding for change of 1% inbreeding by -92.9 and 83.9 g, respectively. Consequently, 6MW male and female lambs decreased to 92.9 and 83.9 g owing to 1% increase in inbreeding. Furthermore, a significant effect was obtained for litter size of 6MW; the regression coefficient of 6MW of single-born lambs on inbreeding was negative, being -95.7 ($p < 0.01$). Hence, 6MW of single-born lambs decreased to 95.7 g due to 1% increase in inbreeding; on the other hand, 6MW of twins was non-significantly negative (-8.1 g, $p > 0.05$). Inbreeding depression of 6MW was up to 79 g (Ghavi Hossein-Zadeh, 2012). Also, the coefficient of inbreeding regression was significantly different in terms of the lamb sex and litter size for 6MW (Ghavi Hossein-Zadeh, 2012). Inconsistent with the current findings, Pedrosa *et al.* (2010) and Eteqadi *et al.* (2014) obtained inbreeding depression of -204 and +20.1 g, respectively. Eteqadi *et al.* (2014) reported significant inbreeding reduction only for 6MW of twin lambs as -413.5 g, although inbreeding depressions were 20.9, 17.1, and 22.3 g for male, female, and single-born lambs, respectively. Moreover, Dorostkar *et al.* (2012) did not report any significant inbreeding effects on 6MW.

Inbreeding depression of 9MW was -77.4 g ($p < 0.01$) per 1% increase in inbreeding coefficient. Also, 9MW of females and single-born lambs decreased up to 233.8 ($p < 0.001$) and 109.5 g ($p < 0.05$), respectively. Similar to the present findings, Eteqadi *et al.* (2014) reported regression coefficient as -45 g per 1% increase in 9MW of individual inbreeding; the non-significant inbreeding depression was -166, -71, and -25 g for twins, males, and females, respectively.

There are few published reports regarding the inbreeding effect on 9MW. There are several methodological and biological factors that determine the estimated inbreeding impact on performance traits. It is well known that both negative and positive effects exist. Hence, in a study undertaken on the population, «bad» and «good» inbreeding effects are mixed (Barczak *et al.*, 2009).

Inbreeding significantly reduced post weaning growth traits, but it had no effect on the pre-weaning period. Thus, inbreeding showed no deleterious effects at early ages, but as the lambs grew older, the effect was remarkable.

Similar findings were reported by Dorostkar *et al.* (2012), as well. Notwithstanding, lower inbreeding trends were assessed by Norberg and Sørensen (2007), Van Wyk *et al.* (2009), Sheikhloo *et al.* (2012) and Rashedi Dehsahraei *et al.* (2013), for Danish, Elsenburg Dormer, Baluchi, and Lori-Bakhtiari sheep breeds, respectively. Concerning inbred population, inbreeding trends were only negative for male and female lambs being -0.010 and -0.001, respectively, which showed a non-significant level ($p > 0.05$).

The difference between the present study and the others might be due to differences amongst breeds, variations of inbreeding level, genetic variation of the base population, population size, and ram to ewe ratio for mating, and different environmental and managing conditions. Also, responding to inbreeding banks on the genetic differences or differentials in levels of inbreeding density and differences between the animals such as age, sex, and litter size. Inbreeding is a special parameter in every population; being different in the various flocks is common. In general, these estimates of inbreeding trend provide a good prospective of the mating system for Lori sheep. Inbreeding level was low in the present study. The deleterious inbreeding effects were not obtained for BW and WW of this breed, but inbreeding depression was observed on 6MW and 9MW, especially for the 9MW females. Continuation of inbreeding trend rising in the whole population over the years warns that mating shall be planned more carefully in the future to avoid mating of close relatives. Furthermore, using the designed mating system could be an appropriate approach to

maintain the inbreeding level under control in the present population.

In this study, positive and significant inbreeding trends were 0.212, 0.223, 0.223, and 0.202 in the whole population for male, female, single born, and twin born lambs, respectively (Table 6). A lower inbreeding trend was observed by Huby *et al.* (2003) and Van Wyk *et al.* (2009). However, a higher inbreeding trend (0.40) was reported by Norberg and Sørensen (2007). The mean of inbreeding for the whole population (0.69) was lower than that in previous reports (Norberg and Sørensen, 2007; Van Wyk *et al.*, 2009; Tahmoorespur and Sheikhloo, 2011; Rashedi Dehsahraei *et al.*, 2013).

Acknowledgements

We would like to thank Parviz Mohammadi Ghalaee, Pharm. D., for providing Spanish and Portuguese versions of the Abstract, and Mr Abouzar Mohammadi, Kashan University of Medical Sciences and Health Services, Kashan and Dr Hassan Khajavy, University of Bojnord, for proofreading our manuscript. Also, we acknowledge the staff of the Agricultural and Natural Resources Research Center of Lorestan province for providing the data for the study.

Conflict of interest

The authors declare they have no conflicts of interest with regard to the work presented in this report.

References

- Akhtar P, Ahmad Z, Mohiuddin G, Abdullah M. Effect of inbreeding on different performance traits of Hissardale sheep in Pak Pak. *Vet J* 2000; 20:169-172.
- Barczak E, Wolc A, Wójtowski J, Slószar P, Szwaczkowski T. Inbreeding and inbreeding depression on body weight in sheep. *Anim Feed Sci* 2009; 18:42-50.
- Dorostkar M, Faraji Arough H, Shodja J, Rafat SA, Rokouei M, Esfandyari H. Inbreeding and inbreeding depression in Iranian Moghani sheep breed. *J Agri Sci Technol* 2012; 14:549-556.
- Ebel ER, Phillips PC. Intrinsic differences between males and females determine sex-specific consequences of inbreeding. *BMC Evol Biol* 2016; 16:36.
- Eteqadi B, Ghavi Hossein-Zadeh N, Shadparvar AA. Population structure and inbreeding effects on body weight traits of Guilan sheep in Iran. *Small Ruminant Res* 2014; 119:45-51.
- Ghavi Hossein-Zadeh N. Inbreeding effects on body weight traits of Iranian Moghani sheep. *Arch Tierzucht* 2012; 55:171-178.
- Huby M, Griffon L, Moureaux S, De Rochambeau H, Danchin Burge C, Verrier E. Genetic variability of six French meat sheep breeds in relation to their genetic management. *Genet Sel Evol* 2003; 35:637-655.
- Jalali Zonour MJ. *Modern Principles of Sheep Husbandry*. 3rd ed. Tehran, Iran (In Persian): Partove Vaghe'e Publishers; 2010.
- Khan MS, Ahmad MD, Ahmad Z, Jadoon JK. Effect of inbreeding on performance traits of Rambouillet sheep. *J Anim Plant Sci* 1995; 5:299-304.
- Lush J L. *Animal Breeding Plans*. Iowa, USA: Iowa State College Press; 1945.
- Meyer K. WOMBAT- A tool for mixed model analyses in quantitative genetics by restricted maximum likelihood (REML). *J Zhej Uni Sci B* 2007; 8:815-821.
- Mirza RH, Abdullah M, Ali I, Hussain R. Effect of inbreeding on body weight in Lohi sheep. *J Anim Plant Sci* 1999; 9: 25-27.
- Mohammadi K, Abdollahi-Arpanahi R, Amraei F, Mirza Mohammadi E, Rashidi A. Genetic parameter estimates for growth and reproductive traits in Lori sheep. *Small Ruminant Res* 2015; 131:35-42.
- Mokhtari MS, Moradi Shahrababak M, Esmailizadeh AK, Moradi Shahrababak H, Gutierrez JP. Pedigree analysis of Iran-Black sheep and inbreeding effects on growth and reproduction traits. *Small Ruminant Res* 2014; 116:14-20.
- Negussie E, Abegaz S and Rege JOE. Genetic trend and effects of inbreeding on growth performance of tropical fat-tailed sheep. *International Proceeding of the 7th World Congress on Genetics Applied to Livestock Production, 2002 Aug 19-23; Montpellier, France, 2002.*
- Norberg E, Sørensen AC. Inbreeding trend and inbreeding depression in the Danish populations of Texel, Shropshire and Oxford Down. *J Anim Sci* 2007; 85:299-304.
- Pedrosa VB, Santana Jr ML, Oliveira PS, Eler JP, Ferraz JBS. Population structure and inbreeding effects on growth traits of Santa Inês sheep in Brazil. *Small Ruminant Res* 2010; 93: 135-139.
- Rashedi Dehsahraei A, Fayazi J, Vatankhah M. Investigating inbreeding trend and its impact on growth traits of Lori-Bakhtiari Sheep. *J Ruminant Res* 2013; 1:65-78.
- Sargolzaei M, Iwaisaki H, Colleau JJ. CFC: a tool for monitoring genetic diversity. *International Proceeding of the 8th World Congress on Genetics Applied to Livestock Production CD-ROM Communication, 2006 Aug 13-18; Belo Horizonte, Brazil, 2006.*
- SAS Institute Inc. *SAS User's Guide: Version 9.2*. SAS Institute Inc., Cary, NC, USA; 2009.

Sheikhloo MR, Tahmoorespur M, Eslaminejad AA. Evaluation of the inbreeding for Baluchi sheep at Abbasabad breeding station, Mashhad. *Iranian J Anim Sci Res* 2012; 3:453-458.

Tahmoorespour M, Sheikhloo M. Pedigree analysis of the closed nucleus of Iranian Baluchi sheep. *Small Ruminant Res* 2011; 99:1-6.

Van Wyk JB, Erasmus GJ, Konstantinov KV. Inbreeding in the Elsenburg Dormer sheep stud. *S Afr J Anim Sci* 1993; 23:77-80.

Van Wyk JB, Fair MD, Cloete SWP. Case study: the effect of inbreeding on the production and reproduction traits in the Elsenburg Dormer sheep stud. *Livest Sci* 2009; 120:218-224.