

Scientific note

Susceptibility of the predator *Euborellia annulipes* (Dermaptera: Anisolabididae) to mycoinsecticides

Susceptibilidad del depredador *Euborellia annulipes* (Dermaptera: Anisolabididae) a micoinsecticidas

FLÁVIA QUEIROZ DE OLIVEIRA¹, JACINTO DE LUNA BATISTA¹, JOSÉ BRUNO MALAQUIAS³, CARLOS HENRIQUE DE BRITO⁴, and EMMERSON PEREIRA DOS SANTOS⁵

Abstract: To evaluate the influence of *Beauveria bassiana* (Balsamo) Vuill and *Metarhizium anisopliae* (Metsch) Sorok on the survival of the predator *Euborellia annulipes* Lucas (Dermaptera: Anisolabididae) were used product concentrations 5.00×10^9 ; 7.50×10^9 ; 10.00×10^9 ; 12.50×10^9 conidia/L from isolates of *B. bassiana* and *M. anisopliae* which were obtained, respectively, through products Boveril® and Metaril® and control (0.0 conidia/L). High rates of hatching nymphs of *E. annulipes* have been recorded in clutches that received applications of *M. anisopliae*. Moreover, *B. bassiana* has affected negatively the hatching rate of nymphs of *E. annulipes*. A slightly harmful effect has been observed in all concentrations of conidia *B. bassiana* in clutches of *E. annulipes*. The survival was 100% in all tested concentrations of *M. anisopliae* in females of *E. annulipes*. The females which received topical application of *B. bassiana* have shown a survival rate ranging from 80.30 (12.50×10^9 conidia) to 100% (5.00×10^9 conidia). In males of *E. annulipes*, the survival rate in females treated with *B. bassiana* ranged from 95.00 (12.50×10^9 conidia) to 100% (5.00×10^9 conidia), while with *M. anisopliae* applications this rate was 96.02 (12.50×10^9 conidia) to 100% (5.00×10^9 conidia). The entomopathogenic fungi *B. bassiana* and *M. anisopliae* did not affect the mortality of nymphs and adults of *E. annulipes*. However, careful measures should be adopted in applications of *B. bassiana* directly into clutches of *E. annulipes*.

Key words: Pathogenicity. Entomopathogenic fungi. Selectivity. Natural enemy.

Resumen: Se evaluó la influencia de *Beauveria bassiana* (Balsamo) Vuill y *Metarhizium anisopliae* (Metsch) Sorok en la supervivencia del depredador *Euborellia annulipes* Lucas (Dermaptera: Anisolabididae). Las concentraciones de los productos evaluados fueron $5,00 \times 10^9$; $7,50 \times 10^9$; $10,00 \times 10^9$; $12,50 \times 10^9$ conidios/L de aislamientos de *B. bassiana* y *M. anisopliae*, respectivamente, con los productos Metaril® y Boveril® y un control (0,0 conidios / L). Se registraron altas tasas de eclosión de ninfas de *E. annulipes* en especímenes que recibieron tratamientos con *M. anisopliae*. Se observó un efecto levemente perjudicial en todas las concentraciones de conidios de *B. bassiana* en posturas de *E. annulipes*, pues *B. bassiana* afectó negativamente la tasa de eclosión de ninfas de *E. annulipes*. La supervivencia fue del 100% en hembras de *E. annulipes* que recibieron todas las concentraciones probadas de *M. anisopliae*. Las hembras que recibieron aplicaciones tópicas de *B. bassiana* presentaron sobrevivencia entre 80,3 ($12,50 \times 10^9$ conidios) y 100% ($5,00 \times 10^9$ conidios). Los machos de *E. annulipes* presentaron una tasa de supervivencia con *B. bassiana* entre 95 ($12,50 \times 10^9$ conidios) y 100% ($5,00 \times 10^9$ conidios) y de 96,0 ($12,50 \times 10^9$ conidios) a 100% ($5,00 \times 10^9$ conidios) al recibir aplicaciones de *M. anisopliae*. Los hongos entomopatógenos *B. bassiana* y *M. anisopliae* no afectaron la mortalidad de ninfas y adultos de *E. annulipes*. Sin embargo, se deben adoptar medidas precisas en aplicaciones de *B. bassiana* directamente en posturas de *E. annulipes*.

Palabras clave: Patogenicidad de hongos. Hongos entomopatógenos. Seletividad. Enemigo natural.

Introduction

Earwigs or Dermaptera constitute several species with high predation capacity due to their aggressive behavior. Among these species, *Euborellia annulipes* (Lucas, 1847) (Dermaptera: Anisolabididae) is an important biological control agent of pests in agriculture. This insect species occurs throughout the neotropical zoogeographical region and has been an potential predator of several insect pests, particularly of eggs and immature stages of insects of the Lepidoptera, Hemiptera, Coleoptera and Diptera (Lemos *et al.* 1998, 1999). In northeast Brazil it is common occurrence of this insect preying on immature stages of pests that occur in soil or in plant structures lying on the ground as case of the larvae and pupae of boll weevil at field of cotton.

Entomopathogens are prevalent in natural systems and should receive greater attention in life-history studies (Hesketh *et al.* 2010; Roy *et al.* 2009). Mycoinsecticides compose only a small component of the biopesticide market, but their use is gaining acceptance as an alternative to the use of traditional contact insecticides for pest control (Ludwig and Oetting 2001). *Beauveria bassiana* (Balsamo) Vuill and *Metarhizium anisopliae* (Metsch) Sorok are among the most studied species of entomopathogenic and are used as bio-control agents of agricultural pests (Alves 1992; Alves *et al.* 1998). In biological control of pests an important pathogen is that one who affects only the target insect and not their natural enemies (Flexner *et al.* 1986; Generoso 2002).

The severity of the effects of microbial insecticides depends on the concentration of spores applied and the stage of

¹ M.Sc. Student in Environmental Science and Technology, Universidade Estadual da Paraíba (UEPB), Brazil. e-mail: flavinha2010@ibest.com.br corresponding author; ² Ph.D., Universidade Federal da Paraíba (UFPB). Campus II - Areia - PB. e-mail: jacinto@cca.ufpb.br; ³ M.Sc. Student in Entomology, Universidade de São Paulo (USP), Brazil. e-mail: jbmalaquias@ig.com.br; ⁴ Researcher/Professor, Universidade Federal da Paraíba (UFPB), Campus II - Areia - PB. e-mail: chbrito1@hotmail.com; ⁵ B.Sc. Universidade Federal da Paraíba (UFPB), Campus II - Areia - PB. e-mail: emersonbiozoo@hotmail.com.

the insect (Ginsberg *et al.* 2002). The use of low concentrations of alternative insecticides may increase the selectivity and help reduce costs of production (Oliveira 2008). In recent studies conducted by Oliveira *et al.* (2010), it was observed that low concentrations of mycoinsecticides isolates of *B. bassiana* and *M. anisopliae* promoted high efficiency control of larvae and pupae of fruit flies. However the direct effects of entomopathogenic fungi in natural enemies have not been well studied and probably have been overestimated. It is also necessary to develop techniques to express accuracy in doses of microbial insecticides to achieve maximum physiological compatibility. Due to the increased use of biological control in integrated pest management and considering that information regarding the selectivity of entomopathogenic fungi in biological control agents is incipient, the aim of this research is to evaluate the influence of *M. anisopliae* and *B. bassiana* in the survival of the predator *E. annulipes*.

Material and Methods

Insects. Specimens of the *E. annulipes* used in this study were obtained from laboratory rearing. The stock colony was maintained in an acclimatized room at $26\pm 1^\circ\text{C}$, of $70\pm 10\%$ r.h. and 12L:12D photoperiod. The bioassays followed the methodology described by Lemos *et al.* (1999). This study used eggs, nymphs of 1st instar, ($\leq 24\text{h}$ old), 2nd instar, and adult insects of both sexes of *E. annulipes*. Individuals have been placed in Petri dishes and kept at a constant temperature of $25\pm 1^\circ\text{C}$, with $70\pm 10\%$ R.H., and a photoperiod of 12:12h (L:D). The insects or clutches has been kept in groups of 5 units per dish. Recently emerged adults ($\leq 24\text{h}$ old) have been sexed by the characteristics of external morphology, the number of abdominal segments and the curvature of the forceps (Tomkins and Simmons 1998).

Bioassay. The following concentrations of conidia tested were 5.00×10^9 , 7.50×10^9 , 10.00×10^9 , 12.50×10^9 conidia/L, and control (0.0 conidia/L). These concentrations are recommended to the pests control, i.e. in applications at soil to control of larvae and pupae of fruit flies (Oliveira 2008). The isolates have been obtained from the products Boveril[®] and Metarril[®], produced by the Company Itaforte Bio products.

Pathogenicity and virulence of isolates. The viability of isolated evidence of pathogenicity and virulence has been evaluated according to the method described by França *et al.* (2006). In the viability test has been used two Petri dishes containing PDA (potato dextrose agar) culture media, incubated in climatized chamber ($25\pm 2^\circ\text{C}$, 12h of photophase and relative humidity of $70\pm 10\%$) for 24 hours, later to perform readings in a light microscope by determining the percentage of germinated conidia and not germinated. For proving the pathogenicity and virulence of the isolates tested in this study, we spraying a suspension of 5.00×10^9 conidia/L on the 100 larvae 1st-instar of *Ceratitis capitata* (Wiedemann, 1824) (Diptera: Tephritidae) for each species of fungus, because it is a highly susceptible to entomopathogenic stage (Oliveira 2008).

Effects on survival and development stages of *E. annulipes*. Fungal suspensions of each fungal product concentration have been applied topically with deposition of $5\mu\text{l}$ from each suspension. The applications have been performed using a

manual spray (1L) just once time on the entire structure of the dorsal body of the specimens, and in the case of the clutches around the whole egg chorion. After 10 days of period application was observed the survival rate. This period was established because the action time of two microorganisms have been also observed and the assessments made after 5 days of application to the insects as young and adult (Bustillo *et al.* 1999) as well as after 10 days, which was the maximum period of hatching in this conditions (Lemos 1998). The mortality was confirmed by sporulation in humide chamber. Then, mortality has been calculated and corrigid to control (un-treated) group.

Data analysis. The experimental design was completely randomized with five replications per concentration/treatment, with each replication consisting of five insects or clutches (100 eggs per clutch). The survival of insects in treatments was calculated using the formula: $S (\%) = 100 - Ma (\%)$. The mortality data ($Ma (\%)$) of the treatments and control were submitted to Abbott's formula (Abbott 1925). Data were also subjected to polynomial logistic regression ($P=0.05$) (PROC GENMOD, SAS Institute 2006). The determination of the harmfulness of the products was based on the guidelines for the insecticides under laboratory conditions by the IOBC/WPRS (Amano and Haseeb 2001): survival $>70\%$, non-harmful, survival $>30\%$ and $<70\%$, slightly harmful; survival $>1\%$ and $<30\%$, moderately harmful and survival $<1\%$ severely harmful.

Results and Discussion

In our test with *C. capitata*, it has been recorded a mortality rate approximately 98% and 97% caused by fungi *M. anisopliae* and *B. bassiana*, respectively. Therefore, the survival of *E. annulipes* was not because the lack of fungi virulence.

High rates of hatching nymphs of *E. annulipes* have been recorded in clutches that received applications of *M. anisopliae*. The hatching percentage nymphs confirmed was 96.23, 92.47, 91.58 and 91.50% in the concentrations of 5.00×10^9 , 7.50×10^9 , 10.00×10^9 and 12.50×10^9 conidia/L of *M. anisopliae*, respectively. Moreover, *B. bassiana* affected the hatching rate of nymphs of *E. annulipes*, the survival ranged from 31.13 (12.50×10^9) to 60.06% (5.00×10^9) (Fig. 1). Results




Figure 1. Hatching percentage *Euborellia annulipes* nymphs with conidia concentrations/L of *Metarhizium anisopliae* (—○— Ma) ($y = 0.92x^2 - 6.108x + 101.32$, $R^2 = 0.98$) and *Beauveria bassiana* (---●--- Bb) ($y = 0.4408x^2 - 11.188x + 103.55$, $R^2 = 0.91$). Observed \pm S.E. (dots) and predict results (line models).




Figure 2. Relationship between the application of different concentrations of conidia of fungi *Metarhizium anisopliae* (—○—Ma) ($y = 0.025x^2 - 0.045x + 96.375$, $R^2 = 0.96$) and *Beauveria bassiana* (—●—Bb) ($y = 0.004x^2 - 0.002x + 96.395$, $R^2 = 0.976$) on the survival of males of *E. annulipes* **A**, and the different concentrations of conidia of fungi *M. anisopliae* (—○—Ma) ($y = 100$) and *B. bassiana* (—●—Bb) ($y = 0.0016x^2 - 0.0744x + 80.916$, $R^2 = 0.958$) on the survival of females of *E. annulipes* **B**. Observed \pm S.E. (dots) and predict results (line models).

obtained by Duso *et al.* (2008) with the predatory mite *Phytoseiulus persimilis* (Athias-Henriot, 1957) (Acari: Phytoseiidae) have shown high hatching rate of this predator species after application of *B. bassiana*, although this researchers have reported mortality of up to 43% in females.

In the other hand, *M. anisopliae* and *B. bassiana* did not affect the survival of nymphs. There was overall survival (100%) of nymphs of 1st and 2nd instars in all tested concentrations of both the fungi *B. bassiana* and *M. anisopliae* (data not shown in figures). Results obtained by Kohno *et al.* (2007) are in agreement with those obtained in this study. These authors verified no effect of *B. bassiana* to nymphs of 2nd instar *Labidura riparia* (Pallas, 1773) (Dermaptera, Labiduridae).

The fungi *M. anisopliae* and *B. bassiana* did not affect the survival of adults of *E. annulipes*. In all tested concentrations of *M. anisopliae* in female *E. annulipes*, survival was 100%. The females which received topical application of *B. bassiana* have been shown a survival rate ranging from 80.3% (12.10×10^9 conidia) up to 100% (5.00×10^9 conidia); whereas, for males of *E. annulipes*, the survival rate in insects treated with *B. bassiana* ranged from 95% (12.10×10^9 conidia) to 100% (5.00×10^9 conidia) and from 96.02% (12.10×10^9 conidia) to 100% (5.00×10^9 conidia) after receiving applications of *M. anisopliae* (Figs. 2a and 2b).

Efficient selectivity is based on physiological differences between pests and their natural enemies, being that pest killed at a concentration of product that does not affect individual benefits. A slightly harmful effect (survival > 30% and < 70%) has been observed in all the concentrations of conidia *B. bassiana* in clutches of *E. annulipes* (Fig. 1). The microbial insecticides tested were not harmful (survival > 70%) for adults (Figs. 2A and 2B) and nymphs of first and second instar. Therefore, the two mycoinsecticides can be used to control pests without significantly affecting nymphs or adults of *E. annulipes*. The greater susceptibility of eggs of *E. annulipes* in relation to nymphs or adults should be the interaction that occurs due to pathophysiological effects on the cuticle of nymphs or adults of the predator, which hindered the process of infection and lessens the potential for invasion of microorganisms for these stages of *E. annulipes*. Moreover, the ability of this insect to survive in areas of land, soil inhabitant galleries can be an important factor to provide

a greater hardening of their cuticle. Other factors such as intestinal pH, biochemical or immunological defense-physiological may be related to the tolerance of nymphs or adults of *E. annulipes* concentrations of mycoinsecticides used in this study.

According Kiselek (1975), *B. bassiana* was harmless to adults *Cryptolaemus montrouzieri* Mulsant, 1853 (Coccinellidae: Coleoptera) while the effect was lethal to larvae (50%) that consumed insects infected by this fungus. In this study, nymphs and adults of *E. annulipes* were not affected by concentrations tested in this study adopted of *B. bassiana* and *M. anisopliae*. In other words, the results of this study have revealed that *B. bassiana* caused a reduction in the rates of hatched nymphs, unlike *M. anisopliae*. Moreover, insecticide applications that are not directly to the soil may enable the adoption of that microorganism in programs of integrated pest management in agroecosystems in which *E. annulipes* is present, because this kind of predator tends to deposit their eggs in soil galleries built by adults (Bharadwaj 1966).

Other studies have also demonstrated the selectivity of entomopathogenic organisms to natural enemies, like *B. bassiana* with *Orius insidiosus* (Say, 1832) (Hemiptera: Anthocoridae) (Cavalcanti 2006); *M. anisopliae* and *B. bassiana* with parasitoid *Oomyzus sokolowskii* (Kurdjumov, 1912) (Hymenoptera: Eulophidae) (Santos Jr. *et al.* 2006). The concentrations adopted in this study of entomopathogenic fungi *B. bassiana* and *M. anisopliae* did not affect the mortality of nymphs and adults of *E. annulipes*. Neither we found ovidal activity of *M. anisopliae* to *E. annulipes*. However, careful measures should be adopted in applications from 5.0×10^9 conidia/L of *B. bassiana* directly into clutches of *E. annulipes*, because from this concentration we found a significant reduction in hatching rate this predator. We also suggest the implementation of other studies, as the effects these bioinsecticides on the fitness of *E. annulipes*.

Acknowledgements

The authors thank the Brazilian agency Coordenadoria de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) to concession scholarship to first author. The authors would like to thank Robério de Oliveira for his help in the laboratory.

We especially thank Tricia Hornsby (Ohio State University) for her valuable help with an important review of English in this manuscript.

Literature Cited

- ABBOTT, W. S. 1925. A method of computing the effectiveness of an insecticide. *Journal of Economic Entomology* 18: 265-267.
- ALVES, S. B. 1992. Perspectivas para utilização de fungos entomopatogênicos no controle de pragas no Brasil. *Pesquisa Agropecuária Brasileira* 27: 77-86.
- ALVES, S. B.; ALMEIDA, J. E. M.; ALMEIDA, A.; MOINO JR., L. F.; ALVES, A. 1998. Técnicas de laboratório. In: ALVES, S. B. (ed.). *Controle microbiano de insetos*, 637-711. Piracicaba, Fealq.
- AMANO, H.; HASEEB, M. 2001. Recently-proposed methods and concepts of testing the effects of pesticides on the beneficial mite and insect species: study limitations and implications in IPM. *Applied Entomology and Zoology* 36: 1-11.
- BHARADWAJ, R. K. 1966. Observation on the bionomics of *Euborellia annulipes* (Dermoptera: Labiduridae). *Annals of the Entomological Society of America* 59: 441-450.
- BUSTILLO, A. E.; BERNAL, M. G.; BENAVIDES, P.; HAVES, B. 1999. Dynamics of *Beauveria bassiana* and *Metarhizium anisopliae* infecting *Hypothenemus hampei* (Coleoptera: Scolytidae) populations emerging from fallen coffee berries. *Florida Entomologist* 82: 491-498.
- CAVALCANTI, R. S. 2006. Associação *Beauveria bassiana* (Bals.) Vuill. - nematóides entomopatogênicos (Rhabditida) - *Orius insidiosus* (Say) no controle de trips (Thysanoptera) em cultivo protegido. Tese (Doutorado). Universidade Federal de Lavras, 132p.
- DUSO, C. V.; MALAGNINI, A.; POZZEBON, M.; CASTAGNOLI, M.; LIGUORIB, S. S. 2008. Comparative toxicity of botanical and reduced-risk insecticides to mediterranean populations of *Tetranychus urticae* and *Phytoseiulus persimilis* (Acari Tetranychidae, Phytoseiidae). *Biological Control* 47: 16-21.
- FLEXNER, J. L.; LIGHTHART, B.; CROFT, B. A. 1986. The effects of microbial pesticides on non-target, beneficial arthropods. *Agriculture, Ecosystems & Environment* 16: 203-254.
- FRANÇA, I. W. B.; MARQUES, E. J.; TORRES, J. B.; OLIVEIRA, J. V. 2006. Efeitos de *Metarhizium anisopliae* (Metsch.) Sorok. e *Beauveria bassiana* (Bals.) Vuill. sobre o percevejo predador *Podisus nigrispinus* (Dallas) (Hemiptera: Pentatomidae). *Neotropical Entomology* 35: 349-356.
- GENEROSO, A. R. 2002. Compatibilidade de *Beauveria bassiana* e *Paecilomyces fumosoroseus* com *Chrysoperla externa* (Neuroptera: Chrysopidae) e metodologia para avaliação da seletividade. Dissertação (Mestrado) - Universidade Estadual Paulista, Jaboticabal. 63p.
- GINSBERG, H. S.; LEBRUN, R. A.; HEYER, K.; ZHIOUA, E. 2002. Potential nontarget effects of *Metarhizium anisopliae* (Deuteromycetes) used for biological control of ticks (Acari: Ixodidae). *Environmental Entomology* 31: 1191-1196.
- HESKETH, H.; ROY, H. E.; EILENBERG, J.; PELL, J. K. HAILS, R. S. 2010. Challenges in modeling complexity of fungal entomopathogens in semi-natural populations of insects. *Biocontrol* 55: 55-73.
- KISELEK, E. V. 1975. The effect of biopreparations on insect enemies. *Zash Rast*, 23p.
- KOHNO, K.; TAKEDA, M.; HAMAMURA, T. 2007. Insecticide susceptibility of a generalist predator *Labidura riparia* (Dermoptera: Labiduridae) *Applied Entomology and Zoology* 42: 501-505.
- LEMONS, W. P.; MEDEIROS, R. S.; RAMALHO, F. S. 1998. Influência da temperatura no desenvolvimento de *Euborellia annulipes* (Lucas) (Dermoptera: Anisolabidae) predador de bicudo-do-algodoeiro. *Anais da Sociedade Entomológica do Brasil* 27: 67-76.
- LEMONS, W. P.; MEDEIROS, R. S.; RAMALHO, F. S. 1999. Exigências térmicas de *Euborellia annulipes* (Lucas) (Dermoptera, Anisolabidae) e sua relação com a presa: bicudo-do-algodoeiro. *Revista Brasileira de Entomologia* 43: 61-68.
- LUDWIG, S. W.; OETTING, R. D. 2001. Susceptibility of natural enemies to infection by *Beauveria bassiana* and impact of insecticides on *Ipheseius degenerans* (Acari: Phytoseiidae), *Journal of Agricultural and Urban Entomology* 18 (3): 169-178.
- OLIVEIRA, F. Q. 2008. Eficiência dos fungos *Beauveria bassiana* e *Metarhizium anisopliae* no controle de moscas das - frutas *Ceratitis capitata* (Diptera: Tephritidae). BSc em Agronomia. Universidade Federal da Paraíba. 37p.
- OLIVEIRA, F. Q.; BATISTA, L. J.; MALAQUIAS, J. B.; ALMEIDA, D. M.; OLIVEIRA, R. 2010a. Determination of the median lethal concentration (LC50) of mycoinsecticides for the control of *Ceratitis capitata* (Diptera: Tephritidae). *Revista Colombiana de Entomologia* 36 (2): 213-216.
- ROY, H. E.; HAILS, R. S.; HESKETH, H.; ROY, D. B.; PELL, J. K. 2009. Beyond biological control: non-pest insects and their pathogens in a changing world. *Insect Conservation and Biodiversity* 2: 65-72.
- SANTOS JR., R.; MARQUES, H. J. G.; BARROS, E. J. R.; GONDIM JR., M. G. C.; ZAGO, H. B.; SILVA, C. M. 2006. Efeito de *Metarhizium anisopliae* (Metsch.) Sorok. e *Beauveria bassiana* (Bals.) Vuill. sobre adultos de *Oomyzus sokolowskii* (Kurdjumov) (Hymenoptera: Eulophidae). *Acta Scientiarum. Agronomy* 28: 241-245.
- SAS. 2006. SAS User's Guide: Statistics. SAS Institute, Cary, NC, USA.
- TOMKINS, J. L.; SIMMONS, L.W. 1998. Female choice and manipulations of forceps size and symmetry in the earwig *Forficula auricularia* L. *Animal Behavior* 56: 347-356.

Received: 23-dec-2010 • Accepted: 2-jun-2011