

Esophagogastric Junction Contractile Integral (EGJ-CI) in Various Phenotypes of Gastroesophageal Reflux Disease (GERD)

Óscar Mariano Pinto-Saavedra,¹  Andrés Felipe Ardila-Hani,²  Albis Cecilia Hani-Amador,³  Gerardo Andrés Puentes-Leal,⁴ 
Óscar Mauricio Muñoz-Velandia.^{5*} 

OPEN ACCESS

Citation:

Pinto-Saavedra OM, Ardila-Hani AF, Hani-Amador AC, Puentes-Leal GA, Muñoz-Velandia OM. Esophagogastric Junction Contractile Integral (EGJ-CI) in Various Phenotypes of Gastroesophageal Reflux Disease (GERD). *Revista. colomb. Gastroenterol.* 2023;38(4):460-466. <https://doi.org/10.22516/25007440.1066>

¹ Gastroenterology fellow, Pontificia Universidad Javeriana. Gastroenterology Unit, Hospital Universitario San Ignacio. Bogotá, Colombia.

² Pontificia Universidad Javeriana, Gastroenterology Unit. Hospital Universitario San Ignacio, Department of Internal Medicine. Bogotá, Colombia.

³ Professor, Pontificia Universidad Javeriana. Gastroenterology Unit and Department of Internal Medicine, Hospital Universitario San Ignacio. Bogotá, Colombia.

⁴ Gastroenterologist, Pontificia Universidad Javeriana. Bogotá, Colombia. Hospital Serena del Mar. Cartagena de Indias, Colombia.

⁵ Associate Professor, Department of Internal Medicine, Pontificia Universidad Javeriana. Department of Internal Medicine, Hospital Universitario San Ignacio. Bogotá, Colombia.

*Correspondence: Óscar Mauricio Muñoz-Velandia.
o.munoz@javeriana.edu.co

Received: 11/04/2023

Accepted: 30/06/2023



Abstract

Introduction: Two parameters of high-resolution esophageal manometry are used to observe the function of the esophagogastric junction (EGJ): the anatomical morphology of the EGJ and contractile vigor, which is evaluated with the esophagogastric junction contractile integral (EGJ-CI). To date, how these parameters behave in different gastroesophageal reflux disease (GERD) phenotypes has not been evaluated. **Materials and methods:** An analytical observational study evaluated patients with GERD confirmed by pH-impedance testing and endoscopy undergoing high-resolution esophageal manometry. The anatomical morphology of the EGJ and EGJ-CI was assessed and compared between reflux phenotypes: acid, non-acid, erosive, and non-erosive. **Results:** 72 patients were included (63% women, mean age: 54.9 years), 81.9% with acid reflux and 25% with erosive esophagitis. In the latter, a decrease in EGJ-CI (median: 15.1 vs. 23, $p = 0.04$) and a more significant proportion of patients with type IIIa and IIIb EGJ (83.3% vs 37.1%, $p < 0.01$) were found. No significant differences existed in the manometric parameters of patients with and without acid and non-acid reflux. **Conclusion:** In our population, EGJ-CI significantly decreased in patients with erosive GERD, suggesting that it could be used to predict this condition in patients with GERD. This finding is also related to a higher proportion of type III EGJ and lower pressure at end-inspiration of the lower esophageal sphincter in this reflux type.

Keywords

Gastroesophageal reflux disease, esophagogastric junction, lower esophageal sphincter.

INTRODUCTION

The measurement and quantification of the functionality of the esophagogastric junction (EGJ) allow the establishment of adequate competence of this barrier mechanism. They can provide valuable information on conditions or pathologies in which it can be altered, such as gastroesophageal reflux disease (GERD)^(1,2), postoperative states

of myotomy in patients with achalasia, postoperative states of anti-reflux surgery, among others.

The Lyon Consensus proposes two parameters of esophageal high-resolution manometry (HRM) to observe the EGJ's function: the anatomical morphology of the EGJ and contractile vigor^(3,4). The morphology of the EGJ is defined by the relationship between the lower esophageal sphincter (LES) and the crura of the diaphragm (CD). Three types of

EGJ morphology are proposed: Type 1 when the LES and CD are superimposed, Type 2 when the LES and CD are separated by less than 2 cm, and Type 3 when the separation is greater than 2 cm⁽³⁻⁵⁾. The contractile vigor of the EGJ or esophagogastric junction contractile integral (EGJ-CI) is calculated using a similar method to that of the distal contractile integral (DCI). The DCI frame is placed over the EGJ to include three respiratory cycles; the average value recorded in mm Hg/cm/s is divided by the duration of the three respiratory cycles to obtain the EGJ-CI in mm Hg/cm (Figure 1)^(3,6).

The study by Nicodeme et al. evaluated the severity of gastroesophageal reflux defined by pH impedance variables such as acid exposure time (AET), the number of reflux episodes in 24 hours, and a symptomatic index greater than 50%. They found that a greater EGJ-CI in the HRM was associated with fewer reflux episodes⁽⁶⁾.

Although different GERD phenotypes have been described based on clinical, pH impedance, and endoscopic parameters, such as acid, non-acid, erosive, and non-erosive ref-

lux, it has not been possible to define factors that can predict these phenotypes. Specifically, it has not been determined whether the assessment of the EGJ competence, especially the EGJ-CI, could be helpful in this context. Defining it would be clinically relevant since the treatment and prognosis of each phenotype differs significantly⁽⁷⁾.

The present study aims to describe the function of the EGJ, including anatomical morphology and EGJ-CI in various GERD phenotypes, using pH-impedance testing and upper GI tract endoscopy as the gold standard.

MATERIALS AND METHODS

We conducted an analytical observational study, which included patients with a diagnosis of GERD confirmed by pH impedance or endoscopic findings that were taken for esophageal manometry in the gastroenterology unit of the Hospital Universitario San Ignacio, a referral hospital in Bogotá (Colombia), between June 2019 and June 2021. Patients over 18 with simultaneous HRM, pH-impedance




Figure 1. The distal contractile integral frame (red frame) is placed over the EGJ to include three respiratory cycles; the value recorded in mm Hg/cm/s is divided by the duration of the three respiratory cycles to obtain the EGJ-CI in mm Hg/cm. Authors' archives.

testing, and upper GI endoscopy were included. Patients with incomplete manometries for any reason, a history of upper GI surgery (total or partial gastrectomy), esophageal interventions, achalasia, outflow tract obstruction, Jackhammer, and hypercontractile esophagus, as well as patients who did not meet the GERD criteria, were excluded. The Hospital Universitario San Ignacio and the Pontificia Universidad Javeriana ethics committee approved the study under code FM-CIE-1992-21/246/2021.

The endoscopic procedures (esophagogastroduodenoscopy) were performed by gastroenterologists attached to the Hospital Universitario San Ignacio. The manometry and pH-impedance studies were conducted by duly trained nursing staff, and gastroenterology fellows analyzed them under the supervision and approval of certified gastroenterologist members of the digestive physiology group. The manometries were analyzed according to the parameters of the Chicago Classification v4.0 and the pH-impedance tests according to the guidelines of the Lyon Consensus^(15,16). Demographic data and the results of the different tests were obtained from the records systematically filled out in the unit and collected using a standardized form.

Non-acid reflux was defined as more than 27 reflux episodes when the patient came without treatment and more than 44 reflux episodes if the patient came with treatment. Acid reflux was determined as AET > 6%, or, in patients with AET in the gray zone (between 4% and 6%), as a number of acid refluxes greater than 80, a DeMeester score greater than 14.7, a post-reflux swallow-induced peristaltic wave index greater than 61% and the mean nocturnal baseline impedance less than 2,292 ohms. Non-erosive reflux was defined as those patients with routine endoscopy, and erosive reflux as the presence of Grade C and D esophagitis according to the Los Angeles Classification, Barrett's esophagus, or peptic stricture⁽¹⁶⁾.

The EGJ-CI measurement used the DCI frame placed over the EGJ to include three respiratory cycles. The value recorded in mm Hg/cm/s is divided by the duration of the three respiratory cycles to obtain the EGJ-CI in mm Hg/cm (Figure 1).

Continuous variables were described using measures of central tendency and dispersion. A Shapiro-Wilk test was used to define whether the normality criterion in the data distribution was met. If met, the mean and standard deviation were described; otherwise, the median and interquartile range. Categorical variables were defined with absolute numbers and percentages. The groups generated according to the erosive, non-erosive, acid, and non-acid phenotypes were compared using a t-test or a Mann-Whitney U test. A Stata 16 statistical package was used for the analysis.

RESULTS

Seventy-two patients were included, with a mean age of 54.9 years (standard deviation [SD]: 14.1) and a predominance of the female sex (63%). Regurgitation and heartburn were the most frequent symptoms (66%), followed by cough and belching, with a frequency of 31% and 11%, respectively. The indication for performing pH-impedance testing in order of frequency was GERD, presurgery, typical symptoms, atypical symptoms, and patients with chest pain and dysphagia (Table 1). Of the patients included in the study, 34.7% did not receive any PPI, and 2% received alginate as a treatment strategy for GERD. The most frequently used PPI was esomeprazole, with 26%, followed by lansoprazole, dexlansoprazole, and pantoprazole (Table 1).

The comparison of manometric variables between patients with and without acid reflux and those with and without non-acid reflux is presented in Table 2; the mean LES pressure, the end-inspiratory LES pressure, the median IRP (integrated relaxation pressure), DCI, and EGJ-CI had no statistically significant differences between the groups.

The type of EGJ was different between patients with a positive versus negative pH impedance study for non-acid reflux ($p = 0.04$), with a more considerable proportion of patients with Type IIIa or IIIb EGJ among patients without that type of reflux (63.1% versus 31.3%). Among patients with acid reflux, no statistically significant differences were found in the variables studied (Table 2).

Lower end-inspiratory LES pressure was detected in patients with erosive reflux than those with non-erosive reflux (median: 6.1 vs. 11.9; $p < 0.01$). There was a similar when evaluating the EGJ-CI (median: 15.1 vs. 23, $p = 0.04$). The proportion of patients with Type IIIa and IIIb EGJ was higher in patients with erosive reflux (83.3% vs. 37.1%; $p < 0.01$) (Table 3).

DISCUSSION

The present study described the function of the EGJ in the different reflux phenotypes (acid, non-acid, erosive, and non-erosive) and found that Type IIIa and IIIb EGJs were more frequent in patients with erosive esophagitis and less common among patients with non-acid reflux. Additionally, EGJ-CI is significantly lower in patients with erosive reflux.

In our study, the symptoms that most frequently occurred in the population with GERD were regurgitation and heartburn, and the highest proportion of patients were women. These results correlate with what was documented by Paramo et al. in a published study of the prevalence of

Table 1. Clinical characteristics and treatment received by the included patients

Variable	Value
	n = 72
Female sex, n (%)	46 (63.9)
Age, mean (SD)	54.9 (14.1)
Symptoms	
- Regurgitation, n (%)	48 (66.7)
- Heartburn, n (%)	48 (66.7)
- Belching, n (%)	8 (11.1)
- Cough, n (%)	23 (31.9)
- Positive SI, n (%)	39 (54.2)
- SAP, n (%)	25 (34.7)
PPI use	
- Does not receive (%)	25 (34.7)
- One dose, n (%)	26 (36.1)
- Double dose, n (%)	21 (29.1)
pH impedance indication	
- GERD, n (%)	47 (65.28)
- Pre-surgical, n (%)	7 (9.7)
- Regurgitation, n (%)	4 (5.5)
- Heartburn, n (%)	7 (9.7)
- Cough, n (%)	3 (4.1)
- Belching, n (%)	2 (2.78)
- Chest pain, n (%)	1 (1.3)
- Dysphagia, n (%)	1 (1.3)
Proton pump inhibitor type	
- Esomeprazole, n (%)	19 (26.3)
- Omeprazole, n (%)	5 (6.94)
- Pantoprazole, n (%)	6 (8.33)
- Lansoprazole, n (%)	8 (11.1)
- Dexlansoprazole, n (%)	7 (9.7)
- No use of PPI, n (%)	25 (34.7)
- Alginate, n (%)	2 (2.78)

SD: standard deviation; n = number. Prepared by the authors.

symptoms of gastroesophageal reflux and associated factors in the Colombian population⁽¹¹⁾.

The PPI most frequently prescribed in our population was esomeprazole, followed by lansoprazole and dexlansoprazole, molecules with proven effectiveness in managing GERD according to different published studies⁽¹²⁾.

When the manometric variables were evaluated in patients with and without acid reflux, we did not find significant differences in EGJ-CI, contrary to what was reported by Gor et al.⁽¹³⁾, who found an inverse correlation between EGJ-CI and gastroesophageal reflux determined by AET. No statistically significant differences were found in the mentioned manometric parameters when the analysis was performed for the non-acid reflux subgroup. This finding could be because all of the patients in that study did not receive antisecretory therapy, while our study included patients with and without antisecretory treatment.

When evaluating the manometric variables between erosive and non-erosive reflux, our results were similar to those by Hyoju Ham et al., who reported that EGJ-CI is significantly correlated with the morphology of the EGJ, establishing the presence of GERD when the morphology of the EGJ is altered. For example, a Type 3 EGJ was associated with a more significant number of reflux episodes, a finding also correlated with EGJ-CI alteration⁽¹⁴⁾. Furthermore, our study demonstrated that the EGJ-CI is significantly decreased in patients with erosive GERD compared to non-erosive GERD ($p = 0.04$), representing a greater impairment of the barrier mechanism in the EGJ. Finally, we documented a significant decrease in end-inspiratory LES pressure among patients with erosive GERD compared to patients with non-erosive GERD ($p < 0.01$). The above shows us that the EGJ-CI is a new manometric parameter altered in patients with erosive and non-erosive GERD, significantly correlated with the alteration of end-inspiratory LES pressure and Type 3 EGJ. These results suggest that alteration of EGJ-CI could predict erosive reflux in our population and allow us to understand how the impairment of the anti-reflux barrier in patients with erosive GERD is more significant than in those with non-erosive GERD.

Wang et al. established a normal cut-off point for EGJ-CI with a mean of 34.7 mm Hg and a range between 26.2 and 58.3 mm Hg⁽¹⁵⁾; these results are compatible with ours, given that in our population with erosive GERD, the EGJ-CI had a median of 15.1 with a range between 7 and 25 mm Hg, decreased compared to this cut-off point established for healthy patients. In patients with non-erosive GERD, a decrease in EGJ-CI was also documented concerning these cut-off points, suggesting a dysfunction of the EGJ barrier mechanism in these patients. A recent

Table 2. Comparison of manometric variables in patients with or without acid and non-acid reflux

Manometric variable	Acid reflux			Non-acid reflux		
	Positive, n = 59	Negative, n = 13	p-value	Positive, n = 34	Negative, n = 38	p-value
Mean LES pressure (mm Hg), median (IQR)	10,4 (3,6 -21,9)	13,1 (11,6 -23,8)	0,08	12,1 (8,5-23,8)	10,1 (3,6 -15)	0,11
End-inspiratory LES pressure (mm Hg), median (IQR)	9,6 (5,9-15,4)	9,4 (5,9-12,9)	0,75	8,4 (5,2 -13,3)	9,85 (6,2 -15,4)	0,53
Median IRP (IQR)	3,9 (2,0-6,7)	4,4 (1,4-6,2)	0,80	4,3 (1,4-6,2)	3,9 (2,1-7,3)	0,48
DCI mm Hg, median (IQR)	1551 (910 -2412)	1476 (774-1862)	0,53	1660 (1021-2134)	1389 (783-2320)	0,52
EGJ-CI, median (IQR)	21 (14-44)	15 (11-28)	0,24	20,5 (12,7-30,0)	20 (14-44)	0,80
EGJ type, n (%)						
- I	8 (13,50)	2 (15, 38)	0,82	6 (17,65)	4 (10,53)	0,04
- II	21 (35,59)	6 (46, 15)		17(50,00)	10 (26,32)	
- IIIa	16 (27,12)	2 (15,38)		4 (11,7)	14 (36,84)	
- IIIb	14 (23,73)	3 (23,08)		7 (20,59)	10 (26,32)	

DCI: distal contractile integral; IQR: interquartile range; IRP: integrated relaxation pressure; LES: lower esophageal sphincter. Prepared by the authors.

Table 3. Comparison of manometric variables between erosive reflux and non-erosive reflux

Manometric variable	Erosive reflux		
	Positive, n = 18	Negative, n = 54	p-value
Mean LES pressure (mm Hg), median (IQR)	9,9 (4,0-11,6)	11,5 (5,0 -23,8)	0,09
End-inspiratory LES pressure (mm Hg), median (IQR)	6,1 (2,8-7,5)	11,9 (6,6-17,4)	< 0,01
Median IRP (IQR)	3,75 (2,3-6,6)	4,3 (2,8-6,3)	0,94
DCI (mm Hg), median (IQR)	1549 (783-2007)	1546 (991-2241)	0,41
EGJ-CI, median (IQR)	15,1 (7-25)	23 (14-42,0)	0,04
EGJ type, n (%)			
- I	2,0 (11,1)	8,0 (14,8)	< 0,01
- II	1,0 (5,6)	26,0 (48,2)	
- IIIa	9,0 (50,0)	9,0 (16,7)	
- IIIb	6,0 (33,3)	11,0 (20,4)	

DCI: distal contractility integral; IQR: interquartile range; IRP: integrated relaxation pressure; LES: lower esophageal sphincter. Prepared by the authors.

study published by Rogers et al., who evaluated the EGJ in healthy patients using the two manometric parameters described (EGJ type and EGJ-CI), established normal cut-off points for the EGJ-CI, very similar to those provided by Wang et al.⁽¹⁶⁾ and Jasper et al.⁽¹⁷⁾.

Tolone et al. explained that 50% of patients with decreased EGJ-CI present with GERD, and 14% have functional heartburn; data suggest that decreased EGJ-CI is associated with a longer AET, greater number of reflux episodes and esophageal mucosal damage documented on endos-

copy⁽¹⁸⁾. These findings were duly noted in our population for patients with erosive and non-erosive GERD.

We can conclude that in the Colombian population, the EGJ-CI objectively evaluates the barrier mechanism of the EGJ since it is decreased in all patients with GERD. Furthermore, as shown in our study, it allowed a characterization of patients with erosive esophagitis, given that the EGJ-CI in this group was found to be significantly decreased, which was correlated with the decrease in end-inspiratory LES pressure and Type IIIa and IIIb EGJ.

CONCLUSION

In our population, decreased EGJ-CI was significantly correlated with erosive GERD, suggesting that a reduction in EGJ-CI values could predict this condition in patients with GERD. This result is appropriately related to the type of EGJ morphology, with Type III being the most frequent, and lower end-inspiratory LES pressure, allowing us to characterize this reflux phenotype better to guarantee an appropriate diagnostic and therapeutic approach.

REFERENCES

1. Vakil N, van Zanten SV, Kahrilas P, Dent J, Jones R, Bianchi LK, et al. The Montreal definition and classification of gastroesophageal reflux disease: A global evidence-based consensus. *Am J Gastroenterol*. 2006;101(8):1900-20; quiz 1943. <https://doi.org/10.1111/j.1572-0241.2006.00630.x>
2. Prakash Gyawali C, Kahrilas PJ, Savarino E, Zerbib F, Mion F, Smout AJPM, et al. Modern diagnosis of GERD: The Lyon Consensus. *Gut*. 2018;67(7):1351-1362. <https://doi.org/10.1136/gutjnl-2017-314722>
3. Prakash Gyawali C, Kahrilas PJ, Savarino E, Zerbib F, Mion F, Smout AJPM, et al. Modern diagnosis of GERD: The Lyon Consensus. *Gut*. 2018;67(7):1351-1362. <https://doi.org/10.1136/gutjnl-2017-314722>
4. Hani A, Bernal W, Leguizamó AM, Zuluaga C, Vargas R, Vergara H, et al. Cómo realizar e interpretar una manometría esofágica de alta resolución usando la clasificación de Chicago 3.0. *Rev Col Gastroenterol*. 2017;32(4):369-78. <https://doi.org/10.22516/25007440.181>
5. Kahrilas PJ, Mittal RK, Bor S, Kohn GP, Lenglinger J, Mittal SK, et al. Chicago Classification update (v4.0): Technical review of high-resolution manometry metrics for EGJ barrier function. *Neurogastroenterol Motil*. 2021;33(10):e14113. <https://doi.org/10.1111/nmo.14113>
6. Nicodème F, Pipa-Muniz M, Khanna K, Kahrilas PJ, Pandolfino JE. Quantifying esophagogastric junction contractility with a novel HRM topographic metric, the EGJ-Contractile Integral: Normative values and preliminary evaluation in PPI non-responders. *Neurogastroenterol Motil*. 2014;26(3):353-60. <https://doi.org/10.1111/nmo.12267>
7. Katzka DA, Pandolfino JE, Kahrilas PJ. Phenotypes of gastroesophageal reflux disease: Where Rome, Lyon, and Montreal meet. *Clin Gastroenterol Hepatol*. 2020;18(4):767-76. <https://doi.org/10.1016/j.cgh.2019.07.015>
8. Costa VA, Pinto Saavedra OM, Hani Amador AC, Leguizamó Naranjo AM, Ardila Hani AF. Actualización en la interpretación de la medición del pH e impedanciometría. *Rev Col Gastroenterol*. 2021;36(1):73-80. <https://doi.org/10.22516/25007440.608>
9. Carlson DA, Gyawali CP, Khan A, Yadlapati R, Chen J, Chokshi RV, et al. Classifying esophageal motility by FLIP panometry: A study of 722 subjects with manometry. *Am J Gastroenterol*. 2021;116(12):2357-2366. <https://doi.org/10.14309/ajg.0000000000001532>
10. Yadlapati R, Kahrilas PJ, Fox MR, Bredenoord AJ, Prakash Gyawali C, Roman S, et al. Esophageal motility disorders on high-resolution manometry: Chicago classification version 4.0 ©. *Neurogastroenterol Motil*. 2021;33(1):e14058. <https://doi.org/10.1111/nmo.14058>
11. Páramo-Hernández DB, Albis R, Galiano MT, de Molano B, Rincón R, Pineda-Ovalle LF, et al. Prevalencia de síntomas del reflujo gastroesofágico y factores asociados: una encuesta poblacional en las principales ciudades de Colombia. *Rev Col Gastroenterol*. 2016;31(4):337-46. <https://doi.org/10.22516/25007440.108>
12. Huerta-Iga F, Bielsa-Fernández MV, Remes-Troche JM, Valdovinos-Díaz MA, Tamayo-de la Cuesta JL, en representación del Grupo para el estudio de la ERGE 2015. Diagnóstico y tratamiento de la enfermedad por reflujo gastroesofágico: recomendaciones de la Asociación Mexicana de Gastroenterología. *Rev Gastroenterol Mex*. 2016;81(4):208-22. <https://doi.org/10.1016/j.rgm.2016.04.003>
13. Gor P, Li Y, Munigala S, Patel A, Bolckhir A, Gyawali CP. Interrogation of esophagogastric junction barrier function using the esophagogastric junction contractile integral: an observational cohort study: EGJ barrier function on HRM. *Dis Esophagus*. 2016;29(7):820-8. <https://doi.org/10.1111/dote.12389>
14. Ham H, Cho YK, Lee HH, Yoon SB, Lim CH, Kim JS, et al. Esophagogastric junction contractile integral and morphology: Two high-resolution manometry metrics of the anti-reflux barrier. *J Gastroenterol Hepatol (Australia)*. 2017;32(8):1443-9. <https://doi.org/10.1111/jgh.13720>
15. Wang D, Patel A, Mello M, Shriver A, Gyawali CP. Esophagogastric junction contractile integral (EGJ-CI)

- quantifies changes in EGJ barrier function with surgical intervention. *Neurogastroenterology and Motility*. 2016;28(5):639-46.
<https://doi.org/10.1111/nmo.12757>
16. Rogers BD, Rengarajan A, Abrahao L, Bhatia S, Bor S, Carlson DA, et al. Esophagogastric junction morphology and contractile integral on high-resolution manometry in asymptomatic healthy volunteers: An international multicenter study. *Neurogastroenterol Motil*. 2021;33(6):e14009.
<https://doi.org/10.1111/nmo.14009>
17. Jasper D, Freitas-Queiroz N, Hollenstein M, Misselwitz B, Layer P, Navarro-Rodriguez T, et al. Prolonged measurement improves the assessment of the barrier function of the esophago-gastric junction by high-resolution manometry. *Neurogastroenterol Motil*. 2017;29(2).
<https://doi.org/10.1111/nmo.12925>
18. Tolone S, de Bortoli N, Marabotto E, de Cassan C, Bodini G, Roman S, et al. Esophagogastric junction contractility for clinical assessment in patients with GERD: A real added value? *Neurogastroenterol Motil*. 2015;27(10):1423-31.
<https://doi.org/10.1111/nmo.12638>