



Original investigation

Coronal knee moments of asymptomatic elderly individuals during gait: Preliminary study[☆]



José David Vélez Uribe*, Ailin Caterin Zuluaga Zuluaga, Ana Cristina Arias Tobón, Nicolás Eugenio Gómez Suárez

Facultad de Fisioterapia, Universidad CES, Medellín, Colombia

ARTICLE INFO

Article history:

Received 25 May 2020

Accepted 21 July 2020

Available online 28 October 2021

Keywords:

Aged

Gait

Kinetics

Knee

Osteoarthritis

ABSTRACT

Introduction: Joint moments are explained by the forces that occur during gait and generate a movement in the joint. The maximum peak adduction moment (MPM) of the knee has been associated with osteoarthritis (OA). In Colombia, however, there are no references to these values for clinical use in an asymptomatic population.

Objective: To describe the characteristics of joint moments in the coronal plane of the knee of asymptomatic elderly whilst walking.

Materials and methods: This is a descriptive study that included 50 participants of both genders and older than 60 years old. They were residents of the metropolitan area del Valle de Aburrá that had volunteered and were capable of following indications without any external aids while walking, and with no pain. Gait analysis was performed when walking at a normal speed on a 6 m pathway on two force plates, and was registered by a system of 8 opto-electric 3D recording cameras. The statistical analysis was performed using SPSS 20.

Results: A mean BMI of 26.34 kg/m² was found, and the mean of the MPM was 0.26 Nm/kg, generating a graph with two peaks, with the first being of greater magnitude that occurred in 17% of the gait cycle.

Conclusions: The MPM of 0.26 Nm/kg found in this population is lower than that reported in the osteoarthritis population. A reference curve for gait analysis has been established in these asymptomatic elderly. It is essential that these values are available, in order to help the professional to make precise and quantitative diagnoses in gait alterations.

© 2021 Published by Elsevier España, S.L.U. on behalf of Asociación Colombiana de Reumatología.

[☆] Please cite this article as: Vélez Uribe JD, Zuluaga Zuluaga AC, Arias Tobón AC, Gómez Suárez NE. Momentos articulares de la rodilla en el plano coronal durante la marcha de adultos mayores asintomáticos: estudio preliminar. Rev Colomb Reumatol. 2021;28:276–281.

* Corresponding author.

E-mail address: jvelezu@ces.edu.co (J.D. Vélez Uribe).

Momentos articulares de la rodilla en el plano coronal durante la marcha de adultos mayores asintomáticos: estudio preliminar

R E S U M E N

Palabras clave:

Anciano
Marcha
Cinética
Rodilla
Osteoartritis

Introducción: Los momentos articulares se explican por las fuerzas que se producen durante la marcha, las cuales producen un giro en la articulación. El pico de momento máximo aductor (PMM) de la rodilla se ha asociado con osteoartritis (OA), sin embargo, en Colombia no se tienen valores de referencia en población asintomática para su uso clínico.

Objetivo: Describir las características del momento articular de la rodilla en el plano coronal durante la marcha de adultos mayores (AM) asintomáticos.

Materiales y métodos: Estudio descriptivo en el que se evaluaron 50 participantes mayores de 60 años, residentes en el área metropolitana del Valle de Aburrá, de ambos géneros, que participaron voluntariamente, capaces de seguir indicaciones, sin ayudas externas durante la marcha ni dolor. Se realizó el análisis de la marcha caminando a velocidad habitual, por una pista de 6 metros sobre dos plataformas de fuerza, y se registró con un sistema de 8 cámaras optoeléctricas 3D. El análisis estadístico se hizo con SPSS 20.

Resultados: Se encontró un IMC de 26,34 kg/m² y un PMM de 0,26 Nm/kg que generó una gráfica con 2 picos, el primero de estos de mayor magnitud, en el 17% del ciclo de la marcha.

Conclusión: El PMM de 0,26 Nm/kg encontrado en esta población es menor a lo reportado en personas con OA. Se establece la curva de referencia en esta población asintomática para el análisis de marcha. Es indispensable que existan dichos valores, puesto que ayudan al profesional a realizar diagnósticos oportunos, precisos y cuantitativos en alteraciones de la marcha.

© 2021 Publicado por Elsevier España, S.L.U. en nombre de Asociación Colombiana de Reumatología.

Introduction

Gait is defined as the bipedal motion of humans, with low effort and minimal energy consumption. In the elderly gait is associated with independence and interaction with the environment, as well as the potential to conduct the activities of daily living (ADL). When free ambulation declines, these activities become limited.^{1,2} Functional alterations of the joints in the lower extremities may affect the ability to walk, particularly when the dynamic and static stability of the knee joint is interrupted due to structural changes such as osteoarthritis (OA), a degenerative disease associated with aging, and affecting the functionality of the affected individuals.³⁻⁵

Knee OA is a musculoskeletal disease which could be idiopathic but is associated with the interaction of biological factors such as genetic predisposition, gender, age and race, or factors of the joint mechanics, of which the most relevant are overweight, previous injuries, the level of physical activity and the adduction and abduction moments of the joint.^{1,6-8} Hence, knee OA may affect between 7 and 17% of the population, with a trend to increase with aging; this factor increases the cost for healthcare systems.^{1,9}

Individuals with OA of the medial or lateral compartment exhibit more varus or valgus alignment, respectively, as compared to controls.¹⁰⁻¹³ Likewise, varus alignment is greater in the most severe cases of OA.¹⁴⁻¹⁶ During the stance phase of gait in people with this condition and varus deformity, there is an increased external moment of knee adduction, which increases the medial compartment forces which is more prone to develop OA.^{4,17}

The forces that cross the knee during the stance phase of gait may be internal or external. The former are produced by muscles, ligaments, and soft tissues, while the latter are produced by the ground reaction and gravity forces. The external forces are applied medially with respect to the center of articulation of the knee and tend to cause a medial rotation of the tibia with respect to the femur in the coronal plane.¹⁷ This phenomenon is called the external adduction moment – of relevance for the elderly population – since the higher it is, the more the compressive forces increase in the medial compartment of the knee, and this is associated with the development of degenerative disease.^{9,17-19} The literature describes that this moment presents a peak at the 20% of the gait stance phase,⁹ called maximum peak moment (MPM), measured in Nm/kg. However, to accurately access such data, the *gold standard* should be motion analysis laboratories^{20,21} that have their own reference parameters.²²

Hence, in order to establish the MPM deviations in the Colombian population, international references are used but this may bias the interpretation of the data.²² It is important for each laboratory to have its own reference parameters, since the anthropometric and demographic characteristics, as well as the individual life styles, change according to their nationality. Moreover, the elderly population is growing leading to an increasing demand for healthcare, particularly from practitioners working in the area of human body motion. This highlights the need to procure reference parameters for intervention of the Colombian elderly population.²³ Until now, no studies had been conducted in Colombia – specifically at the motion analysis laboratory of CES University – on the

knee joint moments in the coronal plain during walking of asymptomatic elderly individuals. Consequently, this research project is intended to describe the knee joint movements in the coronal plane, during gait in asymptomatic elderly individuals, residents of the Valle de Aburrá, Antioquia, Colombia.

Materials and methods

Study type and design

A descriptive, cross-sectional study was conducted intended to describe the knee joint moments in the coronal plane during gait of asymptomatic elderly individuals, residents of the Valle de Aburrá.

Population and sample

The reference population for this study was Colombian elderly individuals, aged 60 years and older, residents of the Valle de Aburrá. The inclusion criteria were: both males and females who accepted to voluntarily participate, able to follow instructions and to walk with no external walking aids. The exclusion criteria were: fractures or surgeries of the lower extremities during the past two years; neurological disease, cardiopulmonary instability or experiencing pain at the time of the assessment; uncorrected visual impairment; hearing or sensory limitations; allergies to the instruments used for assessment, such as allergy to double-sided tape. The data from 50 elderly individuals were analyzed, who were selected in a non-probabilistic and convenience mode.

Data collection

Following the signature of the informed consent, each elderly individual was asked to dress in comfortable clothing provided by the team of investigators, suitable for data capture. The measurements were taken using an anthropometer (Lafayette 60) and ergonomic measuring tape (SECA model 201), the del Full Body PiG model, measuring each item on both sides of the body. The measurements were: length of lower limbs, ankle width, knee width, elbow width, wrist with, hand width and shoulder displacement, height and body mass. Subsequently, the 39 markers were entered, including 4 head markers and markers at C7, T10, posterosuperior and anterosuperior iliac spines, acromion, arm, lateral humeral epicondyle, forearm, styloid process of the ulna and radius, head of the second metacarpal, manubrium of the sternum, xyphoid process, thigh, lateral femoral condyle, leg, external malleolus, heel bone, and head of the second metatarsal.²⁴

The participant was asked to walk back and forth over a 6-meter walkway and complete four runs. Two force platforms (AMTI-OR6-7-1000) were placed in the middle and 8 previously calibrated optoelectrical cameras surrounding the area – Bonita 10 (VICON, Oxford, UK). The participant was instructed to “walk as you normally do, and at your normal speed”. No records were made while the participant adapted to the laboratory environment and the markers; then at least 7 valid captures were recorded. This was done when the participant made full contact with a different foot on each of the

platforms. Once the data were captured, the information was processed and the appropriateness of the tests was checked using the Polygon 4.3.3 *software*. Then the data were exported onto an Excel spreadsheet to create the database for analysis using the SPSS V.20 *software*.

Statistical analysis

In order to describe the sociodemographic characteristics and assess the knee joint moments in the coronal plane, the qualitative variables were presented using frequencies and percentages; the quantitative variables were expressed as means and standard deviation or medians and interquartile ranges, according to distribution. Pearson or Spearman were used to assess the relationship of the MPM to the anthropometric variables (size, mass and BMI), spatial-temporal (gait speed, length of step, length of stride, step rate, stance phase) based on distribution.

Ethical considerations

The study was conducted under the principles of the Declaration of Helsinki, the ethical standards for research in humans of the World Health Organization (WHO), Pan American Health Organization (PAHO), the Council of International Organizations of Medical Sciences (CIOMS) and Resolution 8430 of 1993 of the Colombian Ministry of Health, keeping in mind good research practices favoring the participants and respecting intellectual property during the development of the project. The participation was voluntary and those who accepted to be part of the study, previously signed the informed consent.

Results

Sample data

50 asymptomatic elderly individuals with a mean age of 66 years (64–72) were studied; 52% were females ($n=26$). Only the anthropometric variables showed a normal distribution: height 1.60 (± 0.087) m, mass 66.07 (± 13.54) kg, BMI 26.34 kg/m². The median MPM of the knee in the coronal plane was 0.26 (± 0.67) Nm/kg.

Fig. 1 illustrates the behavior of the knee joint moment in the coronal plane during the gait cycle: the vertical shows the positive and negative adduction moments, whilst the horizontal axis shows the percentage of the gait cycle where the stance phase had a duration of 60.42%. The moments were mostly adduction and exhibited a curve with two peaks. The MPM occurred at approximately 17% of the gait cycle, and the second lower intensity peak at 40–60%.

Table 1 shows the references of the time, spatial and spatial-temporal values which presented a non-normal behavior; each variable was reported in terms of the median and its corresponding IQR.

Table 2 shows the correlations between MPM and the anthropometric and spatial-temporal aspects; no statistically significant values or strong correlations were identified among the variables assessed.

Knee ADD/ABD moment




Fig. 1 – Coronal knee joint moments – average reported as a central line and standard deviation.

Table 1 – Univariate analysis of the spatial-temporal variables.

Spatial-temporal variables	Median	IQR
Length of step (m)	0.55	(0.53–0.59)
Length of stride (m)	1.12	(1.06–1.19)
Step cadence (steps/min)	108.79	(100.43–114.42)
Gait speed (m/s)	1.01	(0.93–1.07)
Stance phase (%)	60.42	(59.20–62.05)

IQR: interquartile range.

Table 2 – Correlation between MPM and anthropometric/spatial-temporal variables.

Variables	Stance phase maximum peak		
	Rho	p	95% CI
Size	-0.13*	0.17	(158.7–1618.9)
Mass	-0.18*	0.07	(63.06–68.43)
BMI	-0.17*	0.87	(25.31–27.11)
Length of step	0.09**	0.36	(0.54–0.59)
Length of stride	0.07**	0.48	(1.08–1.13)
Stance phase	0.06**	0.55	(58.57–61.18)
Cadence of step	0.15**	0.11	(107.39–127.74)
Gait speed	0.16**	0.10	(0.98–1.04)

CI: confidence interval; BMI: body mass index.

* Pearson correlation test.

** Spearman correlation test.

Discussion

This preliminary study shows the value of the knee joint moments in the coronal plane, identifying the highest peak in the stance phase; however, no associations were found between each variable associated with gait of the

asymptomatic elderly individuals studied. Until this date, there are no data available in Colombia reporting the values for the coronal knee joint moments during gait in this population. The data herein collected may be interpreted as an initial description which could be the foundation for future studies on knee joint moments and how these may affect the occurrence of musculoskeletal diseases.

The graphical representation of the external knee joint moment in the subjects studied showed a curve with two adduction peaks in the stance phase. The MPM was identified at 17% of the gait cycle, indicating a similar pattern to what Foroughi et al.²⁵ described in their systematic review about coronal knee joint moments, in patients with OA. Despite the consistency of the general morphology of the curves, the MOM in this study was 0.26 Nm/kg in asymptomatic subjects, while in the systematic review the MOM was reported between 0.25 and 0.71 Nm/kg in patients with OA, and a narrower range, but with lower peaks, between 0.33 and 0.55 Nm/kg for controls. According to these data, MPM could be related to the severity and the OA symptoms.²⁵ Moreover, this evidences a significant variation among the various nationalities, which further emphasizes the need to establish population-specific parameters according to symptomatology.

The adduction MPM herein described refers to the maximum force per kilogram that results in the tendency of the tibia to rotate medially with respect to the femur, which actually happens during the stance phase of gait. This rotational force may account for the maximum compression on the medial compartment of the knee. Hence, several authors argue that elevated MPMs are associated with radiographic changes and degeneration of the knee cartilage,^{26,27} due to increased medial contact force which is presumed to be the deleterious mechanical effect for the cartilage.²⁸ Moreover, the adduction MPM, the peak of the maximum moment in the sagittal

plane (flexor moment peak) is also considered a significant contributor to the force of the medial contact of the knee,²⁹ hence showing the importance of the three-dimensional components in the analysis of motion and the need to study these components in populations at increased risk for OA.

Furthermore, the body mass index (BMI) of the participants indicates overweight (BMI = 26.34 kg/m²), a factor which depending on the magnitude, may increase the risk for OA up to four-fold, in contrast to normal weight individuals.³⁰ However, this study failed to show an association with MPM, probably because the participants were asymptomatic. Similarly, Verlaan et al.³⁰ studied the relationship of the adduction moment while going up and down the stairs in patients with knee OA, with and without obesity, and they found no differences in the adduction moment among these participants. The authors believe that their findings are due to the fact that obese individuals slow down their walking speed as an effective strategy to reduce the adduction moment during stair negotiation.

Over time, humans have experienced aged-related musculoskeletal changes that may lead to gait impairment, with a subsequent reduction in gait cadence, length of stride, and speed by 20%.³¹ In this particular study the speed was 1.01 m/s, and although participants were allowed to walk at their preferred speed and were given some time to adapt, the speed was below the reference values according to several authors, with asymptomatic populations of various age groups^{22,32,33} and of elderly individuals, such as the Mündermann et al. trial.¹⁵ The authors observed a group of subjects with OA and a control group; their speeds were 1.25 and 1.23 m/s, respectively. It may be argued that the differences among these studies are due to the assessment environment in the laboratory and the length of the walkway. However, the speed apparently did not affect the MPM, but considering that in this study the MPM and the gait speed were lower than the numbers reported by other authors, it is possible to conclude that reducing the speed increases the duration of the stance phase, which may contribute to reducing the ground reaction force³⁰ and hence the moment.

The limitations of this study may be related to the number of participants; nevertheless, the kinetics of both knees was analyzed and totaled 100 data points. Moreover, it may be valuable to include the analyses of other complementary tests in future studies, to establish the correlation between any existing symptoms and structural changes, including the static load alignment test, as well as the behavior of joint kinematics of the lower extremity, since as already mentioned, it is associated with several factors affecting the progression of OA and the joint moment. Another limitation was that the assessment model failed to consider the muscle activation factor involved in generating the moments in the coronal plane. This value is important to determine the involvement of muscles and other soft tissues for potential therapies to strengthen these structures and their impact on the loads that are injurious to the knee. It should be noted however, that the available bibliography is not very extensive for the analysis of the population studied, since most published articles discuss the behavior of joint moments on each plane in individuals with musculoskeletal pathologies, but not in asymptomatic populations.

Conclusion

This study analyzed the coronal knee joint moments during the stance phase of gait, in 50 asymptomatic elderly subjects, using a 3D analytical system. The behavior curve of the external adduction moment is similar to the findings reported by researchers in other countries, but with an adduction MPM of 0.26 Nm/kg, which in general is less than the figures reported in the population with OA. However, this MPM was not associated with other anthropometric or spatial-temporal variables.

Further studies are needed to expand the reference parameters used by motion analysis laboratories and healthcare professionals, to contribute to making accurate, timely, quantitative and objective diagnoses in gait disorders and to make headway in physiotherapy intervention. Although these technological tools are not always available, this study provides an initial reference on behavior of the knee joint moment in asymptomatic elderly subjects.

Funding

This article corresponds to the study "Reference Parameters of gait in elderly subjects, residents of the Valle de Aburra", financed by the Directorate of Research and Innovation of Universidad CES, through the internal minimum amount call for proposals, code: INV.022018.003.

Conflicts of interest

The authors have no conflicts of interest to disclose.

REFERENCES

1. Safiri S, Kolahi A-A, Smith E, Hill C, Bettampadi D, Mansournia MA, et al. Global, regional and national burden of osteoarthritis 1990–2017: a systematic analysis of the Global Burden of Disease Study 2017. *Ann Rheum Dis*. 2020;79:819–28, <http://dx.doi.org/10.1136/annrheumdis-2019-216515>.
2. Magnani PE, Freire Junior RC, Grecco Zanellato NF, Baena Genovez M, Camargo Alvarenga I, Carvalho de Abreu DC. The influence of aging on the spatial and temporal variables of gait during usual and fast speeds in older adults aged 60 to 102 years. *Hum Mov Sci*. 2019;68:102540, <http://dx.doi.org/10.1016/j.humov.2019.102540>.
3. Park S-K, Kobsar D, Ferber R. Relationship between lower limb muscle strength, self-reported pain and function, and frontal plane gait kinematics in knee osteoarthritis. *Clin Biomech (Bristol, Avon)*. 2016;38:68–74, <http://dx.doi.org/10.1016/j.clinbiomech.2016.08.009>.
4. Fukaya T, Mutsuzaki H, Okubo T, Mori K, Wadano Y. Relationships between the center of pressure and the movements of the ankle and knee joints during the stance phase in patients with severe medial knee osteoarthritis. *Knee*. 2016;23:631–6, <http://dx.doi.org/10.1016/j.knee.2016.02.019>.
5. Martínez Figueroa R, Martínez Figueroa C, Calvo Rodríguez R, Figueroa Poblete D. Osteoarthritis (artrosis) de rodilla. *Rev Chil Ortop Traumatol*. 2015;56:45–51, <http://dx.doi.org/10.1016/j.rchot.2015.10.005>.

6. Pascual Huerta J [Thesis] Determinantes mecánicos de los momentos articulares de la rodilla en el plano frontal durante la fase de apoyo de la marcha. Universidad Complutense de Madrid; 2010. Available from: <https://eprints.ucm.es/11613/>
7. Doherty M. Risk factors for progression of knee osteoarthritis. *Lancet*. 2001;358:775-6, [http://dx.doi.org/10.1016/S0140-6736\(01\)06006-8](http://dx.doi.org/10.1016/S0140-6736(01)06006-8).
8. Dieppe PA, Lohmander LS. Pathogenesis and management of pain in osteoarthritis. *Lancet*. 2005;365:965-73, [http://dx.doi.org/10.1016/S0140-6736\(05\)71086-2](http://dx.doi.org/10.1016/S0140-6736(05)71086-2).
9. Zhang L, Liu G, Han B, Wang Z, Yan Y, Ma J, et al. Knee Joint biomechanics in physiological conditions and how pathologies can affect it: a systematic review. *Appl Bionics Biomech*. 2020;2020:7451683, <http://dx.doi.org/10.1155/2020/7451683>.
10. Janakiraman N, Teichtahl AJ, Wluka AE, Ding C, Jones G, Davis SR, et al. Static knee alignment is associated with the risk of unicompartmental knee cartilage defects. *J Orthop Res*. 2008;26:225-30, <http://dx.doi.org/10.1002/jor.20465>.
11. Briem K, Snyder-Mackler L. Proximal gait adaptations in medial knee OA. *J Orthop Res*. 2009;27:78-83, <http://dx.doi.org/10.1002/jor.20718>.
12. Teichtahl AJ, Cicuttini FM, Janakiraman N, Davis SR, Wluka AE. Static knee alignment and its association with radiographic knee osteoarthritis. *Osteoarthritis Cartilage*. 2006;14:958-62, <http://dx.doi.org/10.1016/j.joca.2007.02.017>.
13. Whelton C, Thomas A, Elson DW, Metcalfe A, Forrest S, Wilson C, et al. Combined effect of toe out gait and high tibial osteotomy on knee adduction moment in patients with varus knee deformity. *Clin Biomech (Bristol, Avon)*. 2017;43:109-14, <http://dx.doi.org/10.1016/j.clinbiomech.2017.02.009>.
14. Mündermann A, Dyrby CO, Hurwitz DE, Sharma L, Andriacchi TP. Potential strategies to reduce medial compartment loading in patients with knee osteoarthritis of varying severity: reduced walking speed. *Arthritis Rheum*. 2004;50:1172-8, <http://dx.doi.org/10.1002/art.20132>.
15. Mündermann A, Dyrby CO, Andriacchi TP. Secondary gait changes in patients with medial compartment knee osteoarthritis: increased load at the ankle, knee, and hip during walking. *Arthritis Rheum*. 2005;52:2835-44, <http://dx.doi.org/10.1002/art.21262>.
16. Thorp LE, Wimmer MA, Block JA, Moio KC, Shott S, Goker B, et al. Bone mineral density in the proximal tibia varies as a function of static alignment and knee adduction angular momentum in individuals with medial knee osteoarthritis. *Bone*. 2006;39:1116-22, <http://dx.doi.org/10.1016/j.bone.2006.05.001>.
17. Reeves ND, Bowling FL. Conservative biomechanical strategies for knee osteoarthritis. *Nat Rev Rheumatol*. 2011;7:113-22, <http://dx.doi.org/10.1038/nrrheum.2010.212>.
18. Nie Y, Wang H, Xu B, Zhou Z, Shen B, Pei F. The relationship between knee adduction moment and knee osteoarthritis symptoms according to static alignment and pelvic drop. *BioMed Res Int*. 2019;2019:7603249, <http://dx.doi.org/10.1155/2019/7603249>.
19. Hall M, Bennell KL, Wrigley TV, Metcalf BR, Campbell PK, Kasza J, et al. The knee adduction moment and knee osteoarthritis symptoms: relationships according to radiographic disease severity. *Osteoarthritis Cartilage*. 2017;25:34-41, <http://dx.doi.org/10.1016/j.joca.2016.08.014>.
20. Mariana Haro D. Laboratorio de análisis de marcha y movimiento. *Rev Med Clin Condes*. 2014;25:237-47, [http://dx.doi.org/10.1016/S0716-8640\(14\)70034-3](http://dx.doi.org/10.1016/S0716-8640(14)70034-3).
21. Willems P-A, Schepens B, Detrembleur C. Marcha normal. *EMC – Kinesiterapia – Med Física*. 2012;33:1-29, [http://dx.doi.org/10.1016/S1293-2965\(12\)61944-6](http://dx.doi.org/10.1016/S1293-2965(12)61944-6).
22. Duque-Ramírez JR, Vélez-Urbe JD, Gómez-Suárez NE, Manrique-Hernández RD. Gait parameters in a sample of healthy Colombian adults aged between 18 and 25 years: a cross-sectional study. *Rev Fac Med*. 2020;68:84-91, <http://dx.doi.org/10.15446/revfacmed.v68n1.71690>.
23. Ministerio de Salud y Protección Social. Sala situacional de la Población Adulta Mayor; 2018. Available from: <https://www.minsalud.gov.co/sites/rid/Lists/BibliotecaDigital/RIDE/DE/PS/sala-situacion-envejecimiento-2018.pdf>
24. Vicon. Full body modeling with plug-in Gait-Nexus 2.8. Vicon Documentation; 2018. Available from: <https://docs.vicon.com/display/Nexus28/Full+body+modeling+with+Plug-in+Gait>.
25. Foroughi N, Smith R, Vanwanseele B. The association of external knee adduction moment with biomechanical variables in osteoarthritis: a systematic review. *Knee*. 2009;16:303-9, <http://dx.doi.org/10.1016/j.knee.2008.12.007>.
26. Bennell KL, Bowles K-A, Wang Y, Cicuttini F, Davies-Tuck M, Hinman RS. Higher dynamic medial knee load predicts greater cartilage loss over 12 months in medial knee osteoarthritis. *Ann Rheum Dis*. 2011;70:1770-4, <http://dx.doi.org/10.1136/ard.2010.147082>.
27. Chehab EF, Favre J, Erhart-Hledik JC, Andriacchi TP. Baseline knee adduction and flexion moments during walking are both associated with 5 years cartilage changes in patients with medial knee osteoarthritis. *Osteoarthritis Cartilage*. 2014;22:1833-9, <http://dx.doi.org/10.1016/j.joca.2014.08.009>.
28. Kutzner I, Trepczynski A, Heller MO, Bergmann G. Knee adduction moment and medial contact force – facts about their correlation during gait. *PLOS ONE*. 2013;8:e81036, <http://dx.doi.org/10.1371/journal.pone.0081036>.
29. Richards RE, Andersen MS, Harlaar J, van den Noort JC. Relationship between knee joint contact forces and external knee joint moments in patients with medial knee osteoarthritis: effects of gait modifications. *Osteoarthritis Cartilage*. 2018;26:1203-14, <http://dx.doi.org/10.1016/j.joca.2018.04.011>.
30. Verlaan L, Boekesteijn RJ, Oomen PW, Liu WY, Peters MJM, Emans PJ, et al. Knee adduction moments are not increased in obese knee osteoarthritis patients during stair negotiation. *Gait Posture*. 2019;73:154-60, <http://dx.doi.org/10.1016/j.gaitpost.2019.07.192>.
31. Agudelo Mendoza AI, Briñez Santamaria TJ, Guarín Urrego V, Ruiz Restrepo JP, Zapata García MC. Marcha: descripción, métodos, herramientas de evaluación y parámetros de normalidad reportados en la literatura. *CES Mov Salud*. 2013;1:29-43.
32. Al-Obaidi S, Wall JC, Al-Yaqoub A, Al-Ghanim M. Basic gait parameters: a comparison of reference data for normal subjects 20 to 29 years of age from Kuwait and Scandinavia. *J Rehabil Res Dev*. 2003;40:361-6, <http://dx.doi.org/10.1682/jrrd.2003.07.0361>.
33. Oberg T, Karsznia A, Oberg K. Basic gait parameters: reference data for normal subjects, 10-79 years of age. *J Rehabil Res Dev*. 1993;30:210-23.