



Original Investigation

Impact of ultrasound on the perception of health quality in adult patients with rheumatoid arthritis[☆]



Luis Javier Cajas Santana^{a,*}, Rafael Barón^b, Carlos A. Guillén-Astete^c

^a Institute of Rheumatology Fernando Chalem, Bogota, Colombia

^b Foundation University of Health Sciences, Bogota, Colombia

^c Department of Rheumatology, Hospital Ramón y Cajal, Madrid, Spain

ARTICLE INFO

Article history:

Received 13 December 2019

Accepted 12 August 2020

Available online 16 March 2022

Keywords:

Rheumatoid arthritis

Ultrasound

Quality of health care

ABSTRACT

Introduction: Ultrasound has shown its usefulness in multiple aspects in the management of inflammatory joint disease and in particular of rheumatoid arthritis (RA). The evidence using patient outcomes and its aspects related to quality of health care is scarce. The aim of this study is to determine the level of satisfaction in the perception of the quality of health care in a group of patients with RA who underwent ultrasound during the consultation, and whether it is higher than those who did not have the ultrasound.

Methods: An observational, cross-sectional descriptive study was performed. Patients older than 18 years with a diagnosis of RA using the ACR/EULAR classification criteria were included. One group underwent skeletal muscle ultrasound to study RA during the outpatient medical consultation, as decided by the attending physician. After the completion of the medical action according to prior verbal acceptance by the patient, the Servqhos questionnaire and an ultrasound questionnaire were completed. A satisfied patient was defined as one who had 70% or more in the responses in the Servqhos questionnaire greater than or equal to 4, and a score of 5 in this questionnaire was defined as maximum satisfaction. There were no significant differences between the number of satisfied patients in the two groups. Univariate analysis was performed according to the distribution in the ultrasound or non-performing groups. Subsequently, a bivariate analysis of the different questions was carried out according to the distribution in the satisfaction and very high satisfaction groups. It was established if there was any degree of association using the Chi squared test for categorical variables, and the parametric tests (Mann Whitney U) or non-parametric tests (Kruskal-Wallis test) for the numerical variables were performed according to the distribution.

Results: A total of 126 patients were obtained, of whom 62 corresponded to the group of patients who underwent ultrasound during the consultation and 62 to the control group in whom no ultrasound was performed. The majority were women (91%). Ultrasound was

* Please cite this article as: Cajas Santana LJ, Barón R, Guillén-Astete CA. Impacto de la ecografía en la percepción de la calidad en salud de pacientes adultos con artritis reumatoide. Rev Colomb Reumatol. 2022;29:85–92.

[☆] Corresponding author.

E-mail address: ljaviercs.77@hotmail.com (L.J. Cajas Santana).

mostly performed to study joint disease (93%), with a third of the time to assess more than one aspect. In those on whom the ultrasound was performed, the number of satisfied patients was 56 (90%) and for the control group it was 48 (77%). The difference in the proportion of satisfied patients (13%) was statistically significant ($P = .05$). A difference was found between the groups in the number of patients with the highest level of satisfaction in the questions regarding presentation of staff and technology ($P < .05$). The vast majority of patients considered that ultrasound was useful during the consultation (93%), and that it generates greater confidence in the treatments and the doctor (93%).

Conclusions: Performing skeletal muscle ultrasound during consultation in patients with RA improves satisfaction rates of health care, perception of the doctor, and treatments.

© 2021 Asociación Colombiana de Reumatología. Published by Elsevier España, S.L.U. All rights reserved.

Impacto de la ecografía en la percepción de la calidad en salud de pacientes adultos con artritis reumatoide

RESUMEN

Palabras clave:

Artritis reumatoide
Ultrasonografía
Calidad de atención en salud

Introducción: La ecografía ha mostrado su utilidad en múltiples aspectos del manejo de la enfermedad articular inflamatoria, particularmente en la artritis reumatoide (AR). Su utilidad usando desenlaces derivados de pacientes y relacionados con aspectos de la calidad de la atención en salud es escasa. El objetivo del estudio es determinar si el grado de satisfacción de la calidad de la atención en un grupo de pacientes con AR, en quienes se realizó ecografía durante la consulta, es superior al de un grupo en los que esta no se llevó a cabo.

Métodos: Se realizó un estudio observacional, descriptivo transversal. Se incluyeron pacientes mayores de 18 años con diagnóstico de AR por criterios clasificatorios ACR/EULAR que posteriormente a la finalización del acto médico respondieron las preguntas del cuestionario Servqhos. Quienes se sometieron a ecografía musculoesquelética para estudio de AR durante la consulta ambulatoria, según decisión del médico tratante, respondieron adicionalmente el cuestionario de ecografía. Se define paciente satisfecho como aquel que tiene un 70% o más en las respuestas del cuestionario Servqhos con un puntaje mayor o igual a 4, y se define como la máxima satisfacción al puntaje de 5 en una pregunta de dicho cuestionario. Se determinó si había diferencias significativas entre las proporciones de pacientes satisfechos en los dos grupos con y sin ecografía. Se realizó un análisis univariado según la distribución en los grupos, y posteriormente se hizo un análisis bivariado de las diferentes preguntas según la distribución en los grupos de satisfacción y muy alta satisfacción. Se estableció si había algún grado de asociación con las pruebas de chi-cuadrado para las variables categóricas, en tanto que para las variables numéricas se llevaron a cabo pruebas paramétricas (U de Mann Whitney) y no paramétricas (test de Kruskal-Wallis), según la distribución.

Resultados: Se obtuvo un total de 126 pacientes, de los cuales 62 corresponden al grupo de aquellos en quienes se realizó ecografía durante la consulta, mientras que otros 62 hacen parte del grupo control, en quienes no se realizó ecografía. La mayoría eran mujeres (91%). En quienes se hizo la ecografía, el número de pacientes satisfechos fue de 56 (90%), en tanto que para el grupo control fue de 48 (77%). La diferencia en la proporción de pacientes satisfechos entre los grupos fue del 13%, siendo estadísticamente significativa ($p = 0,05$). Se encontró diferencia entre los grupos en las preguntas referentes a presentación del personal y la tecnología de los equipos ($p < 0,05$). La gran mayoría de los pacientes consideró que la ecografía fue útil durante la consulta (93%) y que genera mayor seguridad en los tratamientos y en el criterio médico (93%).

Conclusiones: La realización de ecografía musculoesquelética durante la consulta en pacientes con AR mejora los índices de satisfacción de atención en salud, así como la percepción del criterio médico y de los tratamientos.

© 2021 Asociación Colombiana de Reumatología. Publicado por Elsevier España, S.L.U.
Todos los derechos reservados.

Introduction

Ultrasound has shown its usefulness in the management of inflammatory joint disease, particularly of rheumatoid arthritis (RA),¹⁻⁶ both in the early and late phases, not only in the diagnostic process, but also in the assessment of the damage, the progression and the evaluation of treatments.

It has demonstrated to be more sensitive than the physical examination when evaluating the presence of synovitis,⁷⁻¹¹ and for this reason the scientific societies such as the European League Against Rheumatism (EULAR) include it in their recommendations to improve diagnostic certainty.⁹ In addition, in the established disease it detects subclinical activity, which is predictive of both structural damage and appearance of subsequent crises.¹²⁻¹⁴

However, the evidence of its usefulness using patient-derived outcomes is scarce.^{15,16} One study found that there was a relationship between the ultrasound activity status and anxiety, pain, and morning stiffness.¹⁵ Regarding the assessment of the quality of care or the influence that may exist on patients satisfaction with the use of ultrasound, three studies were found that evaluated both patients and physicians. Of these studies, two evaluated the satisfaction of the patient without annotated diagnoses in relation to the use of ultrasound for joint aspiration or intra-articular injection, and found that it helps the vast majority of them to understand the procedure and improves the anxiety related thereof. Ultrasound was perceived to be more effective than non-ultrasound-guided therapy and patients were more likely to undergo similar new procedures.^{17,18}

The third study analyzed the opinion of a group of patients with RA, regarding ultrasound, to which they were subjected rigorously to assess the activity of their disease. The authors found that most patients (80-90%) consider a reasonable ultrasound duration time to be 20-30 min. Likewise, they found greater satisfaction with the care and a perception of the treatment as safer.¹⁹ An inconvenience found was the use of non-endorsed or standardized tools for the evaluation of satisfaction.

The quality of service has several definitions; however, the most widely accepted is that of Parasuramam et al.,²⁰ which considers it a construct resulting from the comparison of the expectations or desires of the consumer vs. those of a provider and his/her perceptions of the service received.²¹ Given its importance, several methods have been developed to measure the quality of service, such as Servperf, Servqual²²⁻²⁴ and Servqhos,^{25,26} the latter for the specific measurement of the health sector. The Servqhos has been practiced in hospital, dialysis, surgical and outpatient areas,²⁷⁻³³ without results in the areas of rheumatology or diagnostic imaging. This instrument is validated in Colombia, where it has been already applied.³⁴

In the Colombian model, governed by Law 100 of 1993, the quality of health care is part of the aspects that must be measured by health providers to achieve quality standards and compete with each other for users.³⁵ The data show that 67% of the people identified as ill who did not use health services had as the first reason the poor quality of the service.³⁶ In addition, the Bulletin 018 of the Office of the Attorney General of

the Nation, 2007, considered that the main failings of health service provider companies were the inhumane treatment given to patients, the discrimination and the disproportionate demands to access the service.

The objectives of the study are to determine the degree of satisfaction with the health care of a group of patients with RA, who underwent an ultrasound during the consultation, as well as to evaluate whether it is superior compared to that of a group of patients who did not undergo this procedure, through the Servqhos questionnaire and with the hypothesis that ultrasound would have positive effects in the assessment of the quality of care and would improve the parameters of satisfaction.

Methods

An observational cross sectional descriptive study which included patients over 18 years of age with a diagnosis of RA by the ACR/EULAR classification criteria, who underwent a musculoskeletal ultrasound for the study of RA during the medical consultation in a Colombian institution in the period between March and August, 2019, was carried out. Patients with RA who did not undergo ultrasound during the consultation were included as a comparator group. Patients in whom ultrasound was performed as a guide to procedures or at a time other than the outpatient consultation were excluded, in addition to those who had difficulties or did not accept to answer the questionnaire.

At the discretion of the treating physician, during the consultation, it was considered to perform an ultrasound to determine clinical aspects related to RA or its musculoskeletal complications. The procedure was carried out by the treating physician himself, and there was no limit for the number of studies or time. A GE Logic S5 equipment, linear transducer of 7-13 MHz, gains of 50-70%, and Doppler with PRF 0.7, adjusted gain, was used. After the medical act and with the acceptance by the patient, the Servqhos questionnaire (Appendix B annex 1) was delivered, whether or not the ultrasound had been performed. In the event that the ultrasound had been carried out, the patient was also asked to answer the non-standardized questionnaire on the perception of ultrasound (Appendix B annex 2).

The Servqhos survey consists of 19 ordinal categorical variables that assess the quality of the healthcare service, each of which is scored on a Likert scale ranging from 1 (it has been much worse than I expected) to 5 (it has been much better than I expected). The ultrasound perception survey is conducted using five ordinal categorical variables, and each of these is scored on a Likert scale ranging from 1 (nothing) to 5 (much).

The information bias was controlled by training the interviewers. It was carried out a pilot test in which the procedures were standardized and changes were allowed in case that is required. Source biases were controlled by sensitizing all respondents on the need of answer the totality of the survey, in addition to provide assistance when necessary. A satisfied patient was defined as one who scored 70% or more on the questions of the Servqhos questionnaire with a score higher than or equal to 4. The maximum satisfaction was defined as a score of 5 in a question of this questionnaire. A sample

size was calculated based on the *a priori* hypothesis of finding differences of 20% between the proportion of satisfied vs. dissatisfied patients, with an alpha error of 0.05 and a two-tailed power of 80%. Thus, a size of 62 research subjects in each group was estimated. In addition, a description of the general population was carried out.

Subsequently, a univariate analysis was made according to the distribution in the groups with or without performance of ultrasound, after which a bivariate analysis of the different questions was performed according to the distribution in the groups of satisfaction and very high satisfaction. The variables were described according to their nature assessed by the Shapiro Wilk test: the categorical ones in relative and absolute frequencies, the numerical ones according to their distribution, the normal ones with mean and standard deviation, and the non-normal ones with median and interquartile range. It was established if there was some grade of association with the chi-square tests for the categorical variables, while in the case of the numerical ones, parametric (Mann Whitney U) and no parametric (Kruskal-Wallis) tests were performed according to the distribution.

The database was collected in Microsoft Excel and the data analysis was performed with the Stata 12.0 statistical software. The work follows the scientific, technical and administrative standards for health research, especially in human beings, set forth in Resolution 8430 of 1993 of the Colombian Ministry of Health, which are in relation to international regulations. It was considered a risk-free study and voluntary and anonymous participation was notified in the questionnaire.

Results

A total of 126 patients were obtained, of whom two were excluded from the analyzes given that they did not answer more than 50% of the questions. Of the remaining 124, 62 correspond to the group of patients to whom an ultrasound was performed during the consultation and 62 to the control group, in whom the procedure was not performed. The mean age of the ultrasound group was 57.4 years and that of the control group was 55.9, with no differences between the groups ($p = 0.36$). Regarding the gender, the majority of patients were women, 93.5% in the ultrasound group and 88% in the control group, with no differences among them ($p = 0.89$).

The reason why the physician considered to perform an ultrasound was for the study of joint pathology in 58 patients (93%), tendon pathology in 14 of them (22.5%) and study of bursae in 8 cases (12%). In 20 patients (32%) ultrasound was used to assess two or more of the described aspects.

In the completion of the Servqhos questionnaire, of a total of 1178 responses in each group, there was no response in 29 in the ultrasound group (2.4%) and in 29 in the control group (2.4%). Question number 18, which makes reference to information to the family, was the one that had the most missing data, with 19%, without being excluded from the analysis.

The results in the ultrasound group are presented in Table 1. The majority of responses were positive in almost all aspects of care. The questions featured with a high number of responses with the best value responses were: medical

equipment, 61%; appearance of the personnel, 67%; information of the physician, 77%; confidence in the doctor, 82%, and kindness of the staff, 72%.

The results of the Servqhos in the control group are presented in Table 2. Most responses were positive, similarly than in the ultrasound group. Based on these results, it was found that in those to whom the ultrasound was performed, the number of satisfied patients was 56 (90%), while in the control group it was 48 (77%).

The difference in the proportion of satisfied patients between the groups was 13%, being statistically significant in favor of the patients to whom the ultrasound was performed ($p = 0.05$). The analysis of the relationship between the age and the state of satisfaction did not show that there was such association ($p = 0.99$), as well as in the case of the gender variable ($p = 0.12$).

Table 3 summarizes the differences between the groups in specific questions of the Servqhos questionnaire that make reference to medical care, and that may be more related to the ultrasound procedure, taking into account the number of patients who rated them with the highest degree of satisfaction, defined as 5. Table 3 shows that in questions 1 and 2 of the questionnaire, which refer to the technology of the medical equipment and the appearance of the staff, there is a statistically significant difference ($p < 0.05$).

The non-standardized survey to assess the perception regarding ultrasound of the patients who underwent the procedure obtained the results that are summarized in Table 4. According to these results, there was a very good acceptance, considering that it was useful, with the highest qualification, in 93%; that greatly improves the safety of their treatments and of their physician, with 92%, and that improves care, with 93%.

Discussion

The level of satisfaction measured is high in both groups, but it is higher in the ultrasound group, being statistically significant, which may be due to the impact on people with the use of equipment during the medical act, the direct visualization of the pathology and the additional diagnostic and resolute possibility that ultrasound allows. In general, this enables the physician to report in a more explicit and detailed manner the health status of each patient, which could be reflected in the analysis of the individual assessment of variables considered more related to the medical act and ultrasound, such as the use of medical equipment and the presentation of the staff, which were higher in the intervention group, in addition to a tendency to be higher in others, such as the medical information or the friendliness of the staff. Likewise, the vast majority of patients consider that it is useful, generates more confidence in the physician and the treatment, in addition to improving care. Similar results were obtained in studies such as the one of Ceponis et al.¹⁹ in patients also with RA who underwent ultrasound at a time different from that of the medical consultation.

This is the first study that shows that ultrasound improves the parameters related to the quality in the provision of medical service and their perception by the patient, which is essential for the individual well-being, favorable results

Table 1 – Distribution of data from the Servqhos survey in the ultrasound group (n = 62).

Variable	1	2	3	4	5	0	M	SD	V
Technology of the medical equipment for diagnoses and treatments	0	0	3 (4.8%)	20 (32%)	38 (61%)	1 (1.6%)	4.5	0.5	0.3
Appearance of the staff	0	0	4 (6.5%)	16 (25%)	42 (67%)	0	4.6	0.6	0.3
Indications for orientation in the institution	0	0	5 (8.0%)	22 (35%)	34 (54%)	1 (1.6%)	4.4	0.6	0.3
If the staff do what they say they are going to do	0	0	4 (6.5%)	14 (23%)	44 (71%)	0	4.6	0.6	0.3
The condition of the consulting rooms	0	1 (1.6%)	7 (11%)	16 (25%)	37 (60%)	1 (1.6%)	4.4	0.7	0.5
Information provided by the doctors	0	1 (1.6%)	3 (4.8%)	10 (16%)	48 (77%)	0	4.6	0.6	0.4
Waiting time to be seen by the doctor	0	2 (3.2%)	12 (19%)	10 (10%)	38 (61%)	0	4.3	0.9	0.8
Ease to get to the hospital	0	2 (3.2%)	10 (16%)	17 (27%)	28 (45%)	5 (8.0%)	4.2	0.8	0.7
Interest of the staff to solve the problems of the patients	0	1 (1.6%)	2 (3.2%)	21 (33%)	38 (61%)	0	4.5	0.6	0.4
Punctuality of the medical consultations	0	3 (4.8%)	5 (8.0%)	15 (24%)	39 (62%)	0	4.4	0.8	0.7
How quickly the user gets what he/she needs or requests	0	1 (1.6%)	8 (12%)	18 (29%)	35 (56%)	0	4.4	0.7	0.6
Willingness of the staff to help the user when he/she needs it	0	0	5 (8.0%)	15 (24%)	41 (66%)	1 (1.6%)	4.5	0.6	0.4
The confidence (security) that the doctor transmits to the patients	0	0	1 (1.6%)	10 (16%)	51 (82%)	0	4.8	0.4	0.1
Friendliness (courtesy) of the staff in their treatment of people	0	0	1 (1.6%)	16 (25%)	45 (72%)	0	4.7	0.4	0.2
Preparation of the staff to perform their work	0	0	2 (3.2%)	20 (32%)	40 (64%)	0	4.6	0.5	0.3
The personalized treatment given to their patients	0	1 (1.6%)	0	18 (29%)	42 (67%)	1 (1.6%)	4.6	0.5	0.3
Ability of the staff to understand the needs of the patients	0	1 (1.6%)	3 (4.8%)	14 (22%)	43 (69%)	1 (1.6%)	4.6	0.6	0.4
The information that doctors give to the relatives	0	1 (1.6%)	6 (9.6%)	10 (16%)	28 (45%)	17 (27%)	4.4	0.8	0.6
Interest of non-medical staff in the patients	0	0	6 (9.6%)	17 (27%)	38 (61%)	1 (1.6%)	4.5	0.6	0.4

SD: standard deviation; M: mean; V: variance.

0. No response (%); 1. Much worse than expected (%); 2. Worse than expected (%); 3. As expected (%); 4. Better than expected (%); 5. Much better than expected (%).

Table 2 – Distribution of data from the Servqhos survey in the control group (n = 62).

Variable	1	2	3	4	5	0	M	SD	V
Technology of the medical equipment for diagnoses and treatments	1 (1.6%)	0	13 (20%)	18 (29%)	20 (32%)	10 (16%)	4.0	0.9	0.8
Appearance of the staff	0	0	10 (16%)	20 (32%)	30 (48%)	2 (3.2%)	4.3	0.7	0.5
Indications for orientation in the institution	0	1 (1.6%)	10 (16%)	19 (30%)	31 (50%)	1 (1.6%)	4.3	0.8	0.6
If the staff do what they say they are going to do	0	1 (1.6%)	6 (9%)	21 (33%)	33 (53%)	1 (1.6%)	4.4	0.7	0.5
The condition of the consulting rooms	0	1 (1.6%)	14 (22%)	13 (20%)	34 (54%)	0	4.2	0.8	0.7
Information provided by the doctors	0	0	8 (12%)	13 (20%)	40 (64%)	1 (1.6%)	4.5	0.7	0.5
Waiting time to be seen by the doctor	2 (3.2%)	0	8 (12%)	20 (32%)	32 (51%)	0	4.2	0.9	0.8
Ease to get to the hospital	1 (1.6%)	0	9 (14%)	21 (33%)	30 (48%)	1 (1.6%)	4.2	0.8	0.7
Interest of the staff to solve the problems of the patients	1 (1.6%)	1 (1.6%)	8 (12%)	13 (20%)	38 (61%)	1 (1.6%)	4.4	0.9	0.8
Punctuality of the medical consultations	2 (3.2%)	1 (1.6%)	8 (12%)	15 (24%)	35 (56%)	1 (1.6%)	4.3	1	1
How quickly the user gets what he needs or requests	1 (1.6%)	0	13 (20%)	12 (19%)	34 (54%)	2 (3.2%)	4.3	0.9	0.8
Willingness of the staff to help the user when he/she needs it	1 (1.6%)	0	6 (9%)	15 (24%)	40 (64%)	0	4.5	0.8	0.6
The confidence (security) that the doctor transmits to the patients	0	0	3 (5%)	13 (20%)	46 (75%)	0	4.6	0.5	0.3
Friendliness (courtesy) of the staff in their treatment of people	0	0	7 (11%)	10 (16%)	45 (72%)	0	4.6	0.6	0.4
Preparation of the staff to perform their work	0	0	7 (11%)	12 (19%)	43 (69%)	0	4.5	0.6	0.4
The personalized treatment given to their patients	0	0	7 (11%)	9 (14%)	46 (74%)	0	4.6	0.6	0.4
Ability of the staff to understand the needs of the patients	0	7 (11%)	6 (9%)	13 (20%)	42 (67%)	0	4.5	0.7	0.5
The information that the doctors give to the relatives	0	0	5 (8%)	14 (22%)	35 (56%)	8 (12%)	4.5	0.6	0.4
Interest of non-medical staff in the patients	0	1 (1.6%)	10 (16%)	19 (30%)	31 (50%)	1 (1.6%)	4.3	0.8	0.6

SD: standard deviation; M: mean; V: variance.

0. No response (%); 1. Much worse than expected (%); 2. Worse than expected (%); 3. As expected (%); 4. Better than expected (%); 5. Much better than I expected (%).

Table 3 – High level of satisfaction in each specific question of the Servqhos.

Variable	Eco (n = 62)	No eco (n = 62)	p
Technology of the medical equipment for diagnoses and treatments	38 (62%)	20 (38%)	0.01
Appearance of the staff	42 (67%)	30 (50%)	0.04
Information provided by the doctors	48 (77%)	40 (65%)	0.14
Waiting time to be seen by the doctor	38 (61%)	32 (51%)	0.27
Interest of the staff to solve the problems of the patients	38 (61%)	38 (61%)	0.9
Punctuality of the medical consultations	39 (62%)	35 (57%)	0.53
The confidence (security) that the doctor transmits to the patients	51 (82%)	46 (74%)	0.43
Friendliness (courtesy) of the staff in their treatment of people	45 (72%)	45 (72%)	1
Preparation of the staff to perform their work	40 (64%)	43 (69%)	0.56
The personalized treatment given to their patients	42 (67%)	46 (74%)	0.51
Ability of the staff to understand the needs of the patients	43 (69%)	42 (67%)	0.74

Table 4 – Distribution of the data of the survey on ultrasound (n = 62).

Variable/n-value (%)	1	2	3	4	5	0	M	SD	V
Consider that the ultrasound was useful during the consultation	0	1	1	2	58 (93%)	0	4.8	0.48	0.23
Consider that ultrasound gives more security in their treatments	0	0	1	3	57 (92%)	1	4.9	0.34	0.12
Consider that ultrasound wastes time unnecessarily	47 (75%)	3	0	2	4	6	1.4	1.1	1.3
Consider that ultrasound gives more security of their doctor	0	0	1	0	58 (93%)	3	4.9	0.3	0.9
Consider that ultrasound improves the healthcare	0	0	1	2	58 (93%)	1	4.9	0.3	0.1

SD: standard deviation; M: mean; V: variance.

1. Nothing; 2. Almost nothing; 3. Few; 4. More than few; 5. Much.

also demonstrated in the use of ultrasound for patients who undergo arthrocentesis.^{17,18}

One of the strengths of the study is the use of the Servqhos tool, an instrument validated in various populations, including that of Colombia,³⁴ which will allow to make comparisons in future studies, in addition to the use of musculoskeletal ultrasound during consultation by the treating physician himself, which is important for the short-term medical decision making.

Regarding the limitations, there is a possible selection bias, since there is no control over whom the ultrasound is performed or the reasons for its use. Likewise, it cannot be generalized to other population groups, with diagnoses other than RA, a high degree of satisfaction with care in general, which may make the difference much smaller, and the use of the non-validated survey for the perception of ultrasound.

The conclusion of the study is that in patients with RA, musculoskeletal ultrasound at the time of medical consultation improves the perception of quality in health in general, as well as the perception specifically regarding the medical equipment and the presentation of the personnel. In addition, the perception of the ultrasound in those who undergo the procedure is good, it gives them more confidence in their doctor and improves the quality of care.

Conflict of interest

The authors declare that they have no conflict of interest.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.rcreue.2020.08.005>.

REFERENCES

1. Mandl P, Navarro-Compan V, Terslev I, Aegeuter P, van der Heijde D, d'Agostino MA, et al. EULAR recommendations for the use of imaging in the diagnosis and management of spondyloarthritis in clinical practice. Ann Rheum Dis. 2015;74:1327-39, <http://dx.doi.org/10.1136/annrheumdis-2014-206971>.
2. Combe B, Landewe R, Daien CI, Hua C, Aletaha D, Álvaro-Gracia JM, et al. 2016 update of the EULAR recommendations for the management of early arthritis. Ann Rheum Dis. 2017;76:948-59, <http://dx.doi.org/10.1136/annrheumdis-2016-210602>.
3. Nam JL, d'Agostino MA. Role of ultrasound imaging in individuals at risk of RA. Best Pract Res Clin Rheumatol. 2017;31:71-9, <http://dx.doi.org/10.1016/j.berh.2017.09.002>.
4. Van der Ven M, van der Veer-Meerkirk M, ten Cate DF, Rasappu N, Kok MR, Csakvari D, et al. Absence of ultrasound inflammation in patients presenting with arthralgia rules out the development of arthritis. Arthritis Res Ther. 2017;19:202, <http://dx.doi.org/10.1186/s13075-017-1405-y>.
5. D'Agostino MA, Terslev I, Wakefield R, Østergaard M, Balint P, Naredo E, et al. Novel algorithms for the pragmatic use of ultrasound in the management of patients with rheumatoid arthritis: From diagnosis to remission. Ann Rheum Dis. 2016;75:1902-8, <http://dx.doi.org/10.1136/annrheumdis-2016-209646>.

6. Horton SC, Wakefield RJ, Freeston JE, Buch MH, Emery P. Ultrasound-detectable grey scale synovitis predicts future fulfilment of the 2010 ACR/ EULAR RA classification criteria in patients with new-onset undifferentiated arthritis. *RMD Open.* 2017;3:e000394, <http://dx.doi.org/10.1136/rmdopen-2016-000394>.
7. Wakefield RJ, Green MJ, Marzo-Hor-Tega H, Conaghan PG, Gibbon WW, McGonagle D, et al. Should oligoarthritis be reclassified? Ultrasound reveals high prevalence of subclinical disease. *Ann Rheum Dis.* 2004;63:382-5, <http://dx.doi.org/10.1136/ard.2003.007062>.
8. Szkudlarek M, Narvestad E, Klarlund M, Court-Payen M, Thomsen HS, Østergaard M. Ultrasonography of the metatarsophalangeal joints in rheumatoid arthritis: Comparison with magnetic resonance imaging, conventional radiography, and clinical examination. *Arthritis Rheum.* 2004;50:2103-12, <http://dx.doi.org/10.1002/art.20333>.
9. Colebatch AN, Edwards CJ, Ostergaard M, van der Heijde D, Balint PV, D'Agostino MA, et al. EULAR recommendations for the use of imaging of the joints in the clinical management of rheumatoid arthritis. *Ann Rheum Dis.* 2013;72:804-14, <http://dx.doi.org/10.1136/annrheumdis-2012-203158>.
10. Lage-Hansen PR, Lindegaard H, Chrysidis S, Terslev L. The role of ultrasound in diagnosing rheumatoid arthritis, what do we know? An updated review. *Rheumatol Int.* 2017;37:179-87, <http://dx.doi.org/10.1007/s00296-016-3587-z>.
11. Takase-Minegishi K, Horita N, Kobayashi K, Yoshimi R, Kirino Y, Ohno S, et al. Diagnostic test accuracy of ultrasound for synovitis in rheumatoid arthritis: systematic review and meta-analysis. *Rheumatology (Oxford).* 2018;57:49-58, <http://dx.doi.org/10.1093/rheumatology/kex036>.
12. Brown AK, Conaghan PG, Karim Z, Quinn MA, Ikeda K, Peterfy CG, et al. An explanation for the apparent dissociation between clinical remission and continued structural deterioration in rheumatoid arthritis. *Arthritis Rheum.* 2008;58:2958-67, <http://dx.doi.org/10.1002/art.23945>.
13. Scirè CA, Montecucco C, Codullo V, Epis O, Todoerti M, Caporali R. Ultrasonographic evaluation of joint involvement in early rheumatoid arthritis in clinical remission: Power Doppler signal predicts short-term relapse. *Rheumatology (Oxford).* 2009;48:1092-7, <http://dx.doi.org/10.1093/rheumatology/kep171>.
14. Lamers-Karnebeek FB, Luime JJ, Ten Kate DF, Teerenstra S, NWAA Swen, Gerards AH, et al. Limited value for ultrasonography in predicting flare in rheumatoid arthritis patients with low disease activity stopping TNF inhibitors. *Rheumatology (Oxford).* 2017;56:1560-5, <http://dx.doi.org/10.1093/rheumatology/kex184>.
15. Van der Ven M, Kuijper TM, Gerards AH, Tchetverikov I, Weel AE, van Zeben J, et al. No clear association between ultrasound remission and health status in rheumatoid arthritis patients in clinical remission. *Rheumatology (Oxford).* 2017;56:1276-81, <http://dx.doi.org/10.1093/rheumatology/kex080>.
16. Závada J, Hánová P, Hurňáková J, Szczuková L, Uher M, Forejtová Š, et al. The relationship between synovitis quantified by an ultrasound 7-joint inflammation score and physical disability in rheumatoid arthritis - A cohort study. *Arthritis Res Ther.* 2017;19, <http://dx.doi.org/10.1186/s13075-016-1208-6>.
17. Sahbudin I, Bell J, Kumar K, Raza K, Filer A. Observing real-time images during ultrasound-guided procedures improves patients' experience. *Rheumatology (Oxford).* 2016;55:585-6, <http://dx.doi.org/10.1093/rheumatology/kev368>.
18. El Miedany Y. Musculoskeletal US: Examining the joints. *Br J Nurs.* 2012;21:340-4, <http://dx.doi.org/10.12968/bjon.2012.21.6.340>.
19. Ceponis A, Onishi M, Bluestein HG, Kalunian K, Townsend J, Kavanaugh A. Utility of the ultrasound examination of the hand and wrist joints in the management of established rheumatoid arthritis. *Arthritis Care Res (Hoboken).* 2014;66:236-44, <http://dx.doi.org/10.1002/acr.2219>.
20. Parasuraman A, Zeithaml V, Berry L. A conceptual model of service quality and its implications in future research. *J Mark.* 1985;49:41-50, <http://dx.doi.org/10.2307/1251430>.
21. Losada M, Rodríguez A. Calidad del servicio de salud: una revisión a la literatura desde la perspectiva del marketing. *Cuad Adm (Bogotá).* 2007;20:237-58.
22. Parasuraman A, Zeithaml VA, Berry LL. Servqual: A multiple-item scale for measuring consumer perceptions of service quality. *J Retail.* 1988;64:12-40.
23. Babakus E, Mangold WG. Adapting the Servqual scale to hospital services: An empirical investigation. *Health Serv Res.* 1992;26:767-86.
24. Bowers MR, Swan JE, Koehler WF. What attributes determine quality and satisfaction with health care delivery? *Health Care Manage Rev.* 1994;19:45-9.
25. Mira JJ, Aranaz J, Rodríguez-Marín J, Buil JA, Castell M, Vialler J. Servqhos: un cuestionario para evaluar la calidad percibida de la asistencia hospitalaria. *Medicina Preventiva.* 1998; 4:8-12.
26. Numpaque-Pacabaque A, Rocha-Buelvas A. Modelos Servqual y Servqhos para la evaluación de calidad de los servicios de salud. *Rev Fac Med.* 2016;64:715-20, <http://dx.doi.org/10.15446/revfacmed.v64n4.54839>.
27. Barragán JA, Manrique FG. Validez y confiabilidad del Servqhos para enfermería en Boyacá, Colombia. *Av Enferm.* 2010;28:48-61.
28. Barragán JA, Moreno C. Calidad percibida por usuarios de enfermería en tres hospitales públicos. *Enferm Global.* 2013;12:217-30.
29. Guinovart E, Bateman F. Estudio multicéntrico sobre la calidad asistencial percibida por los pacientes en programa de hemodiálisis de Tarragona. *Rev Soc Esp Enferm Nefrol.* 2011;14:172-80, <http://dx.doi.org/10.4321/S1139-13752011000300005>.
30. Rodríguez-González D, Sanjuán-Hernández D. Satisfacción percibida por los pacientes en el servicio de hemodiálisis del hospital general de La Palma. *Enferm Nefrol.* 2013;16:75-82, <http://dx.doi.org/10.4321/S2254-28842013000200002>.
31. Braña-Marcos B, Carrera-Martínez D, de la Villa-Santoveña M, Avanzas-Fernández S, Gracia-Corbato M, Vegas-Pardavila E. Analysis of satisfaction in women with recent diagnosis of breast cancer. Multicenter study in oncology services. *Rev Calid Asist.* 2011;26:305-14, <http://dx.doi.org/10.1016/j.cal.2011.04.006>.
32. Jorge-Cerrudo J, Ramón JM, Goday A, González S, Sánchez P, Benaiges D, et al. Estudio descriptivo de la mejora de la calidad percibida en los pacientes sometidos a cirugía bariátrica tras la implantación de una vía clínica. *Rev Calid Asist.* 2011;27:226-32, <http://dx.doi.org/10.1016/j.cal.2012.01.007>.
33. Hidalgo M, Rodríguez C, López M, Moreno E, Pulido M. Valoración de la satisfacción materna con la analgesia epidural para el trabajo de parto. *Rev Calid Asist.* 2011;26:251-5, <http://dx.doi.org/10.1016/j.cal.2011.03.003>.
34. Henao Nieto DE, Giraldo Villa A, Yépes Delgado CE. Instrumentos para evaluar la calidad percibida por los usuarios en los servicios de salud. *Rev Gerenc Polit Salud.* 2018;17:69-80, <http://dx.doi.org/10.11144/Javeriana.rgps17-34.iecp>.
35. Cardona J, Hernández A, Yépes F. La reforma a la seguridad social en salud en Colombia: ¿avances reales? *Rev Gerenc Polit Salud.* 2005;4:81-99.
36. Velandia F, Ardón N, Cardona JM, Jara MI, Pérez N. Oportunidad, satisfacción y razones de no uso de los servicios de salud en Colombia, según la encuesta de calidad de vida del DANE. *Colombia Médica.* 2001;32:4-9.