

ORIGINAL RESEARCH

DOI: <http://dx.doi.org/10.15446/revfacmed.v66n4.66132>

Association between metabolic risk markers, body composition and different manifestations of strength in young adults

Asociación entre marcadores de riesgo metabólicos y diferentes manifestaciones de la fuerza en adultos jóvenes

Received: 05/07/2017. Accepted: 15/11/2017.

Jairo Alejandro Fernández-Ortega¹ • Luz Amelia Hoyos-Cuartas² Flor Alba Ruiz-Arias³¹ Universidad Pedagógica Nacional - Faculty of Physical Education - Exercise Physiology Laboratory - Bogotá D.C. - Colombia.² Universidad Pedagógica Nacional - Faculty of Physical Education - Bachelor's degree in Sports - Bogotá D.C. - Colombia.³ Colegio Mayor Nuestra Señora del Rosario - Faculty of Natural Sciences and Mathematics - Department of Mathematics - Bogotá D.C. - Colombia.

Corresponding author: Jairo Alejandro Fernández-Ortega. Faculty of Physical Education, Universidad Pedagógica Nacional. Calle 183 No. 54D-esquina. Telephone number: +57 1 3471153, ext.: 220. Bogotá D.C. Colombia. Email: jairofdz@pedagogica.edu.co.

| Abstract |

Introduction: For decades, low levels of muscle strength (MS) have been associated with an increased risk of mortality from all causes.

Objective: To identify the correlation between values of different manifestations of MS and metabolic risk markers (MRM) in young university students.

Materials and methods: The study included 50 participants (37 men, 13 women), aged between 19 and 23 years. The grip strength of both hands and the muscular strength of upper and lower limbs were evaluated, as well as height, weight, body mass index (BMI), skin folds, HDL, cholesterol and triglycerides.

Results: No significant associations between MRM and the different manifestations of MS were found.

Conclusion: MS and MRM are separately associated with mortality risk factors and with suffering from cardiovascular diseases

Keywords: Triglycerides; Cholesterol; Lipoproteins, HDL; Muscle (MeSH).

| Resumen |

Introducción. Desde hace varias décadas, los bajos niveles de fuerza muscular (FM) han sido asociados al incremento en el riesgo de mortalidad por todas las causas.

Objetivo. Identificar la relación entre los valores de diferentes manifestaciones de FM con biomarcadores de riesgo metabólico (BRM).

Materiales y métodos. El estudio estuvo compuesto por 50 participantes (37 hombres, 13 mujeres) con edades entre 19 y 23 años. Se evaluó la fuerza prensil de ambas manos, la potencia y la fuerza máxima de miembros superiores e inferiores, colesterol y triglicéridos. Además, se midió talla, peso, índice de masa corporal y pliegues cutáneos.

Resultados. No se identificaron relaciones significativas entre los BRM y las diferentes manifestaciones de la FM.

Conclusión. La FM y los BRM se relacionan con el riesgo de mortalidad y enfermedades cardiovasculares de forma separada.

Palabras clave: Triglicéridos; Colesterol; Lipoproteínas HDL; Fuerza muscular (DeCS).

.....
Fernández-Ortega JA, Hoyos-CuartasLA, Ruiz-Arias FA. Association between metabolic risk markers, body composition and different manifestations of strength in young adults. Rev. Fac. Med. 2018;66(4):565-70. English. doi: <http://dx.doi.org/10.15446/revfacmed.v66n4.66132>.

.....
Fernández-Ortega JA, Hoyos-CuartasLA, Ruiz-Arias FA. [Asociación entre marcadores de riesgo metabólicos y diferentes manifestaciones de la fuerza en adultos jóvenes]. Rev. Fac. Med. 2018;66(4):565-70. English. doi: <http://dx.doi.org/10.15446/revfacmed.v66n4.66132>.

Introduction

Among physical characteristics, muscle strength (MS) has gained great importance within the context of health thanks to the progress made regarding knowledge of muscle functions, as it is considered an endocrine and immune organ and because of its role in metabolism. (1) At first, muscles were thought to be involved only in movement

and regulation of energy metabolism; however, in the last decade, new evidence has suggested that skeletal muscle acts as an endocrine organ because it expresses and produces several cytokines and peptides in response to muscle contractions. (2,3)

Currently, there is sufficient evidence of the association between MS and the reduction of the risk of mortality from all causes. For decades, good levels of MS have been associated with a preventive

effect against chronic and metabolic diseases, while low muscle performance is associated with metabolic syndrome, cancer and increased risk of mortality from all causes, both in men and women. (4-7) This association has been confirmed in the meta-analyses of Volaklis *et al.* (8) and Cooper *et al.* (9), who observed an inverse correlation between MS and mortality, even after adjusting factors for some confounding variables such as physical activity levels and cardiorespiratory fitness, both in adult and younger populations.

Timpka *et al.* (10) conducted a prospective study with 38 588 Swedish adolescents (16-19 years), and concluded that the risk of mortality from all causes decreased by 20% in participants with the best results in grip strength and leg extension, compared to those who obtained low MS results. Other studies (11-12) reported similar results and identified that subjects with moderate or high muscular capacity have a risk reduction of 35% and 44%, respectively, compared to those with low MS levels. However, up to date, few studies have evaluated the association between muscle strength and metabolic syndrome. (13)

The studies by Jurca *et al.* (14) and Aoyama *et al.* (15) reported an inverse correlation between MS and metabolic risk ($\beta = -0.37$; $p < 0.0001$), which was marginally non-significant after further adjustment with aerobic fitness.

For this review, several search strategies were used including searches in specialized journals, in the Cochrane Central Register of Controlled Trials, and in the MEDLINE, PubMed, ProQuest, Hinari, SPORTDiscus, Embase, CINAHL, PsycINFO, PEDro and Ovid databases. After gathering all the data, it was observed that, until now, few studies have evaluated the association between MS and metabolic syndrome, and the results are contradictory. These observations were confirmed by Wijndaele *et al.* (13) and a recent meta-analysis conducted by Volaklis *et al.* (8)

In the Colombian context, several studies have addressed this issue. Ramírez-Vélez *et al.* (16), in a cohort study conducted with 6 095 adults aged between 18 and 40 years, observed that the group of participants with low and moderate levels of MS/body mass (BM) presented higher values in the cardiometabolic risk score index, regardless of their body mass index (BMI) ($p < 0.001$). On the other hand, the group with high MS/BM values had higher levels of cardiometabolic protection.

On the other hand, García-Hermoso *et al.* (17) demonstrated that adults with good MS levels and low fat have significantly lower levels of metabolic risk compared to those with low MS levels. Ramírez-Vélez *et al.* (18) studied the association between MS and cardiovascular health in 1 199 children and young people and observed that high levels of MS (absolute or relative) are significantly ($p < 0.001$) associated with better cardiovascular health.

The purpose of this study was to identify the correlation between the different manifestations of MS and the plasma concentrations of cholesterol, triglycerides, HDL-C and LDL-C in a group of young adults. This approach has been little studied, since most research works focus on the correlation between aerobic capacity and metabolic risk biomarkers, whose results are contrary to those related with MS, as concluded by Jurca *et al.* (14), who report that MS and cardiorespiratory fitness are independently and inversely related to metabolic syndrome.

Materials and methods

Design and population

This is an observational, descriptive and correlational cross-sectional study that was carried out in the Exercise Physiology Laboratory of

Universidad Pedagógica Nacional in 2016. The group of participants consisted of 50 university students (37 men and 13 women) aged between 19 and 25 years, who participated voluntarily in the study. Inclusion criteria were: being a university student, men and women with low levels of physical activity and capable of performing physical exercise; exclusion criteria were: suffering from any type of pathology, being habitual smokers or drinkers, and having participated or being involved in structured programs of physical exercise or training in the last year.

This study was designed following the deontological norms recognized by the Declaration of Helsinki (19) and the Resolution 8430 of 1993 of the Ministry of Health of Colombia (20), which regulate clinical research in humans. The Research Ethics Committee of the Universidad Pedagógica Nacional (VGU-400, March 28, 2017) approved the study. In addition, all participants were informed of the details of the study and signed the informed consent.

Variables

For the development of the study, the following variables were evaluated:

Body composition

Anthropometric measurements were total BM, evaluated using an electronic scale (Healht Metter 599 KL with 0.50g precision); height, measured with a height rod (Detecto D52, USA); and BMI, estimated using the formula $BMI = \text{weight (kg)} / \text{size (m)}^2$.

Maximum strength

Days before the assessments, the procedure of each MS and peak anaerobic power test was carefully explained to the subjects. Six minutes of specific warming-up of the involved muscle groups was performed before the evaluations.

The maximum muscular strength of the upper limbs was evaluated using the bench press test on a Smith machine by directly determining the one-repetition maximum (1RM). The test began with a load of 40% of the body weight of each subject and the participants were asked to perform 4-6 repetitions (21); at the end, the OMNI-RES scale was applied. (22) According to the score assigned to the effort, progressive increases of 10% were made. Maximum strength was determined when the subjects were only able to perform one repetition-maximum with said load. There was a 3-minute recovery period between each series to avoid fatigue.

The maximum strength of the lower limbs was evaluated performing a half squat on a Smith machine, using the same upper limb protocol, but starting with a load of 60% of body weight.

Peak anaerobic power

Peak anaerobic power of upper limbs was assessed using the Wingate test on a Monark 871E bicycle (Monark exercise, Varberg Sweden). Height and distance between the subject and the pedals of the bicycle were adjusted in order to align the glenohumeral joint with the center of the pedals; then three warming-up rounds were performed at 40RPM with a load of 0.10% of body weight and a sprint was executed for 5 seconds at minutes 2, 3 and 4. After 3 minutes of recovery, the participants performed the test using a load of 4.5% of body weight for men and 3.5% for women. (23)

Peak anaerobic power of the lower limbs was assessed by using the Wingate test on a Monark 834E bicycle (Monark exercise,

Varberg Sweden). Previously, the participants warmed-up for 5 minutes at 40RPM, with a load of 0.30% of body weight and a sprint was executed for 5 seconds at minutes 2, 3 and 4. After 3 minutes of recovery, the participants performed the test using a load of 7.5% of body weight for men and 6.7% for women. Saddle height was adjusted to ensure that knee flexion was not greater than 5 degrees.

The Optojump Microgate® Germany was used to estimate flight times (precision of 1/1000 seconds) with two types of jumps: countermovement jump (CMJ) and squat jump (SJ). Two attempts of each type of jump were made at intervals of 6 minutes between both, selecting the best record for statistical analysis.

Isometric strength

Grip strength of both hands was assessed by dynamometry method, using a Takei Grip-A dynamometer T.K.K 5001 (Scientific Instruments Co. Ltd, Tokyo, Japan) with a measuring range of 0-100kg and precision of 0.5kg. Two attempts were made with each hand, with recovery periods of 3 minutes; the best result was recorded. Likewise, isometric leg strength was evaluated with a Takei Back-A dynamometer T.K.K 5002 (Scientific Instruments Co. Ltd., Tokyo, Japan) with a measuring range of 0-300kg and precision of 0.5kg, using the same procedure.

Metabolic biomarkers

The analysis of plasma concentrations of triglycerides, HDL and total cholesterol (TC) was carried out on an empty stomach, asking the subjects not to consume any food after 11:00 pm the night before taking the blood sample. Blood was extracted from the cephalic vein, placing the subjects in supine position. Triglycerides were analyzed using the lipase/glycerol kinase/glycerol phosphate oxidase enzymatic method. HDL-C cholesterol was analyzed using the polyanion/cholesterol esterase/homogeneous oxidase enzymatic method. Measurements were made on an Olympus AU5400 analyzer (Olympus Diagnostica, Hamburg, Germany).

Statistical analysis

The results were analyzed using the SPSS software (version 20.0, SPSS, Inc., Chicago, IL). For the anthropometric, clinical, biochemical and strength condition variables, mean, standard deviation and coefficient of variation were calculated for each study group. On the other hand, to determine the correlations between variables of MS and metabolic risk markers, bivariate correlations were calculated through the Pearson coefficient for the total number of participants in the sample. The normality of the variables was verified by Kolmogorov-Smirnov test, using a level of significance of 5%. A qualitative rating scale was used according to the magnitude of the correlation observed: weak for values >0.40; moderate, between 0.41 and 0.60; strong, between 0.61 and 0.80; and very strong, between 0.81 and 1.

Results

The population that participated in the study had an average age of 21.8±2.3 years and 74% were men. Table 1 shows the descriptive statistics of the general anthropometric, biochemical, strength and peak anaerobic power characteristics of the study population.

Table 1. Morphological, biochemical, strength and peak anaerobic power characteristics of the population evaluated.

Characteristic	Men		Women		Difference by sex	%
	(n=37)	CV	(n=13)	CV		
Age (years)	21.4±2.9	0.13	22.2±4.8	0.22	0.8	4%
Weight (kg)	64.7±6.3	0.1	57±6.3	0.11	7.7	12%
Height (cm)	171±0.5	0.03	160±10	0.04	0.11	6%
Body mass index (kg/m ²)	22.2±2	0.09	22.1±1.9	0.09	0.1	0%
Sum of skinfolds (mm)	102.8±31.8	0.31	175±30	0.17	72.3	41%
Triglycerides (mg/dl)	103.7±63.9	0.62	89.9±48.9	0.54	13.8	13%
HDL (mg/dl)	43.4±8	0.18	53.9±11.2	0.21	10.5	19%
Cholesterol (mg/dl)	167.9±37.4	0.22	169±35.1	0.21	1.4	1%
Grip strength in left hand (kgf)	39.1±6.3	0.16	30.6±4.4	0.15	8.5	22%
Grip strength in right hand (kgf)	40.3±7.4	0.18	31.9±4.7	0.15	8.4	21%
Isometric strength of lower limbs (kgf)	145.9±35.5	0.24	111±32.2	0.29	34	23%
Half squat (kg)	116.6±17.4	0.15	85.2±18.3	0.22	31.4	27%
Bench press (kg)	61.7±9.9	0.16	36.2±10.1	0.28	25.5	41%
Peak anaerobic power upper limbs (watts)	250.6±50.7	0.2	136±33.2	0.24	114.2	46%
Relative peak power upper limbs (watts/kg)	3.7±0.9	0.24	2.3±0.4	0.19	1.4	38%
Peak anaerobic power lower limbs (watts)	513±87.7	0.17	360±69.8	0.19	152.3	30%
Relative peak power lower limbs (watts/kg)	7.7±1.3	0.17	6.1±1	0.17	1.6	21%
Countermovement jump (cm)	26.4±5.3	0.2	18.2±4.5	0.25	8.2	31%
Squat jump (cm)	33.1±4.1	0.13	25.7±3.9	0.15	7.4	22%

CV: covariance.

Source: Own elaboration.

Table 2 presents the partial correlations between HDL-C values and strength and power indicators. HDL-C values have low and negative correlations with values $p < 0.01$ and $p < 0.05$ for muscle strength and power tests. No sex differences were identified in the distribution of total cholesterol. A higher proportion of women with HDL-C values >40 mg/dL and triglycerides >200 mg/dL were observed ($p < 0.05$). No strong associations of HDL-C with the strength and power variables were found in the group of men or women.

Table 2. Correlation of HDL cholesterol with strength and power indicators.

HDL cholesterol	Men and women
Grip strength in right hand (kgf)	$r = -0.312^*$
Bench press (kg)	$r = -0.510^\dagger$
Grip strength in left hand (kgf)	$r = -0.363^*$
Half squat (kg)	$r = -0.381^*$
Peak anaerobic power upper limbs (watts)	$r = -0.333^*$
Peak anaerobic power upper limbs (watts/kg)	$r = -0.351^*$

* The correlation is significant at 0.05 (bilateral).

† The correlation is significant at 0.01 (bilateral).

Source: Own elaboration.

Discussion

The present study provides elements to discuss the relationship between different manifestations of MS and plasma values of triglycerides, total cholesterol, LDL-C and HDL-C. Evidence on this regard is scarce and contradictory, and the results of certain studies are difficult to interpret due to some weaknesses and the inadequate control of factors that may influence lipid metabolism and statistical power. (24) This statement is confirmed in the review carried out by Durstine *et al.* (25) and in the meta-analyses of Kelley *et al.* (26) and Zhang *et al.* (27), who analyzed workout plans in different populations and found few interventions that used MS training as a physical activity strategy. Most studies used low intensity aerobic work.

Blood levels of HDL-C are 4-24 mg/dL, but they are higher in subjects who perform physically demanding work or perform resistance exercise on a regular basis, compared to less active people. The relative differences of HDL-C between these two groups range between 9% and 59%. (25,28) The same pattern is observed with TC and LDL-C. (26) However, this situation is not the same when MS is the physical activity performed.

Zhang *et al.* (27) observed, in a population of young adults, a low to moderate negative correlation between the different manifestations of strength and HDL-C, as well as a lack of correlations with triglycerides, TC and LDL-C. These results agree with the results of this study and the reports of most of the previous works.

Jurca *et al.* (14) identified a moderate inverse association between the prevalence of metabolic syndrome and MS ($\beta = -0.37$, $p < 0.0001$), which decreases ($\beta = -0.08$, $p < 0.01$) when it is adjusted with cardiorespiratory fitness. (29)

Wijndaele *et al.* (13) reported low inverse correlations in men and moderate correlations in women ($p < 0.05$) between MS and triglycerides, cholesterol and HDL-C.

Artero *et al.* (30) identified moderate inverse correlations between grip strength and triglycerides ($p < 0.01$), and HDL-C and cholesterol ($p < 0.001$).

Steene-Johannessen *et al.* (31) conducted a study to identify the independence between MS and cardiorespiratory fitness for the risk of metabolic disease. The researchers evaluated standing long jump-explosive power, grip strength, abdominal strength and cardiorespiratory capacity directly using the peak oxygen uptake. The results indicate a negative association of MS and metabolic risk ($\beta = -0.337$, $p < 0.001$), and a weak negative correlation of grip strength with triglycerides and positive with HDL-C.

Regarding the methodology, the most commonly used tests in previous studies for the assessment of maximum strength and peak power were hand dynamometry and long jump. For this study, dynamic strength tests were used based on the 1RM methodology, such as bench press or squats, and explosive tests such as the Wingate test, CMJ or SJ, which could make the results between the studies little comparable. For example, in this study, the best correlation of HDL-C was obtained with bench press and not with hand dynamometer, which had similar correlation values to power in upper limbs. This could indicate that the level of correlation between HDL-C and MS depends on the type of test used to assess the maximum strength.

Fernández *et al.* (32), in a study on the correlation between the different manifestations of strength, observed that although the musculature of the upper or lower limbs is under evaluation, there is great variability in the correlation between the tests. This difference could be explained by the contribution of the muscle groups involved in the movements to each of the tests, which may not be the same. This would indicate that the information provided in each of the tests is specific to the muscle group and movement, and cannot be

extrapolated to another muscle group or movement. This difference can also be attributed to the neuromuscular implications of the different types of muscle actions involved in each test.

However, the reason why there are no positive correlations between high levels of strength and plasma levels of HDL-C, as is the case of cardiorespiratory capacity, remains unresolved. In this regard, Prabhakaran *et al.* (24) carried out an MS training for 14 weeks, with two sets of eight repetitions and bench press, leg extension and rowing workouts performed at 85% of 1RM. The results of the program led to significant reductions of cholesterol by 9% ($p < 0.05$) and LDL-C by 14.3% ($p < 0.05$) in the TC/HDL ratio. Triglycerides and HDL-C did not report significant changes.

Likewise, Kokkinos *et al.* (33) carried out a strength-training program in 37 subjects with three types of exercises. The results indicated that none of the proposed workout plans generated significant changes in plasma concentrations of triglycerides, total cholesterol, HDL-C and HDL2-C. Later, the same authors carried out another study with trained subjects performing strength work at 70% of 1RM and did not observe significant changes in plasma concentrations of triglycerides, TC, HDL-C or LDL-C. (34)

In the study by Manning *et al.* (35), 16 sedentary obese women trained 3 times a week for 12 weeks performing three sets of 6-8 repetitions at 60-70% of 1RM. No significant changes were observed in body weight, BMI or CT, HDL-C, LDL-C, TG, TC/HDL-c, apo AI or apo B-100 levels. (35)

Goldberg *et al.* (36) developed a free-weight workout plan for 16 weeks in a group of sedentary young men and women. Women presented a 9.5% reduction in cholesterol, 17.9% in LDL-C and 28.3% in triglycerides. The total cholesterol/HDL-C and LDL/HDL rates were reduced by 14.3% and 20.3%, respectively. In men, the LDL-C cholesterol was reduced by 16.2%, while the total cholesterol/HDL and LDL/HDL levels were reduced by 21.6% and 28.9%, respectively.

Only one study (37) that observed an increase in HDL-C and a decrease in TC and LDL-C in a similar way was identified; this study was conducted through strength training and resistance training. There, the results would indicate that, although participants presented significant gains in MS, strength training did not alter the plasma levels of TC, HDL-C, LDL-C, triglycerides, nor the TC/HDL-C ratio, as is the case of aerobic training programs. One possible explanation is the specificity of the adaptive response to physical activity, which, according to the level and type of stimulation, generates changes in hormonal profiles and inflammatory and immune parameters, as well as intrinsic changes in skeletal and cardiac muscles.

Although the mechanisms underlying the effect of physical activity on the lipid profile are not clear, one hypotheses raised several decades ago about the cause of the increase in plasma HDL-C due to the effects of aerobic exercise is related to the metabolic adaptations that the muscle undergoes due to an increase in enzymatic activity, which leads to an increase in the capacity to oxidize fatty acids in comparison with glycogen. This better use of plasma triglycerides occurs because of the increase in the lipoprotein lipase (LPL) activity that is located in the lumen of blood capillaries and hydrolyzes triglycerides, allowing the uptake of free fatty acids (FFA) and the increase in muscle angiogenesis. (38)

One of the possible mechanisms involved in HDL-C modulation is reverse cholesterol transport, which removes cholesterol from the circulation and distributes it to peripheral tissues and the liver. Two important events occur during aerobic exercise: 1) the increase of lecithin-cholesterol acyltransferase (LCAT), an enzyme responsible for ester transfer to HDL cholesterol, and 2) the reduction of cholesterol ester transfer protein (CETP), the enzyme responsible for the transfer of HDL cholesterol to the other lipoproteins. (39,40) This increases

the activity of lipoprotein lipase, although data on this issue are inconsistent. (41) All this increase in the degradation of lipoproteins rich in triglycerides, by means of apoproteins transfer could lead to an increase in the plasma concentration of HDL-C.

In physiological terms, aerobic physical activity performed for a long period creates resistance and keeps the cardiac pump elevated at a constant rate, which can boost cardiac function and increase HDL. (28)

This series of biochemical and physiological mechanisms is not activated in the same way during strength workout. Each type of workout poses a challenge for the organism's control systems to generate an adequate adaptive response. In this sense, and from a functional and metabolic point of view, muscle cells are not a homogeneous tissue and the cell signaling mechanisms and endocrine responses for the two types of workouts are different. Fatty acids are oxidized mostly in type I oxidative fibers that are primarily activated during low to moderate intensity physical activity. On the contrary, strength workout with moderate or high loads increases the activation of glycolytic fibers, therefore decreasing the participation of oxidative fibers. Similarly, cardiovascular function during SM workout is intermittent, since a high heart rate pump is not maintained at a constant rate, but for short periods (2-3 seconds) followed by recovery.

These statements are supported by the work of Aellen *et al.* (42), who studied the effects of anaerobic and aerobic training performed 4 times a week for 9 weeks on lipoprotein concentrations in a group of 62 healthy untrained men. 33 subjects performed physical activity at low intensity, 16 at an intensity above the anaerobic threshold (blood lactate concentration >4 mmol) and 12 served as controls. The caloric expenditure calculated from both training groups was similar; TC, HDL-C, the subtractions of HDL-C (HDL2-c, HDL3-c) and LDL-C were evaluated in the three groups. With anaerobic training, these variables turned to the opposite direction compared to aerobic training, which influenced the lipoprotein profile in the desired direction. The authors concluded that training above the anaerobic threshold has no effect on the blood lipoprotein profile and that changes are achieved with moderate training intensities. The study by Said *et al.* (43) obtained the same results. However, Lira *et al.* (44), in their study on acute response to different percentages of 1RM, observed that the concentration of HDL-C was significantly higher after training at 50% -1RM and at 75% -1RM compared to 110%.

Conclusions

This research identified a moderate to low inverse correlation between HDL-C and the different manifestations of MS. However, these results are preliminary and should be observed with caution due to several study limitations that may have generated a bias. Therefore, it is suggested to carry out studies with a larger sample size, but with the same participation of men and women to verify if similar responses are presented between them.

Along the same lines, a nutritional analysis should be included to quantify the energy expenditure of each subject, as well as the quantification of the level of physical activity and the determination of the ethnic group of each participant. With regard to women, it is important to know whether they are menstruating during the collection of the samples and which phase they are in due to the effect it has on the fluctuations of the lipid profile.

Due to the lack of evidence and the contradictory results of different works carried out on the subject, this study contributes to the consolidation of the evidence of the relationship between MS and metabolic risk markers and also indicates the need to establish this relationship from broader valuations of MS.

Conflicts of interest

None stated by the authors.

Funding

This study was financed by the Universidad Pedagógica Nacional.

Acknowledgements

To the group of young people who participated in the study and to the Biology Laboratory of Universidad Pedagógica Nacional for the analysis of the samples.

References

1. Pedersen BK, Febbraio MA. Muscle as an endocrine organ: focus on muscle-derived interleukin-6. *Physiol Rev*. 2008;88(4):1379-406. <http://doi.org/ffdgx4>.
2. Pratesi A, Tarantini F, Di Bari M. Skeletal muscle: an endocrine organ. *Clin Cases Miner Bone Metab*. 2013;10(1):11-4. <http://doi.org/cp87>.
3. Febbraio MA, Pedersen BK. Contraction-induced myokine production and release: is skeletal muscle an endocrine organ? *Exerc Sport Sci Rev*. 2005;33(3):114-9. <http://doi.org/c9gg2j>.
4. Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, Lee IM, *et al.* American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc*. 2011;43(7):1334-59. <http://doi.org/c6hwt6>.
5. Cheung CL, Nguyen US, Au E, Tan KC, Kung AW. Association of handgrip strength with chronic diseases and multimorbidity: a cross-sectional study. *Age (Dordr)*. 2013;35(3):929-41. <http://doi.org/gb9zbp>.
6. Volaklis KA, Halle M, Thorand B, Peters A, Ladwig KH, Schulz H, *et al.* Handgrip strength is inversely and independently associated with multimorbidity among older women: Results from the KORA-Age study. *Eur J Intern Med*. 2016;31:35-40. <http://doi.org/f8px4v>.
7. Artero EG, Lee DC, Ruiz JR, Sui X, Ortega FB, Church TS, *et al.* A prospective study of muscular strength and all-cause mortality in men with hypertension. *J Am Coll Cardiol*. 2011;57(18):1831-7. <http://doi.org/bw9x9d>.
8. Volaklis KA, Halle M, Meisinger C. Muscular strength as a strong predictor of mortality: A narrative review. *Eur J Intern Med*. 2015;26(5):303-10. <http://doi.org/f3gqvs>.
9. Cooper R, Kuh D, Hardy R, Mortality Review Group, FALCon and HAL-Cyon Study Teams. Objectively measured physical capability levels and mortality: systematic review and meta-analysis. *BMJ*. 2010;341:c4467. <http://doi.org/dzn7wm>.
10. Timpka S, Petersson IF, Zhou C, Englund M. Muscle strength in adolescent men and risk of cardiovascular disease events and mortality in middle age: a prospective cohort study. *BMC Medicine*. 2014;12:62. <http://doi.org/f244db>.
11. Ortega FB, Silventoinen K, Tynelius P, Rasmussen F. Muscular strength in male adolescents and premature death: cohort study of one million participants. *BMJ*. 2012;345:e7279. <http://doi.org/f249b9>.
12. FitzGerald SJ, Barlow CE, Kampert JB, Morrow JR, Jackson AW, Blair SN. Muscular Fitness and All-Cause Mortality: Prospective Observations. *J Phys Act Health*. 2004;1(1):7-18. <http://doi.org/cp88>.
13. Wijndaele K, Duvigneaud N, Matton L, Duquet W, Thomis M, Beunen G, *et al.* Muscular strength, aerobic fitness, and metabolic syndrome risk in Flemish adults. *Med Sci Sports Exerc*. 2007;39(2):233-40. <http://doi.org/bztvx2>.

14. Jurca R, Lamonte MJ, Church TS, Earnest CP, Fitzgerald SJ, Barlow CE, *et al.* Associations of muscle strength and fitness with metabolic syndrome in men. *Med Sci Sports Exerc.* 2004;36(8):1301-7. <http://doi.org/df3dvs>.
15. Aoyama T, Asaka M, Ishijima T, Kawano H, Cao ZB, Sakamoto S, *et al.* Association between muscular strength and metabolic risk in Japanese women, but not in men. *J Physiol Anthropol.* 2011;30(4):133-9. <http://doi.org/d5f8h2>.
16. Ramírez-Vélez R, Correa-Bautista JE, Lobelo F, Izquierdo M, Alonso-Martínez A, Rodríguez-Rodríguez F, *et al.* High muscular fitness has a powerful protective cardiometabolic effect in adults: influence of weight status. *BMC Public Health.* 2016;16(1):1012. <http://doi.org/cp89>.
17. García-Hermoso A, Carrillo HA, González-Ruiz K, Vivas A, Triana-Reina HR, Martínez-Torres J, *et al.* Fatness mediates the influence of muscular fitness on metabolic syndrome in Colombian collegiate students. *PLoS One.* 2017;12(3):e0173932. <http://doi.org/f9t2hj>.
18. Ramírez-Vélez R, Tordecilla-Sanders A, Correa-Bautista JE, Peterson MD, García-Hermoso A. Handgrip Strength and Ideal Cardiovascular Health among Colombian Children and Adolescents. *J Pediatr.* 2016;179:82-9.e1. <http://doi.org/f9hd3m>.
19. Asociación Médica Mundial. Declaración de Helsinki de la Asociación Médica Mundial. Principios éticos para las investigaciones médicas en seres humanos. Fortaleza: 64.ª Asamblea General de la AMM; 2013 [cited 2017 Dec 9]. Available from: <https://goo.gl/hvf711>.
20. Colombia. Ministerio de Salud. Resolución 8430 de 1993 (octubre 4): Por la cual se establecen las normas científicas, técnicas y administrativas para la investigación en salud. Bogotá D.C.; octubre 4 de 1993 [cited 2017 Dec 9]. Available from: <https://goo.gl/agV1mY>.
21. Naclerio FJ, Colado JC, Rhea MR, Bunker D, Triplett NT. The influence of strength and power on muscle endurance test performance. *J Strength Cond Res.* 2009;23(5):1482-8. <http://doi.org/ccqeph>.
22. de Freitas M, Andrade-Paz G, Souza J, Miranda H. Parâmetros de rendimento de força miotônica e adaptação de diferentes sequências de exercícios durante séries emparelhadas agonista-antagonista. *Apunts Med Esport.* 2015;50(187):103-10.
23. Colantonio E, Vilela-Barros R, Peduti-dal Molin-Kiss MA. Oxygen uptake during Wingate tests for arms and legs in swimmers and water polo players. *Rev Bras Med Esporte.* 2003;9(3):141-4. <http://doi.org/bjknkn>.
24. Prabhakaran B, Dowling EA, Branch JD, Swain DP, Leutholtz BC. Effect of 14 weeks of resistance training on lipid profile and body fat percentage in premenopausal women. *Br J Sports Med.* 1999;33(3):190-5. <http://doi.org/brp59s>.
25. Durstine JL, Grandjean PW, Davis PG, Ferguson MA, Alderson NL, DuBose KD. Blood lipid and lipoprotein adaptations to exercise: a quantitative analysis. *Sports Med.* 2001;31(15):1033-62. <http://doi.org/fcvq87>.
26. Kelley GA, Kelley KS, Tran ZV. Aerobic exercise and lipids and lipoproteins in women: a meta-analysis of randomized controlled trials. *J Womens Health.* 2004;13(10):1148-64. <http://doi.org/brtwcx>.
27. Zhang Y, Xu L, Zhang X, Yao Y, Sun Y, Qi L. Effects of different durations of aerobic exercise intervention on the cardiovascular health in untrained women: a meta-analysis and meta-regression. *J Sports Med Phys Fitness.* 2017;57(5):595-603. <http://doi.org/cp9c>.
28. Mann S, Beedie C, Jiménez A. differential effects of aerobic exercise, resistance training and combined exercise modalities on cholesterol and the lipid profile: review, synthesis and recommendations. *Sports Med.* 2014;44(2):211-21. <http://doi.org/f5r5nc>.
29. Jurca R, Lamonte MJ, Barlow CE, Kampert JB, Church TS, Blair SN. Association of muscular strength with incidence of metabolic syndrome in men. *Med Sci Sports Exerc.* 2005;37(11):1849-55. <http://doi.org/ch52rj>.
30. Artero EG, Ruiz JR, Ortega FB, España-Romero V, Vicente-Rodríguez G, Molnar D, *et al.* Muscular and cardiorespiratory fitness are independently associated with metabolic risk in adolescents: the HELENA study. *Pediatr Diabetes.* 2011;12(8):704-12. <http://doi.org/dh267b>.
31. Steene-Johannessen J, Anderssen SA, Kolle E, Andersen LB. Low muscle fitness is associated with metabolic risk in youth. *Med Sci Sports Exerc.* 2009;41(7):1361-7. <http://doi.org/cfstxc>.
32. Fernández-Ortega JA, Hoyos-Cuartas LA. Relaciones entre diversas manifestaciones de fuerza en diferentes grupos musculares en adultos jóvenes. *Revista U.D.C.A Actualidad & Divulgación Científica.* 2017;20(1):33-42.
33. Kokkinos PF, Hurley BF, Vaccaro P, Patterson JC, Gardner LB, Ostrove SM, *et al.* Effects of low- and high-repetition resistive training on lipoprotein-lipid profiles. *Med Sci Sports Exerc.* 1988;20(1):50-4. <http://doi.org/bewfrs>.
34. Kokkinos PF, Hurley BF, Smutok MA, Farmer C, Reece C, Shulman R, *et al.* Strength training does not improve lipoprotein-lipid profiles in men at risk for CHD. *Med Sci Sports Exerc.* 1991;23(10):1134-9. <http://doi.org/fwc7k6>.
35. Manning JM, Dooly-Manning CR, White K, Kampa I, Silas S, Kesselhaut M, *et al.* Effects of a resistive training program on lipoprotein-lipid levels in obese women. *Med Sci Sports Exerc.* 1991;23(11):1222-6. <http://doi.org/c9s5xt>.
36. Goldberg L, Elliot DL, Schutz RW, Kloster FE. Changes in lipid and lipoprotein levels after weight training. *JAMA.* 1984;252(4):504-6. <http://doi.org/czj24f>.
37. Fahlman MM, Boardley D, Lambert CP, Flynn MG. Effects of endurance training and resistance training on plasma lipoprotein profiles in elderly women. *J Gerontol A Biol Sci Med Sci.* 2002;57(2):B54-60. <http://doi.org/bnw3zs>.
38. Earnest CP, Artero EG, Sui X, Lee DC, Church TS, Blair SN. Maximal estimated cardiorespiratory fitness, cardiometabolic risk factors, and metabolic syndrome in the Aerobics Center Longitudinal Study. *Mayo Clin Proc.* 2013;88(3):259-70. <http://doi.org/f2xh93>.
39. Calabresi L, Franceschini G. Lecithin:cholesterol acyltransferase, high-density lipoproteins, and atheroprotection in humans. *Trends Cardiovasc Med.* 2010;20(2):50-3. <http://doi.org/czbwnh>.
40. Riedl I, Yoshioka M, Nishida Y, Tobina T, Paradis R, Shono N, *et al.* Regulation of skeletal muscle transcriptome in elderly men after 6 weeks of endurance training at lactate threshold intensity. *Exp Gerontol.* 2010;45(11):896-903. <http://doi.org/fmde6w>.
41. Harrison M, Moyna NM, Zderic TW, O'Gorman DJ, McCaffrey N, Carson BP, *et al.* Lipoprotein particle distribution and skeletal muscle lipoprotein lipase activity after acute exercise. *Lipids Health Dis.* 2012;11:64. <http://doi.org/cp9f>.
42. Aellen R, Hollmann W, Boutellier U. Effects of aerobic and anaerobic training on plasma lipoproteins. *Int J Sports Med.* 1993;14(7):396-400. <http://doi.org/c69jdv>.
43. Said M, Lamy N, Olfa N, Hamda M. Effects of high-impact aerobics vs. low-impact aerobics and strength training in overweight and obese women. *J Sports Med Phys Fitness.* 2017;57(3):278-88. <http://doi.org/cp9g>.
44. Lira FS, Yamashita AS, Uchida MC, Zanchi NE, Gualano B, Martins E Jr., *et al.* Low and moderate, rather than high intensity strength exercise induces benefit regarding plasma lipid profile. *Diabetol Metab Syndr.* 2010;2:31. <http://doi.org/cm6b3f>.