

## ORIGINAL RESEARCH

DOI: <http://dx.doi.org/10.15446/revfacmed.v66n4.65199>

## Use of sexual health services at hospitals by cleaning workers in Bogotá D.C. 2016

*Uso de los servicios de salud sexual por parte de trabajadoras de servicios generales en hospitales en Bogotá D.C. 2016*

Received: 23/05/2017. Accepted: 10/25/2017.

Diana Marcela Hernández-Pérez<sup>1</sup> • María Natalia Moreno-Ruiz<sup>1</sup> • Anderson Rocha-Buelvas<sup>2</sup> • Arsenio Hidalgo-Troya<sup>3</sup>

<sup>1</sup> Fundación Universitaria del Área Andina - Faculty of Health Sciences - Master's Degree in Public Health and Social Development - Bogotá D.C. - Colombia.

<sup>2</sup> Universidad de Nariño - Faculty of Health Sciences - Technology Program in Health Promotion - Pasto - Colombia.

<sup>3</sup> Universidad de Nariño - Faculty of Exact and Natural Sciences - Department of Mathematics and Statistics - Pasto - Colombia.

Corresponding author: Anderson Rocha-Buelvas. Technology Program in Health Promotion, Faculty of Health Sciences Universidad de Nariño. Ciudad Universitaria Torobajo, calle 18 carrera 50, building 7, floor 2. Telephone Number: +57 7314552. Pasto. Colombia. Email: rochabuelvas@gmail.com.

### | Abstract |

**Introduction:** Poverty and social inequalities together with sexually transmitted diseases have a negative impact on women's health, which is considered to be a public health problem.

**Objective:** To analyze barriers to accessing sexual and reproductive health services in cleaning workers.

**Materials and methods:** A survey was administered to a sample of 37 female cleaning workers at a hospital in Bogotá D.C. A bivariate analysis was performed with chi-square test, as well as a multivariate analysis with binomial logistic regression.

**Results:** Need factors showed greater association with non-use of sexual health services. All married women had accessed the service over the past 12 months, but there were 5.9 less possibilities of using sexual and reproductive health services when there was no awareness about risk behaviors of sexually transmitted diseases.

**Conclusion:** The determining factor for the utilization of sexual health services is the health care need factor. Variables such as perception of risk behaviors and appropriateness of health care significantly influence the use of the service.

**Keywords:** Housekeeping Hospital Service; Sexually Transmitted Diseases; Accessibility of Health Services; Sexual Health; Gender Identity; Women (MeSH).

### | Resumen |

**Introducción.** La pobreza y las desigualdades sociales, junto a infecciones de transmisión sexual, tienen un impacto negativo en la salud de la mujer, lo que se considera un problema de salud pública.

**Objetivo.** Analizar las barreras de acceso para la utilización de servicios de salud sexual de las mujeres trabajadoras en servicios generales.

**Materiales y métodos.** Se aplicó una encuesta a una muestra de 37 mujeres trabajadoras en servicios generales en un hospital de Bogotá D.C. Se realizó análisis bivariado con prueba chi cuadrado y multivariado con regresión logística binomial.

**Resultados.** Los factores de necesidad tuvieron mayor asociación con la no utilización de los servicios de salud sexual. Todas las mujeres casadas accedieron al servicio en los últimos 12 meses y existe 5.9 menos posibilidades de utilizar los servicios de salud sexual y reproductiva si se desconocen las conductas de riesgo de las infecciones de transmisión sexual.

**Conclusión.** El principal factor determinante para la utilización de los servicios de salud sexual es el factor de necesidad de atención en salud, cuyas variables, como la percepción de conductas de riesgo y la pertinencia de la atención en salud, influyen significativamente en la utilización del servicio.

**Palabras clave:** Servicio de limpieza en hospital; Enfermedades de transmisión sexual; Accesibilidad a los servicios de salud; Salud sexual; Identidad de Género; Mujeres (DeCS).

Hernández-Pérez DM, Moreno-Ruiz MN, Rocha-Buelvas A, Hidalgo-Troya A. Use of sexual health services at hospitals by cleaning workers in Bogotá D.C. 2016. Rev. Fac. Med. 2018;66(4):617-22. English. doi: <http://dx.doi.org/10.15446/revfacmed.v66n4.65199>.

Hernández-Pérez DM, Moreno-Ruiz MN, Rocha-Buelvas A, Hidalgo-Troya A. [Uso de los servicios de salud sexual por parte de trabajadoras de servicios generales en hospitales en Bogotá D.C. 2016]. Rev. Fac. Med. 2018;66(4):617-22. English. doi: <http://dx.doi.org/10.15446/revfacmed.v66n4.65199>.

## Introduction

In recent history, women have legitimized sexual rights as human rights, promoting gender equity as a determining factor for sexual and reproductive health research and public policy planning. (1,2) The Colombian health care model is strongly influenced by the competition within the supply and demand of health services; moreover, market laws compromise the equality and universality of these services. (3)

According to a systematic review of studies on access to health services in Colombia, access has not increased as there are barriers associated with insurance, economic income, educational attainment and geographical, organizational and quality-service accessibility. (4) In this regard, the efforts of national and international health organizations to fight the occurrence of sexually transmitted diseases (STDs) specifically have not prevented persistent reproductive and sexual health risks. (5)

UNAIDS estimates show that although the incidence of HIV has decreased in recent decades, it is still a public health problem. (6,7) Both women and men living with HIV and receiving treatment have increased significantly (8); it is also a matter of concern that around 1.4 million maternal infections and 520 000 congenital infections caused by syphilis have been reported worldwide, including 304 000 perinatal deaths from 2009 to 2012 among working populations and disadvantaged social classes of Latin America. (9,10) Furthermore, the people most prone to contract these diseases are the most disadvantaged human groups. (10,11)

It should be noted that barriers to accessing sexual health services are determined by internal factors that depend on each person's perception and decision while looking for a health service, as well as external factors that can derive from administrative, legal and structural difficulties, lack of awareness about the health system, socio-economic situation, culture, communication and education. (12-14) Identifying the barriers to accessing health services largely contributes to monitor inequalities and generate proposals to mitigate impacts on their health condition. (15)

The utilization of health services comes from the interaction of three main factors: 1) predisposing factors: they are inherent to the patient, do not depend on the pathology and evaluate the perception of health and disease related to socioeconomic and demographic variables (age, sex, occupation, ethnicity, etc.) (16); 2) mediating or capacity factors: they can facilitate or limit access to services and are subject to changes in health policies, and 3) need factors: they include aspects related to the health-disease process and address the need in two ways, namely, the patient's perspective and the evaluation by the health staff. (17-19)

Therefore, the objective of this study was to analyze the barriers to accessing sexual and reproductive health services by female cleaning workers at a hospital center in Bogotá D.C. in 2016; these women are one of the most marginalized human groups given their low educational attainment, low income and lack of knowledge about their health rights.

## Materials and methods

The study was developed under an observational, descriptive cross-sectional design. A survey was conducted based on the behavioral model of health services proposed by Aday & Andersen and adapted for sexual health. (17) Need factor variables were: knowledge of sexual health services, sexual and reproductive rights, STDs, contraception and quality of health care received. The predisposing

factor variables were: age, marital status, ethnicity, educational attainment, socio-economic level and social support. Finally, capacity factor variables were: family income, commuting time and location of the Healthcare Service Provider (IPS by its acronym in Spanish), out-of-pocket expenses, among others. The query "sexual health services over the past 12 months" was established for the exposure variable.

Similarly, a pilot study (10 surveys) was conducted to check whether the questionnaire was understood, as well as the coherence and relevance of questions concerning the topic under evaluation.

The inclusion criteria used were: women who were not pregnant, female hospital cleaning workers hired by the JDR company (contracting party) at Fundación Hospital San Carlos, an institution where 37 women work in cleaning services. It was relevant to conduct the study in this vulnerable population because there was no prior record.

Frequency measurements were performed to determine the sample distribution based on age, marital status, ethnicity, employment status, educational attainment and socio-economic level. Predisposing, capacity and need factors concerning the use of sexual and reproductive health services were identified through bivariate and multivariate analyses; the goal was to develop an explanatory model based on the individual determining factors of the Aday & Andersen model.

The chi square test was performed in the bivariate analysis, where the significant statistical evidence were values of  $p < 0.10$ . Effect confusion and modification were assessed through a multivariate analysis by selecting the variables with values  $p < 0.25$  in the bivariate analysis; this analysis was performed using the logistic regression model. The Wald test was used to establish statistical significance and measure the strength of association with the odds ratio (OR) values. The process was executed in the SPSS® software version 14.

The study was approved by the Research Committee of the Master's Degree in Public Health and Social Development of Fundación Universitaria del Área Andina and all the participants signed and informed consent. On the one hand, the research followed the guidelines of the Declaration of Helsinki (20) on the participation of the subjects and the researcher's confidentiality duties both at the beginning of and during the study; on the other, the research followed the guidelines established in the Resolution 8430 of the Colombian Ministry of Health (21), which sets forth the scientific, technical and administrative standards for health research.

## Results

Regarding predisposing factors, 18.9% of the sample had not completed secondary studies, and only 2.7% had completed higher education studies, thus confirming that most of them have a low educational attainment; no participant was identified within an ethnic group (Table 1). 35.1% of the respondent women were 21 to 40 years old and the bivariate analysis showed that age influenced the utilization of sexual health services before the 12 months; for example, 21 to 30-year old women used the service the most (66.7%), while 41 to 50-year old women used it less (12.5%). The marital status variable showed that 100% of married women attended the appointments, while only 14.3% of unmarried women attended the appointments (based on the p-value of the chi square test); 40.5% of the interviewed population lived in cohabitation. Finally, the multivariate analysis did not find any variables of predisposing factors that were closely related to the utilization of health services.

**Table 1.** Determining factors for the utilization of sexual and reproductive health services by cleaning workers at a hospital in Bogotá D.C.. 2015.

Factor	Variables		n	%	P
Predisposing factors	Age (years)	≤20	6	16.2%	0.094
		21-30	6	16.2%	
		31-40	13	35.1%	
		41-50	8	21.6%	
		>50	4	10.8%	
	Educational attainment	Illiterate	0	0.0%	0.206
		Incomplete primary	3	8.1%	
		Completed primary	4	10.8%	
		Incomplete secondary	7	18.9%	
		Completed secondary	22	59.5%	
		Higher education	1	2.7%	
	Marital Status	Single	14	37.8%	0.087
		Married	2	5.4%	
		Cohabitation	15	40.5%	
		Widow	1	2.7%	
		Divorced	5	13.5%	
Capacity factors	Socio-economic level	1	16	43.2%	0.625
		2	20	54.1%	
		3	1	3.7&	
		4	0	0.0%	
	Income	1 CLMW	37	100.0%	0.973
		From 1 to 2 CLMW	0	0.0%	
		More than 2 CLMW	0	0.0%	
	Additional support	Yes	6	16.2%	0.633
No		31	83.8%		
Need factors	Knowledge about sexual and reproductive rights	Yes	21	26.8%	0.143
		No	16	43.2%	
	Condom use	Yes	10	27.0%	0.539
		No	14	37.8%	
		Sometimes	13	35.1%	
	Partner interested in using a condom	Yes	12	32.4%	0.123
		No	19	51.4%	
		Sometimes	6	16.2%	
	Appointment allocation time	Low	37	100.0%	0.152
		Normal	0	0.0%	
	Discrimination	Yes	4	10.8%	0.230
		No	33	89.2%	
	General quality of care	Low	14	37.8%	0.137
		Normal	22	59.5%	

CLMW: Current legal minimum wage

Source: Own elaboration.

With respect to capacity factors, 54.1% of the respondents belonged to the socio-economic level 2, 100% had an income of one current legal minimum wage, 59% had 1 to 2 children, 83% did not have any additional financial support and 89.2% considered that their income was insufficient to cover their monthly expenses. Moreover, 48.6% mentioned that the IPS was close to their workplace. It is worth

mentioning that 67.6% stated that they have economic difficulties to use health services and nobody mentioned any difficulties related to work for such access (Table 1). The bivariate analysis showed that family income is related to lack of attendance to health services in 75.7% of the sample and that no variables in the multivariate analysis were strongly associated with the use of services.

Need factors showed that 43.2% is not aware of their sexual and reproductive rights and 64.9% is not aware of the sexual health services offered by their health promotion organization. Concerning knowledge about STDs, it was observed that 91% know what STDs are, since most of them have received information directly from health professionals at their workplace. It was evident that the respondents associate infidelity and having multiple sexual partners to a risk behavior leading to HIV infection (45.9%), as well as not using condoms (67%).

Concerning quality of health care, 100% of the participants gave a low rating to the opportunity they have to get appointments and the explanations given by health care professionals, 10.8% felt discriminated and 13.5% felt judged during the appointment; in short, the quality of health care in general had a low rating from 37.8% of the respondents.

Moreover, only 27% use condoms on a regular basis during their sexual relations, 24.3% did not consider it necessary and 21.6% did not like it. 54% had taken an HIV test, the majority (27%) had been tested more than 3 years before the execution of the study and did not deem it necessary to repeat it. 89% had undergone a cytology test; however, 78.4% had not taken this test in the last year. Taking into account the time of the last appointment for sexual health services, it was found that 37.8% had never accessed these services and that 37.8% had not used them in the last 12 months (Table 1).

Multivariate analysis showed that the need factors related to knowledge about STDs, purchase of condoms and problems related to access to sexual health services have a statistically significant correlation with the utilization of sexual and reproductive health services by women. That is to say, those who do not have proper knowledge about HIV/AIDS risk behaviors use sexual health services 5.9 times less than women with such knowledge. Furthermore, purchasing condoms in places such as pharmacies and supermarkets has an impact on access to the services, since those buyers access the services 0.8 times less than those who get condoms at their IPS. Finally, when there are issues to make sexual health appointments, women access 2.1 times less than those who do not have any difficulty in doing so (Table 2).

**Table 2.** Logistic regression analysis. Explanatory model of self-perception of sexual health and its correlation with quality of life among female cleaning workers.

Factor	Variable	OR *	CI95%	p
Need factors	HIV risk behavior	5.9	0.764-46.4	0.089
	Place of condom purchase	0.76	0.02-2.40	0.143
	Issues to make an appointment	2.16	0.038-121.5	0.092
	Adjustment quality	Nagelkerke R <sup>2</sup>	0.467	
		Hosmer-Lemeshow test	0.89	

\* OR adjusted by variables of knowledge about HIV transmission behaviors, issues to request for services, and place where condoms are purchased. Source: Own elaboration.

## Discussion

The state of the art of research on access to health services in Colombia shows that Colombians experience mainly administrative, economic and cultural barriers, where the perception of health condition and the

need for care together with a low educational attainment and the lack of information on matters related to the right to health play a key role in the utilization of health services. (22,23) This study demonstrated that the perception of risk in health care is related to the utilization of services and is influenced by the social and economic context of the individual, as well as by variables such as age, marital status, educational attainment and ethnicity.

The above has been shown in research on access to health services, mostly among women who suffer a high-cost disease such as cervical cancer or breast cancer. (24-31) It is also essential to recognize the role of social support in the utilization of health care, since it was demonstrated that married women accessed services more frequently than unmarried women or women in cohabitation. In contrast to these results, a research on taking Pap test stated that one of the main factors that limits access to said test is the pressure that women are subjected to by their partner. (32)

Geographic and travelling access barriers have also been identified, particularly in rural areas (24); this demonstrates that the use of health services in cities is not linked to the travelling time, since the servicing IPS are located close to the participants' residence or workplace. (12,33,34)

Out-of-pocket expenditure represents an economic burden that the most vulnerable populations are not able to afford. (35) In the case of Colombia, this translates into not guaranteeing full access to services due to factors such as copayment, travelling and treatments that are not covered by the Mandatory Health Plan (POS for its acronym in Spanish). The bivariate analysis of this research showed that capacity factors are related to and represent barriers to accessing sexual health services; however, they are not the most representative factors.

STDs have been studied in depth worldwide, demonstrating that they are preventable diseases transmitted primarily by sexual contact (36); nevertheless, STDs and their negative impacts on health, which may include chronicity, infertility, cancer or even death, have not been controlled. (37,38) Consequently, STDs continue to be a priority for public health and a change that focuses on preventing contagion and providing timely treatment (39-41) that encompasses the perception of risk is required, since there is a higher probability of non-access to health services when risk behaviors related to STD transmission are not known, condoms are purchased in commercial places and issues arise while making an appointment.

Furthermore, some research studies have found a correlation between the social stigma generated by HIV in communities and the access to and effective use of health services for the diagnosis and treatment of the disease. (42) Also, gender status is considered to be one of the factors that lead to health inequalities, since it has been defined as a social construction that takes the female and male concepts beyond biological characteristics, relates to all economic, social and private aspects of life of individuals, and determines features and roles depending on sex or on how society sees the subject. (1,43-46)

Access to health services has been studied throughout the world; however, the lack of a specific characterization of populations is evident; although population groups with similarities are addressed, it is not possible to extrapolate their special features. (47) This study allows advancing in knowledge about access barriers and determining factors that influence access to sexual health services by a group of women that has not been extensively studied nor intervened in a timely manner by the health system, even less when they are immigrants, considering essential variables such as predisposing factors, capacity factors and need factors to evaluate the utilization of health services, which are modified over time. Thus, it is imperative to generate continuous monitoring to determine the actual needs of the population.

Identifying access barriers contributes largely to monitor inequalities and generate proposals to mitigate impacts on health condition. (48) Therefore, one of the limitations of this study lies on the fact that the results obtained show a correlation between the perceived need to attend health services, the knowledge about sexual health rights and the risk behaviors of STD transmission; these are, however, not enough to demonstrate the cultural, social, political and economic reasons influencing the use of health services. Therefore, it is advisable to perform a qualitative study to identify and go in depth regarding other determining factors related to women's sexual health.

## Conclusion

The determining factor identified for the utilization of sexual health services by female cleaning workers at a hospital in Bogotá D.C. is health care need; associated variables such as perception of risk behaviors and appropriateness of health care significantly influence the use of the service.

## Conflicts of interest

None stated by the authors.

## Funding

None stated by the authors.

## Acknowledgments

We would like to thank the Master's Degree in Public Health and Social Development of the Fundación Universitaria del Área Andina.

## References

1. Alfonso-Rodríguez AC. Salud sexual y reproductiva desde la mirada de las mujeres. *Rev Cuba Salud Publica*. 2006 [cited 2018 Jun 22];32(1):1-15. Available from: <https://goo.gl/x1JVhN>.
2. Castro-Jiménez MA, Vera-Cala LM, Posso-Valencia HJ. Epidemiología del cáncer de cuello uterino: estado del arte. *Rev Colomb Obstet y Ginecol*. 2006 [cited 2018 Jun 22];57(1):182-9. Available from: <https://goo.gl/NExmJP>.
3. Bejarano-Daza JE, Hernández-Losada DF. Fallas del mercado de salud colombiano. *Rev. Fac. Med.* 2017;65(1):107-13. <http://doi.org/ctdn>.
4. Vargas-Lorenzo I, Vázquez-Navarrete ML, Mogollón-Pérez AS. Acceso a la atención en salud en Colombia. *Rev. salud pública*. 2010;12(5):701-12. <http://doi.org/b3drq9>.
5. Colombia. Ministerio de Salud y Protección Social. Boletín epidemiológico, situación del VIH/Sida Colombia 2013. Bogotá D.C.: MinSalud; 2013.
6. GBD 2015 HIV Collaborators Estimates of global, regional, and national incidence, prevalence, and mortality of HIV, 1980-2015: the Global Burden of Disease Study 2015. *Lancet HIV*. 2016;3(8):e361-87. <http://doi.org/bm2p>.
7. Davies B, Turner KME, Frølund M, Ward H, May MT, Rasmussen S, et al. Risk of reproductive complications following chlamydia testing: a population-based retrospective cohort study in Denmark. *Lancet Infect Dis*. 2016;16(9):1057-64. <http://doi.org/f8xpbv>.
8. Álvarez-Castaño LS. Los determinantes sociales de la salud : más allá de los factores de riesgo. *Rev. Gerenc. Polit. Salud, Bogotá*. 2011;8(17):69-79.
9. Wijesooriya NS, Roach RW, Kamb ML, Turlapati P, Temmerman M, Broutet N, et al. Global burden of maternal and congenital syphilis in 2008 and 2012: a health systems modelling study. *Lancet Glob Health*. 2016;4(8):e525-33. <http://doi.org/f8x4pn>.
10. Bones-Rocha K, Muntaner C, González-Rodríguez MJ, Bernales-Baksai P, Vallebuona C, Borrell C, et al. Clase social, desigualdades en salud y conductas relacionadas con la salud de la población trabajadora en Chile. *Rev Panam Salud Publica*. 2013 [cited 2018 Jun 22];33(5):340-8. Available from: <https://goo.gl/fg1J2R>.
11. Pérez-Valbuena GJ, Silva-Ureña A. Una mirada a los gastos de bolsillo en salud para Colombia. Bogotá D.C.: Banco de la República, Documento de trabajo sobre economía regional No.218; 2015.
12. Garcés-Palacio IC, Rubio-León DC, Scarinci IC. Factores asociados con el tamizaje de cáncer de cuello uterino en mujeres de nivel socioeconómico medio y bajo en Bogotá, Colombia. *Rev Fac Nac Salud Pública*. 2012;30(1):7-16.
13. Díaz-Grajales C, Zapata-Bermúdez Y, Aristizábal-Grisales JC. Acceso a los servicios preventivos en los regímenes contributivo y subsidiado de salud en un barrio estrato dos de la ciudad de Cali. *Rev. Gerenc. Polit. Salud*. 2011;10(21):153-75.
14. Yao X, Dembe AE, Wickizer T, Lu B. Does time pressure create barriers for people to receive preventive health services? *Prev Med*. 2015;74:55-8. <http://doi.org/f69vp2>.
15. Andersen RM. Revisiting the behavioral model and access to medical care: does it matter? *J Health Soc Behav*. 1995;36(1):1-10. <http://doi.org/dhmtg>.
16. Arredondo A, Melendez V. Modelos explicativos sobre la utilización de servicios de salud: revisión y análisis. *Salud Publica de México*. 1992;34(1):36-49.
17. Aday LA, Andersen R. A Framework for the Study of Access to Medical Care. *Health Serv Res*. 1974;9(3):208-20.
18. Arrivillaga M, Borrero YE. Visión comprensiva y crítica de los modelos conceptuales sobre acceso a servicios de salud, 1970-2013. *Cad. Saúde Pública*. 2016;32(5):e00111415. <http://doi.org/crdb>.
19. Tovar-Cuevas LM, Arrivillaga-Quintero MA. Estado del arte de la investigación en acceso a los servicios de salud en Colombia, 2000-2013: revisión sistemática. *Rev. Gerenc. Polit. Salud*. 2014;13(27):12-26.
20. Asociación Médica Mundial. Declaración de Helsinki de la Asociación Médica Mundial. Principios éticos para las investigaciones médicas en seres humanos. Fortaleza: 64.ª Asamblea General de la AMM; 2013 [cited 2018 Aug 28]. Available from: <https://goo.gl/hvf711>.
21. Colombia. Ministerio de Salud. Resolución 8430 de 1993 (octubre 4): Por la cual se establecen las normas científicas, técnicas y administrativas para la investigación en salud. Bogotá D.C.; octubre 4 de 1993 [cited 2018 Aug 28]. Available from: <https://goo.gl/agV1mY>.
22. Hirmas-Adaury M, Poffald-Angulo L, Jasmen-Sepúlveda AM, Aguilera-Sancheza X, Delgado-Becerra I, Vega-Morales J. Barreras y facilitadores de acceso a la atención de salud: una revisión sistemática cualitativa. *Rev Panam Salud Publica*. 2013;33(3):223-9.
23. Mejía-Mejía A, Sánchez-Gandur AF, Tamayo-Ramírez JC. Equidad en el Acceso a Servicios de Salud en Antioquia, Colombia. *Rev. Salud Pública*. 2007;9(1):26-38.
24. Sánchez G, Laza C, Estupiñán C, Estupiñán L. Barreras de acceso a los servicios de salud: narrativas de mujeres con cáncer de mama en Colombia. *Rev. Fac. Nac. Salud Pública*. 2014;32(3):305-13.
25. Vargas J, Molina G. Acceso a los servicios de salud en seis ciudades de Colombia: limitaciones y consecuencias. *Rev Fac Nac Salud Pública*. 2009;27(2):121-30.
26. Lucumí-Cuesta DI, Gomez-Gutiérrez LF. Accesibilidad a los servicios de salud en la práctica de citología reciente de cuello uterino en una zona urbana de Colombia. *Rev. Esp. Salud Publica*. 2004;78(3):367-77.
27. Garcés-Palacio IC, Rubio-León DC, Ramos-Jaraba SM. Barreras y facilitadores del sistema de salud relacionadas con el seguimiento de anomalías citológicas, Medellín-Colombia. *Rev. Gerenc Polit Salud*. 2014;13(27):200-11. <http://doi.org/cs67>.
28. Triviño Z, Stiepvovich J, Merino JM. Factores predictores de conductas promotoras de salud en mujeres peri- post-menopáusicas de Cali , Colombia. *Colomb Med*. 2007;38(4):395-407.



29. **Rodríguez-González D, Pérez-Piñero J, Sarduy-Nápoles M.** Infección por el virus del papiloma humano en mujeres de edad mediana y factores asociados. *Rev Cuba Obstet Ginecol.* 2014;40(2):218-32.
30. **López-Torres Z, Ochoa-Marín SC, Alcaraz-López G, Leyva-Flores R, Ruiz-Rodríguez M.** Vulnerabilidad a infecciones de transmisión sexual y SIDA en mujeres en situación de desplazamiento forzado. Medellín, Colombia. *Investig y Educación en Enfermería.* 2010 [cited 2018 Jun 22];28(1):11-22. Available from: <https://goo.gl/HAZjRy>.
31. **Wiesner-Ceballos C, Vejarano-Velandia M, Caicedo-Mera JC, Tovar-Murillo SL, Cendales-Duarte R.** La citología de cuello uterino en Soacha, Colombia: representaciones sociales, barreras y motivaciones. *Rev Salud Pública.* 2006;8(3):185-96.
32. **Rubio-Mendoza ML.** Equidad en el Acceso los servicios de salud y equidad en la financiación de la atención en Bogotá. *Rev salud pública.* 2008;10(suppl 1):29-43.
33. **Kobayashi D, Otsubo T, Imanaka Y.** The effect of centralization of health care services on travel time and its equality. *Health Policy.* 2015;119(3):298-306. <http://doi.org/f259bg>.
34. **Chandrasekhar V.** Social determinants of health and health equity. *Indian J Public Health.* 2009;53(2):79-82.
35. **Godoy P.** La vigilancia y el control de las infecciones de transmisión sexual: todavía un problema pendiente. *Gac Sanit.* 2011;25(4):263-6.
36. **Blomquist PB, Miari VF, Biddulph JP, Charalambous BM.** Is gonorrhea becoming untreatable? *Future Microbiol.* 2014;9(2):189-201. <http://doi.org/f5x7hm>.
37. **Barbee LA.** Preparing for an era of untreatable gonorrhea. *Curr Opin Infect Dis.* 2014;27(3):282-7. <http://doi.org/f56q8n>.
38. **Mayaud P, Mabey D.** Approaches to the control of sexually transmitted infections in developing countries: old problems and modern challenges. *Sex Transm Infect.* 2004;80(3):174-82. <http://doi.org/dm4d5k>.
39. Organización Mundial de la Salud. Estrategia Mundial De Prevención Y Control De Las Infecciones De Transmisión Sexual: 2006-2015. Ginebra: OMS; 2007.
40. **Poku NK.** UN political declaration on HIV and AIDS: where to begin? *Lancet.* 2016;388(10046):743-4. <http://doi.org/crdc>.
41. **Bermúdez-Román V, Bran-Piedrahita L, Palacios-Moya L, Posada-Zapata IC.** Influencia del estigma en torno al VIH en el acceso a los servicios de salud. *Salud Publica Mex.* 2015;57(3):252-9.
42. **Farah-Quijano MA.** Cambios en las relaciones de género en los territorios rurales: aportes teóricos para su análisis y algunas hipótesis. *Cuadernos Des. Rural.* 2008;5(61):71-91.
43. **Castañeda-Abascal IE.** Reflexiones teóricas sobre las diferencias en salud atribuibles al género. *Rev Cuba Salud Publica.* 2007;33(2):1-20.
44. **Gómez-Gómez E.** Equidad, género y salud: retos para la acción. *Rev Panam Salud Pública.* 2002;11(5/6):454-61.
45. **Guarnizo-Herreño CC, Agudelo C.** Equidad de Género en el Acceso a los Servicios de Salud en Colombia. *Rev salud pública.* 2008;10(1):44-57.
46. **Llop-Gironés A, Vargas-Lorenzo I, Garcia-Subirats I, Aller MB, Vázquez-Navarrete ML.** Acceso a los servicios de salud de la población inmigrante en España. *Rev. Esp. Salud Pública.* 2014;88(6):715-34. <http://doi.org/crdd>.
47. **Restrepo-Zea JH, Silva-Maya C, Andrade-Rivas F, Vh-Dover R.** Acceso a servicios de salud: análisis de barreras y estrategias en el caso de Medellín, Colombia. *Rev. Gerenc. Polit. Salud.* 2014;13(27):242-65.
48. **Arrivillaga M, Correa D, Tovar LM, Zapata H, Varela MT, Hoyos PA.** Infecciones de transmisión sexual en la región Pacífica colombiana: implicaciones para población en situación de vulnerabilidad étnica, social y económica. *Pensamiento Psicológico.* 2011;9(16):145-52.