

ORIGINAL RESEARCH

DOI: <http://dx.doi.org/10.15446/revfacmed.v67n4.67159>

Correlation of HOMA-IR with BMI-for-age percentile in children and adolescents from the Soconusco region of Chiapas, Mexico

Correlación del HOMA-IR con el índice de masa corporal percentil en niños y adolescentes de la región Soconusco de Chiapas, México

Received: 20/09/2017. Accepted: 16/03/2018.

Consuelo Chang-Rueda¹ • Ana Cañas-Urbina² • Karina Trujillo-Murillo³ • Marisol Espinoza-Ruiz¹ • Jorge Feliciano-Díaz⁴ • Miguel Vázquez-Moreno⁵ • Ángel Lugo-Trampe³

¹ Universidad Autónoma de Chiapas - Faculty of Chemical Sciences - Department of Biochemistry and Immunology - Tapachula - Mexico.

² Universidad Autónoma de Chiapas - General Direction of Research and Postgraduate Studies - Tuxtla Gutiérrez - Mexico.

³ Universidad Autónoma de Chiapas - School of Human Medicine - Department of Biomedicine - Tapachula - Mexico.

⁴ Hospital General de Tapachula - Outpatient Service - Tapachula - Mexico.

⁵ Centro Médico Nacional Siglo XXI - Hospital de Especialidades - Biochemistry Medical Research Unit - Mexico City - Mexico.

Corresponding author: Consuelo Chang-Rueda. Departamento de Bioquímica e Inmunología, Facultad de Ciencias Químicas, Universidad Autónoma de Chiapas. Carretera a Puerto Madero km 1.5. Phone: +52 962 6251555, ext.: 112. Tapachula. Mexico. Email: consuelo.chang@unach.mx.

| Abstract |

Introduction: Insulin resistance (IR) is a major risk factor for developing diabetes *mellitus* type 2 and cardiovascular diseases. In pediatrics, morbidity and mortality associated with these diseases highlights the diagnostic importance of IR for primary care.

Objective: To determine Homeostatic Model Assessment Insulin Resistance (HOMA-IR) values and their correlation with BMI-for-age percentile in children and adolescents of the Soconusco region of Chiapas, Mexico.

Materials and methods: Cross-sectional study. Overweight and obesity prevalence was determined based on the Body Mass Index (BMI) percentile of 112 children (5-19 years old). Glucose and fasting insulin values were quantified and used for estimation of HOMA-IR.

Results: The combined prevalence of obesity and overweight was 66%, with insulin ($p=0.010$) and HOMA-IR ($p=0.015$) values higher than those of the normal weight group. The HOMA-IR values correlated positively with age ($r=0.636$), weight ($r=0.569$), height ($r=0.578$) and BMI percentile ($r=0.198$).

Conclusions: In the study population, HOMA-IR has a moderately significant correlation with an increase in BMI percentile.

Keywords: Insulin Resistance; Obesity; Diabetes *mellitus* type 2 (MeSH).

| Resumen |

Introducción. La resistencia a la insulina es un factor importante en el desarrollo de diabetes *mellitus* tipo 2 y de enfermedades cardiovasculares. En pediatría, su morbilidad y mortalidad resalta la importancia diagnóstica con fines de atención primaria.

Objetivo. Determinar los valores del homeostatic model assessment insulin resistance (HOMA-IR) y su relación con el índice de masa corporal percentil (IMCp) en niños y adolescentes de la región Soconusco, Chiapas.

Materiales y métodos. Estudio transversal. Se determinó sobrepeso y obesidad por IMCp en 112 pacientes pediátricos (5-19 años); se determinaron concentraciones de glucosa y de insulina sérica para estimar el HOMA-IR.

Resultados. Se encontró una prevalencia combinada de obesidad y sobrepeso de 66% con valores de insulina ($p=0.010$) y de HOMA-IR ($p=0.015$) más elevados que los del grupo de peso normal. El HOMA-IR se correlacionó positivamente con la edad ($r=0.636$), el peso ($r=0.569$), la talla ($r=0.578$) y el IMCp ($r=0.198$).

Conclusión. En la población de estudio, el HOMA-IR presenta una correlación moderadamente significativa con el aumento del IMCp.

Palabras clave: Resistencia a la insulina; Obesidad; Diabetes *mellitus* tipo 2 (DeCS).

Chang-Rueda C, Cañas-Urbina A, Trujillo-Murillo K, Espinoza-Ruiz M, Feliciano-Díaz J, Vázquez-Moreno M, et al. Correlation of HOMA-IR with BMI-for-age percentile in children and adolescents from the Soconusco region of Chiapas, Mexico. Rev. Fac. Med. 2019;67(4):447-50. English doi: <http://dx.doi.org/10.15446/revfacmed.v67n4.67159>.

Chang-Rueda C, Cañas-Urbina A, Trujillo-Murillo K, Espinoza-Ruiz M, Feliciano-Díaz J, Vázquez-Moreno M, et al. [Correlación del HOMA-IR con el índice de masa corporal percentil en niños y adolescentes de la región Soconusco de Chiapas, México]. Rev. Fac. Med. 2019;67(4):447-50. English. doi: <http://dx.doi.org/10.15446/revfacmed.v67n4.67159>.

Introduction

Insulin resistance (IR) is defined as a decreased response to the effect of the hormone, mainly on the liver, skeletal muscle and adipose tissue, and its pathophysiology has been strongly associated with the etiology of obesity. (1) IR has been described as a major risk factor for type 2 diabetes *mellitus* and cardiovascular disease, conditions that have high morbidity and mortality rates, which explains its diagnostic importance. (2)

To this end, the standard method used is the euglycemic clamp, which evaluates fasting glucose regulation based on insulin concentrations in vivo (3); however, this method is costly and invasive, and this hinders the frequency of its use in clinical practice. Other methods have been developed to determine IR indirectly, including Homeostatic Model Assessment Insulin Resistance (HOMA-IR), Quantitative Insulin Sensitivity Check Index (QUICKI) and the McAuley Index. (4) The first two use fasting glucose and insulin levels to establish a diagnosis, and the third is based on a relationship of fasting triglyceride and insulin levels. (5)

The use of HOMA-IR in routine clinical diagnosis has allowed observing that reference values range from 2.6 to 3.8 in adult and pediatric populations in different regions of the world (6-9); therefore, determining reference values for specific populations is highly suggested. (10-13)

The state of Chiapas has a total population of 5 217 908 inhabitants, of whom 74.7% live in poverty and 33.5% lack access to health services. (14,15) Regarding overweight and obesity, a combined prevalence in adolescents from urban areas of the state of 33% has been reported (16), with a HOMA-IR range for the capital between 0.4-7.5 and 21% in obese adolescents. (17) These is relevant when taking into account that more than 80% of deaths from noncommunicable diseases such as type 2 diabetes occur in low- and middle-income countries such as Mexico. (18)

The objective of this work was to determine HOMA-IR values and their correlation with the body mass index (BMI) for age percentiles in children and adolescents from the Soconusco region of Chiapas, Mexico.

Materials and methods

A cross-sectional study was conducted on individuals who used the clinical laboratory of the Tapachula Regional Hospital, Chiapas, from May to July 2015. The protocol was approved and filed on June 29, 2010 by the Research Committee of the General Directorate of Research and Graduate Studies of the Universidad Autónoma de Chiapas under code 02/QUI/RPR/290/10. This work complied with the ethical principles of the Declaration of Helsinki (19) and all participants and parents/guardians signed an informed consent. Children and adolescents of both sexes, between 5 and 19 years of age, presenting signs of metabolic control without presumptive diagnosis of the disease, were included in the study.

For biochemical determination (glucose and fasting insulin), a 6mL sample of peripheral blood without anticoagulant was taken from each study subject, with an 8-10 hour fast. The variables age, sex, height, weight and body mass index (BMI) were obtained by applying a physical examination questionnaire and taking anthropometric measurements with trained personnel at the Tapachula Regional Hospital. To measure weight and height, a clinical scale with a stadiometer was used (Clínica-160, Básculas Nuevo León®, México) and the Centers for Disease Control and Prevention (CDC) BMI percentiles were used to determine obesity and overweight (20); BMI between the 5th and 85th percentile was taken as normal weight, and BMI ≥ 85 was considered as obesity and overweight.

Glucose was determined using glucose oxidase-peroxidase (Glicemia enzyme AA, Wiener Lab., Argentina) and insulin was determined using the ELISA immunoassay (Insulin ELISA test, Diagnostic Automation, INC., USA); the automated ChemWell® 2910 (Awareness Technology Inc., USA) was used for both determinations. The HOMA-IR calculation was performed by adapting the formula proposed by Matthews *et al.* (21):

$$\text{Fasting glucose (mg/dL)} \times \text{fasting insulin } (\mu\text{U/mL}) / 405$$

The SPSS 21.0 software was used for statistical analysis, with 95% confidence. Data was described based on the following three parameters: frequencies, mean $\pm$ standard deviation, or median (interquartile range). For the comparison of the variables between groups, student's *t* and Mann-Whitney U tests were used according to the normality of the data, and Spearman's rank correlation coefficient was used to determine the correlation between variables.

Results

The study sample consisted of 112 subjects, 55 girls (49%), with a median age of 12 (9-13) years. Average weight, height and BMI were 47.0 ± 12.8 kg, 1.4 ± 0.1 m and 21.9 ± 3.8 kg/m², respectively. Regarding the comparison by sex, both weight and height, and BMI were homogeneous ($p=0.473$, $p=0.190$ and $p=0.696$). The general population showed a combined prevalence of overweight and obesity of 66%: 70% for males and 62% for females ($p=0.350$). The analysis of the biochemical variables revealed a mean glucose concentration of 80.5 ± 8.4 mg/dL, an insulin median of 7.7 $\mu\text{U/mL}$ (4.8-15.02), and an HOMA-IR of 1.6 (0.90-3.06). The three variables presented homogeneity between sexes (Table 1).

Table 1. Anthropometric and metabolic description of 112 children and adolescents in the Soconusco region of Chiapas, Mexico.

Measurement	Total n= 112	Female n= 55	Male n= 57	<i>p</i>
Age (years) *	12 (9-13)	12 (9-12)	12 (11-13)	0.095
Weight (kg) †	47.0 ± 12.8	47.8 ± 11.3	46.1 ± 14.1	0.473
Height (m) †	1.4 ± 0.1	1.4 ± 0.1	1.4 ± 0.1	0.190
BMI (kg/m ²) †	21.9 ± 3.8	22.1 ± 3.8	21.8 ± 3.9	0.696
Overweight/obesity (n, %) ‡	79.7	34.6	40.7	0.350
Glucose (mg/dL) †	80.5 ± 8.4	79.1 ± 7.5	82.0 ± 9.0	0.073
Insulin ($\mu\text{U/mL}$) *	7.7 (4.84-15.02)	9.0 (5.79-15.25)	7.0 (3.84-14.91)	0.100
HOMA-IR *	1.6 (0.90-3.06)	1.7 (1.08-2.97)	1.4 (0.75-3.11)	0.176

BMI: body mass index; HOMA-IR: homeostatic model assessment insulin resistance.

* Description with median. Contrast: Mann-Whitney U

† Description with median $\pm$ standard deviation. Contrast: Student's *t*

‡ Description of frequency. Contrast: Chi square

Source: Own elaboration.

According to the body weight diagnosis, the mean glucose concentration was homogeneous between groups (normal weight = 79.9 ± 8.1 mg/dL vs. overweight/obesity = 80.9 ± 8.6 mg/dL, $p=0.1456$). The overweight/obesity group presented insulin and HOMA-IR medians of 9.16 $\mu\text{U/mL}$ (5.5-15.99) and 1.8 (1.0-3.2), respectively, which were significantly higher than the values of the normal weight group at 5.9 $\mu\text{U/mL}$ (3.5-12.7) ($p=0.010$) and 1.3 (0.67-2.50) ($p=0.015$), respectively.

The correlation analysis found that HOMA-IR is positively and significantly related to the increase of the BMI percentile ($r=0.198$, $p=0.037$) (Figure 1) .




Figure 1. Correlation analysis of HOMA-IR with age, weight, height and body mass index percentile in 112 children and adolescents from the Soconusco region of Chiapas, Mexico. A) age; B) weight; C) height; D) BMI percentile. Log HOMA-IR: logarithm of homeostatic model assessment insulin resistance; BMI percentile: body mass index percentile. Source: Own elaboration.

Discussion

The combined prevalence of overweight and obesity (66%) of the study population is higher than the prevalence values reported at state and national level for adolescents by the 2012 National Health and Nutrition Survey (29%) (16) and by the 2016 National Health and Nutrition Survey of Medio Camino (36%). (22) This fact is highly relevant when considering that the prevalence found is close to that reported in Mexican adults (72%). (22) Finding such high values in early ages lead to infer early development of complications such as atherosclerosis and type 2 diabetes *mellitus*, the two diseases of greatest concern in Mexico. (23)

The glucose and insulin values found showed homogeneity between sexes. Glucose concentration (80.5 ± 8.4 mg/dL) coincides with previous studies conducted on adolescents in the states of Chiapas (85.8 ± 11.8) and Coahuila (73.2 ± 8.7) (17,24), while insulin levels 7.7 (4.8 - 15.02) μ U/mL and HOMA-IR 1.6 (0.90 - 3.06) are below what has been reported in those two states (13.0 ± 7.9 y 13.2 ± 10.5 μ U/mL, and 2.3 ± 1.4 and 2.9 ± 2.5 , respectively).

Even though the populations studied in Tuxtla Gutiérrez, Coahuila and Tapachula are mestizo, they have a different food culture. Consequently, insulin concentrations and HOMA-IR values of the former are probably higher because insulin levels become a compensatory mechanism for high glucose levels after consuming foods with high sugar content. (25)

On the other hand, the correlation of HOMA-IR with the BMI percentile ($r=0.198$; $p=0.037$) found in this study is lower than that reported with anthropometric parameters in adolescents from Jalisco, Mexico (BMI: $r=0.27$; $p<0.001$), Brazil (BMI: $r=0.366$; $p=0.031$) and Chile (BMI: $r=0.327$; $p<0.0001$). (26-28) According to the body weight analysis, the overweight and obese group had higher insulin and HOMA-IR levels than the normal weight group ($p=0.010$ and $p=0.015$), trend that has been reported in adolescents from Peru and Venezuela. (29,30)

For several years, IR has been described as a high impact factor for the alteration of the level of triglycerides and HDL cholesterol, lipids closely related to the development of metabolic syndrome in both adult and adolescent population. (31-34)

Conclusions

In the study population, the HOMA-IR presented a moderately significant correlation with the increase of the BMI percentile, reason why the present study can serve as evidence to propose the early diagnosis of IR as a preventive measure for the development of diabetes *mellitus* type 2 and cardiovascular disease in the adolescent population of Chiapas.

Conflicts of interest

None stated by the authors.

Funding

Project financed by the Unidades de Vinculación Docente program of the Universidad Autónoma de Chiapas 2010.

Acknowledgements

The authors would like to express their gratitude for the financial support provided by the Unidades de Vinculación Docente program of the Universidad Autónoma de Chiapas 2010, and the collaboration of the staff of the Hospital Regional de Tapachula, Chiapas.

References

1. Samuel VT, Shulman GI. The pathogenesis of insulin resistance: integrating signaling pathways and substrate flux. *J Clin Invest*. 2016;126(1):12-22. <http://doi.org/f75xws>.
2. Watve MG, Yajnik CS. Evolutionary origins of insulin resistance: a behavioral switch hypothesis. *BMC Evol Biol*. 2007;7:61. <http://doi.org/d7pvmq>.
3. Gungor N, Saad R, Janosky J, Arslanian S. Validation of surrogate estimates of insulin sensitivity and insulin secretion in children and adolescents. *J Pediatr*. 2004;144(1):47-55. <http://doi.org/d8tkdf>.

4. **Rijks J, Karnebeek K, van Dijk JW, Dorenbos E, Gerver WJ, Stouthart P, et al.** Glycaemic profiles of children with overweight and obesity in free-living conditions in association with cardiometabolic risk. *Sci Rep.* 2016;6:31892. <http://doi.org/f8x5k7>.
5. **Keskin M, Kurtoglu S, Kendirci M, Atabek ME, Yazici C.** Homeostasis model assessment is more reliable than the fasting glucose/insulin ratio and quantitative insulin sensitivity check index for assessing insulin resistance among obese children and adolescents. *Pediatrics.* 2005;115(4):500-3. <http://doi.org/b5pj8v>.
6. **Qu HQ, Li Q, Rentfro AR, Fisher-Hoch SP, McCormick JB.** The definition of insulin resistance using HOMA-IR for Americans of Mexican descent using machine learning. *PLoS One.* 2011;6(6):e21041. <http://doi.org/b633sd>.
7. **Gayoso-Diz P, Otero-González A, Rodríguez-Álvarez MX, Gude F, García F, De Francisco A, et al.** Insulin resistance (HOMA-IR) cut-off values and the metabolic syndrome in a general adult population: effect of gender and age: EPIRCE cross-sectional study. *BMC Endocr Disord.* 2013;13:47. <http://doi.org/gbd6db>.
8. **Yun KJ, Han K, Kim MK, Park YM, Baek KH, Song KH, et al.** Insulin resistance distribution and cut-off value in Koreans from the 2008-2010 Korean National Health and Nutrition Examination Survey. *PLoS One.* 2016;11(4):e0154593. <http://doi.org/f9wdnz>.
9. **Gutch M, Kumar S, Razi SM, Gupta KK, Gupta A.** Assessment of insulin sensitivity/resistance. *Indian J Endocrinol Metab.* 2015;19(1):160-4. <http://doi.org/c5s3>.
10. **Kim JW, Park SH, Kim Y, Im M, Han HS.** The cutoff values of indirect indices for measuring insulin resistance for metabolic syndrome in Korean children and adolescents. *Ann Pediatr Endocrinol Metab.* 2016;21(3):143-8. <http://doi.org/f9jskw>.
11. **Dev SB.** Unsolved problems in biology - The state of current thinking. *Prog Biophys Mol Biol.* 2015;117(2-3):232-9. <http://doi.org/f7d9r5>.
12. **Van der Aa MP, Fazeli-Farsani S, Knibbe CA, de Boer A, van der Vorst MMJ.** Population-Based Studies on the Epidemiology of Insulin Resistance in Children. *J Diabetes Res.* 2015;(2015). <http://doi.org/f72459>.
13. **Koh YK, Lee JH, Kim EY, Moon KR.** Acanthosis nigricans as a clinical predictor of insulin resistance in obese children. *Pediatr Gastroenterol Hepatol Nutr.* 2016;19(4):251-8. <http://doi.org/c5s4>.
14. **Instituto Nacional de Estadística y Geografía (INEGI).** Anuario estadístico y geográfico de Chiapas 2017. México: INEGI; 2017.
15. **Consejo Nacional de Evaluación de la Política de Desarrollo Social (CONEVAL).** Informe de pobreza y evaluación en el estado de Chiapas 2012. Ciudad de México: CONEVAL; 2012.
16. **Gutiérrez JP, Rivera-Dommarco J, Shamah-Levy T, Villalpando-Hernández S, Franco A, Cuevas-Nasu L, et al.** Encuesta Nacional de Salud y Nutrición 2012. Resultados Nacionales. Cuernavaca: Instituto Nacional de Salud Pública (México); 2012.
17. **Velasco-Martínez RM, Jiménez-Cruz A, Higuera-Domínguez F, Domínguez-de la Piedra E, Bacardí-Gascón M.** Obesidad y resistencia a la insulina en adolescentes de Chiapas. *Nutr Hosp.* 2009;24(2):187-92.
18. **Organización Mundial de la Salud (OMS)** [internet]. Enfermedades no transmisibles. OMS; 2018 [cited 2019 May 15]. Available from: <https://bit.ly/2HVkcOr>.
19. **Asociación Médica Mundial.** Declaración de Helsinki de la Asociación Médica Mundial. Principios éticos para las investigaciones médicas en seres humanos. Fortaleza: 64.ª Asamblea General de la AMM; 2013 [cited 2019 May 15]. Available from: <https://goo.gl/hvf711>.
20. **Centros para el Control y la Prevención de Enfermedades (CDC)** [internet]. Acerca del índice de masa corporal para niños y adolescentes. CDC; 2015 [cited 2019 May 15]. Available from: <https://bit.ly/2HyOYjQ>.
21. **Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC.** Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia.* 1985;28(7):412-9.
22. **Shamah-Levy T, Ruiz-Matus C, Rivera-Dommarco J, Kuri-Morales P, Cuevas-Nasu L, Jiménez-Corona ME, et al.** Encuesta Nacional de Salud y Nutrición de Medio Camino 2016. Resultados Nacionales. Cuernavaca: Instituto Nacional de Salud Pública (México); 2017.
23. **Instituto Nacional de Salud Pública (INSP)** [internet]. Diabetes, causa principal de muerte en México. INSP; 2015 [cited 2019 May 15]. Available from: <https://bit.ly/2QRCR2y>.
24. **González-Zavala MA, Velasco-Morales A, Terrazas-Flores JJ, de la Cruz-Galicia MG, Cepeda-Nieto AC, Hernández-del Río A.** Levels of insulin and HOMA-IR in adolescents in Saltillo, Coahuila, Mexico. *Medicina Universitaria.* 2015;17(67):80-7. <http://doi.org/f3g6nb>.
25. **Pinheiro AC, Canaan FA, Gonçalves R.** Insulinemia, ingesta alimentaria y metabolismo energético. *Rev Chil Nutr.* 2008;35(1):18-24. <http://doi.org/bjpk5>.
26. **Ortega-Cortes R, Trujillo X, Hurtado-López EF, López-Beltrán AL, Colunga-Rodríguez C, Barrera-de Leon JC, et al.** Models predictive of metabolic syndrome components in obese pediatric patients. *Arch Med Res.* 2016;47(1):40-8. <http://doi.org/f8h7tg>.
27. **Morais PR, Sousa AL, Jardim T, Nascente FM, Mendonça KL, Povoa TI, et al.** Correlation of insulin resistance with anthropometric measures and blood pressure in adolescents. *Arq Bras Cardiol.* 2016;106(4):319-26. <http://doi.org/c5s5>.
28. **Barja S, Arnaiz P, Domínguez A, Villarroel L, Cassis B, Castillo O, et al.** Insulinemia e índice HOMA en niños y adolescentes chilenos. *Rev Med Chile.* 2011;139(11):1435-43. <http://doi.org/3cs>.
29. **Rojas-Gabullí MI, Núñez O, Del Águila C, Briceño M, Valenzuela N.** Resistencia a insulina en adolescentes obesos. *An Fac Med.* 2010;71(1):13-7.
30. **Ruiz N, Rangel A, Rodríguez C, Rodríguez L, Rodríguez V, Varela I.** Relación circunferencia de cintura/talla: Predictor de insulino-resistencia y riesgo cardiometabólico agrupado en adolescentes. *Arch Venez Pueric Pediatr.* 2015;78(1):6-12.
31. **Simental-Mendía LE, Castañeda-Chacón A, Rodríguez-Morán M, Aradillas-García C, Guerrero-Romero F.** Relationship between elevated triglyceride levels with the increase of HOMA-IR and HOMA-β in healthy children and adolescents with normal weight. *Eur J Pediatr.* 2015;174(5):597-605. <http://doi.org/f69pcf>.
32. **Medina-Urrutia A, Juárez-Rojas JG, Martínez-Alvarado R, Jorge-Galarza E, Posadas-Sánchez R, Cardoso-Saldaña G, et al.** High-density lipoprotein subclasses distribution and composition in Mexican adolescents with low HDL cholesterol and/or high triglyceride concentrations, and its association with insulin and c-reactive protein. *Atherosclerosis.* 2008;201(2):392-7. <http://doi.org/cn54wx>.
33. **Wall-Medrano A, Ramos-Jiménez A, Hernández-Torres RP, Villalobos-Molina R, Tapia-Pancardo DC, Jiménez-Flores JR, et al.** Cardiometabolic risk in young adults from northern Mexico: Revisiting body mass index and waist-circumference as predictors. *BMC Public Health.* 2016;16:236. <http://doi.org/f8ddh9>.
34. **Cifuentes-Goches JC, Gómez-López JD, Hernández-Ancheyta L, Flores-Huerta SE, Incháustegui-Arias JL, Cañas-Urbina AO.** Hipertrigliceridemia e hipoalipoproteinemia. Su impacto para diagnosticar síndrome metabólico. *Rev Med Inst Mex Seguro Soc.* 2012;50(3):301-6.