

# Determination of predominant soluble salts in soils of the irrigation district Alto Chicamocha of Boyacá

Determinación de sales solubles predominantes en suelos del distrito de riego Alto Chicamocha de Boyacá

doi: 10.15446/rfnam.v71n3.72375

Ingrid Walteros Torres<sup>1\*</sup>, Germán Cely Reyes<sup>1</sup> and Diego Moreno Pérez<sup>1</sup>

## ABSTRACT

### Keywords:

Anions  
Cations  
Geostatistics  
E.C  
Water table

In Boyacá, the Alto Chicamocha irrigation and drainage district (DRACH, by its initials in Spanish) is the department's main agricultural production unit, covering an area of 8016.78 ha and due to the natural conditions and the management that has been given to the high basin of the Chicamocha River, salinization has been recognized as a limiting factor. Therefore, we sought to determine the predominant soluble salts in the soils that comprise the DRACH. Based on the chemical soil analysis information of 301 samples, obtained from studies conducted by GISSAT-UPTC and Corpoica, E.C, pH, anions and cations present in the soil were determined. For the spatial analysis of the variables studied, the ArcGis 10.3 software was used. Thirty-one water samples were collected in wells of the phreatimetric network to carry out the chemical characterization of the water. It was found that 48.01% of the soils of the district were non-saline, 22.93% slightly saline, 14.74% moderately saline and saline 14.33%. The main soluble salts in the soil were  $\text{Na}_2\text{SO}_4$ ,  $\text{Ca}_2\text{SO}_4$ ,  $\text{NaCl}_2$  and  $\text{CaCl}_2$ , which are related to the lacustrine origin and the presence of thermal springs in the region. The areas with greater problem of salinization occur in the municipalities of Tibasosa, Patrocinio, Ucacá, Las Vueltas; in Santa Rosa de Viterbo in the village of Salitre; in Duitama in the villages Cebadero and Higuera with E.C. greater than 2 dS  $\text{m}^{-1}$ .

## RESUMEN

### Palabras clave:

Aniones  
Cationes  
Geoestadística  
C.E  
Nivel freático

El distrito de riego y drenaje del Alto Chicamocha (DRACH por sus siglas en español), es la principal unidad de producción agropecuaria del departamento de Boyacá, Colombia, abarca un área de 8016,78 hectáreas y debido a las condiciones naturales y al manejo que se le ha venido dando a la cuenca alta del Río Chicamocha, la salinización ha sido reconocida como limitante. Por lo anterior se buscó determinar las sales solubles predominantes en los suelos que comprenden el DRACH. Con base en la información de análisis químicos de suelos de 301 muestras, obtenidas de estudios realizados por el GISSAT-UPTC y Corpoica, se determinó C.E, pH, aniones y cationes presentes en el suelo. Para el análisis espacial de las variables estudiadas se usó el software ArcGis 10.3. Se colectaron 31 muestras de agua en pozos de la red freática para realizar la caracterización química del agua. Se encontró que el 48,01% de los suelos del distrito eran no salinos, el 22,93 % ligeramente salino, el 14,74% moderadamente salino y salinos el 14,33%. Las principales sales solubles en el suelo fueron el  $\text{Na}_2\text{SO}_4$ ,  $\text{Ca}_2\text{SO}_4$ ,  $\text{NaCl}_2$  y  $\text{CaCl}_2$ , las cuales están relacionadas con el origen lacustre y la presencia de termales en la región. Las zonas con mayor riesgo de salinización se presentan en los municipios de Tibasosa, veredas Patrocinio, Ucacá, las Vueltas; en Santa Rosa de Viterbo en la vereda Salitre; en Duitama en las veredas Cebadero e Higuera con C.E. mayores a 2,00 dS  $\text{m}^{-1}$ .

<sup>1</sup> Facultad de Ciencias Agropecuarias. Universidad Pedagógica y Tecnológica de Colombia. Avenida Central del Norte 39-115, 150003 Tunja, Colombia.

\* Corresponding author: <ingrid.walteros@uptc.edu.co>

Irrigation is considered a fundamental element in agriculture due to its effect on increasing production, improving the quality of products, sustainable intensification of land use and its contribution to safety food (FAO, 2000; 2011). In 2012, there were more than 324 million hectares in the world equipped for irrigation, of which approximately 85% or 275 million are effectively irrigated; by 2010, China became the country with the largest area of irrigation, and together with India, it covers 42% of the world's irrigation (AQUASTAT-FAO, 2014). In this way, irrigation has reduced the dependence on seasonal agriculture, thus achieving high agricultural production (Rhoades *et al.*, 1992). In soils such as those in semi-arid regions where water requirements for crops are high and are necessarily supplied by irrigation districts, the use of these causes serious problems of environmental degradation that hampers the growth of crops and regional production in general (Abbas *et al.*, 2013). Although irrigation districts bring advantages to agricultural systems, they have also brought with them environmental problems, such as: Salinization of soils, contamination of surface and groundwater bodies, changes in the landscape, and stress on plants.

The mobilization of salts in the soil is very variable among the different irrigation zones and according to Chedlia *et al.* (2012). Salinity depends on several factors such as the amount of salts present in the water, the texture of the soil, the distance from the intake to the district, the hydrogeology of the area, irrigation and drainage, and climatic conditions (rainfall regime, average temperature) (Duncan *et al.*, 2008). For Girón-Ríos *et al.* (2009), salinity is a complex process of chemical degradation, which influences significant changes in the physical properties of soils and is affected mainly by the presence of salts in irrigation water and efficiency of the same (Aragüés *et al.*, 2011).

Colombia has not been unrelated to the problem of salinization, in the different soil studies conducted, it is common to find that the soils have degradation phenomena, with erosion and salinization being the main problems in the territory. In this regard, the country ratified the United Nations Convention to Combat Desertification and Drought (UNCCD), which considers as an important strategy the identification

and monitoring of soil degradation processes at the national, regional and local levels (FAO, 2000; 2011). Currently, public institutions of national order have been making important efforts to diagnose salinity problems in the country's soils, such as the national map of soil degradation by salinization carried out by IDEAM in 2017. According to this study, 14,041,883 ha (12.3%) of the country's soils (continental and insular areas) present some degree of degradation due to salinization. The very severe and severe degradation occupies 2,726,757 ha (2.4%); the moderate degree 8,885,369 ha (7.8%) and with 2,449,757 ha slightly (2.1%).

In Boyacá, the most important agricultural production unit in the department and the largest sprinkler irrigation district that has been built in a cold climate zone, is the DRACH, which has eleven irrigation units (Pacheco *et al.*, 2004) and covers an area of 8016.78 ha. The problem of salinization has been recognized as a limiting factor in the agricultural soils that currently correspond to DRACH since 1960, the Colombian Institute for Agrarian Reform – INCORA (by its initial in Spanish), carried out drainage works to enable land to agriculture. The soils of the Tundama - Suamox valley are very vulnerable to salinization due to the natural conditions and the management that has been given to the upper basin of the Chicamocha River. For 2012 after an extreme rainfall weather event where 3000 ha were affected, the main victims were the agricultural producers who benefit from the DRACH and both urban and rural inhabitants of the municipalities of Paipa, Duitama, Nobsa, Sogamoso and to a greater extent Tibasosa (GISSAT, 2012), after this extreme weather event the problem of salinization was evidenced, where high electrical conductivities and saline scabs were observed. Therefore, with this research we sought to determine which were the most predominant soluble salts in the soils that comprise the DRACH and its special distribution.

## MATERIALS AND METHODS

### Location

The research was conducted in the DRACH which covers the towns of Paipa, Duitama, Tibasosa, Santa Rosa de Viterbo, Sogamoso and Nobsa (Figure 1) over the eleven irrigation units (Table 1).




Figure 1. Location of the Irrigation and Drainage District of Alto Chicamocha.

Table 1. Irrigation units that define the DRACH.

N°	Irrigation Unit	Town	Rural áreas
1	Monquirá	Tibasosa Nobsa Sogamoso	Patrocinio Caleras Siatame, Área Urbana
2	Ministerio	Tibasosa Nobsa	Suescún, Centros, Patrocinio, Resguardo Ucuenga, Caleras
3	Tibasosa	Tibasosa Nobsa	Suescún, Boyera, Centros, Área Urbana. Ucuenga
4	Las Vueltas	Tibasosa	Vueltas
5	San Rafael	Tibasosa Santa Rosa de Viterbo Nobsa	Peña Negra, Suescún Salitre Punta Larga, San Martín, Dicho, Ucuenga
6	Cucho	Santa Rosa de Viterbo  Duitama Nobsa	Cucho, Salitre, Cachavita, Creciente, Tungaquita, La Chorrera Tocogua San Martín
7	Duitama	Duitama  Tibasosa	San Lorenzo de Abajo, Aguatendida, Tocogua, Área Urbana Chorrito
8	Ayalas	Duitama Tibasosa	San Lorenzo de Abajo Ayalas
9	Surba	Duitama	San Lorenzo de Arriba, San Lorenzo de Abajo
10	Holanda	Paipa	Toibita, Cruz de Bonza, Romita, Caños, Paipa Área Urbana
11	Pantano de Vargas	Paipa	Rincón De Vargas, Pantano De Vargas, Varguitas, Caños

The DRACH is located in the department of Boyacá, approximately 180 km from Bogotá, with an average temperature of 14 °C, with an average rainfall of 778 mm, an average altitude of 2560 m and a relative humidity of 70% (Martínez *et al.*, 2008). Most of the soils are of agricultural vocation, however, dairy cattle stands out and, on a smaller scale, dual-purpose cattle (GISSAT, 2012).

In the DRACH data were taken for soils and waters which were georeferenced and projected to the Magna Colombia Bogotá coordinate system. The samples were faced in the dry period, after the La Niña phenomenon (2010-2012).

### Soils analysis for Salinity

Based on the chemical soil analysis information from 301 samples, obtained from studies conducted by the Interinstitutional Research Group on Tropical Acid Sulphate Soils (GISSAT-UPTC, 2012) and Corpoica; electrical conductivity (E.C) by saturation extract – conductivity meter, hydrogen potencial (pH) by the 1:1 ratio method and the soluble salts in the soil were determined, for this the cations were obtained by atomic absorption.

To establish the spatial tendency of salinity, the geostatistical analysis of the E.C, pH, Sulfates ( $\text{SO}_4^-$ ), Sodium ( $\text{Na}^{2+}$ ), was carried out. For the spatial analysis of the variables studied, the Geostatistical analyst tool of the ArcGis 10.3 software was used, the interpolation method used was ordinary Kriging. To classify the salinity of DRACH soils, the following ranges of electrical conductivity (E.C.) were used: saline ( $\geq 2 \text{ dS m}^{-1}$ ), moderately saline ( $\geq 1.5 < 2 \text{ dS m}^{-1}$ ), slightly saline ( $\geq 1 < 1.5 \text{ dS m}^{-1}$ ) and not saline ( $< 1 \text{ dS m}^{-1}$ ), according to Castro and Gómez (2010). Regarding the classification used for sulfates ( $\text{SO}_4^-$ ), the following ranges were used: Under 11 ppm, Optimum 38 ppm and High 64 ppm according to Castro and Gómez (2010). For sodium, the ranges were used: low  $0.1 \text{ cmol kg}^{-1}$ , medium  $0.1-0.5 \text{ cmol kg}^{-1}$  and high  $> 0.5 \text{ cmol kg}^{-1}$  according to Castro and Gómez (2010).

### Groundwater analysis for Salinity

Thirty-one water samples were collected wells of the phreatimetric network to perform the chemical

characterization corresponding to pH with potentiometer, E.C with conductivimeter, sulfates by colorimetry, and by titulation were obtained the chlorids, carbonates, Bicarbonates and nitrates, the cationes ( $\text{Na}^{2+}$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  and K) by atomic absorption. This information allowed to determine the quality of these waters and their possible relationship with the salinization processes. The samples were analyzed in the soil and water laboratory of the Faculty of agricultural sciences of the UPTC (by its initials in Spanish), with the analytical methodology (Analytical Control Soil Laboratory- ICONTEC).

The research had a non-experimental design, since independent variables could not be manipulated, where the phenomenon of salinization was observed in its natural context within the irrigation District. For each of the variables, the basic descriptive statistics were calculated, in which the mean, median, kurtosis and asymmetry were calculated, as well as the parameters of the semivariogram (Sill, Nugget and Range) and the cross validation (Table 2).

## RESULTS AND DISCUSSION

As it is mentioned by Narváez *et al.* (2014), classical statistics and geostatistics are some of the tools for the analysis of soil salinity that contributes to the identification of affected areas and the monitoring of spatio-temporal variations, in the Table 2 the results for the geostatistical analysis of the variables studied are observed.


The statistical model that was best adjusted for the variables pH (Figure 2), E.C (Figure 3),  $\text{Na}^{2+}$  (Figure 4) was the exponential, while for the sulfates was the Rational Quadratic model (Figure 5). These variables were studied because they are the most representative soluble salts of the zone.

According to the Kolmogorov-Smirnov normality test, the variables analyzed show a non-normal behavior except for pH (Table 2). For Cressie (1993), normality is not a mandatory requirement for the analysis of geostatistical data, however, it must be considered that the distribution of the data don't show a very long tail, since it compromises the results for the Kriging estimates.


Variability is an intrinsic characteristic of each property and its specific behavior for each soil condition, use and

**Table 2.** Descriptive and spatial statistics of the variables studied.

Statistics	Variables	E.C.	pH	Na <sup>2+</sup>	SO <sub>4</sub> <sup>=</sup>
Mean		1.5	5.411	4.446	2.456
Sill (CO+C)		0.64	0.9483	1.119	0.909
Range		2640.84	1280	3055.31	3874.58
Nugget (CO)		0.35	0.0531	0.725	0.494
Cross-validation (Error mean)		0.0178	-0.03	0.010	-0.079



**Figure 2.** Semivariogram of pH.



**Figure 3.** Semivariogram of E.C in the DRACH.

management; therefore, interpreting the autocorrelation of a variable along its distribution in one area, and its correlation with respect to another variable or variables, is a study that has a large number of immersion factors,

and requires clear knowledge of the factors that condition it (Jaramillo *et al.*, 2008). The Range is the zone of influence and corresponds to the distance from which two observations are independent (Giraldo, 2002).



All the variables had a moderate spatial dependence (Table 2) according to the classification proposed by Cambardella *et al.* (1994), since the relationship between the plateau and the nugget effect  $[C/(Co+C)]$  is between

0.25 and 0.75, the pH showed a range of 1287.93 m, the E.C a range of 2048.9 m,  $Na^{2+}$  a range of 3055.31 and the  $SO_4^{=}$  a range of 3874.58, indicating that there is a spatial correlation in each of the variables at that distance.




Figure 4. Semivariogram of  $Na^{2+}$




Figure 5. Semivariogram of  $SO_4^{=}$

The results of the descriptive analysis for the soil variables showed that the E.C. of the soil saturation extract, presented an average value of 1.77  $dS\ m^{-1}$ ; the highest value was registered in the town of Santa Rosa rural area "El Salitre" with an E.C. of 6.57  $dS\ m^{-1}$  and the lowest in the municipality of Paipa with 0.16  $dS\ m^{-1}$ . Figure 6 shows the behavior of the mean of the E.C. values for the districts that make up the district.

It should be noted that 48.01% of the district's soils are found in the range of E.C.  $< 1\ dS\ m^{-1}$ , which classifies

it as non-saline (Table 3), 22.93% between  $\geq 1.0 < 1.5\ dS\ m^{-1}$  which is classified as slightly saline, 14.74% between  $\geq 1.5 < 2\ dS\ m^{-1}$  which are moderately saline and 14.33% have values higher than  $\geq 2\ dS\ m^{-1}$  are saline soils (Figure 7), which indicates that they are soils that begin to have salt problems, according to Castro and Gómez (2010). These values coincide with those reported by Mercado *et al.* (2011) who in the Doctrina Irrigation District, Colombia found values higher than 3  $dS\ m^{-1}$ , which indicated salinity problems for those areas. The electrical conductivity of the soil is a function of the clay content

and water content (Kurtulus *et al.*, 2009, Narváez *et al.*, 2014), with the topography, and the phreatic level which allows high variability to be present for this parameter (Ulset *et al.*, 1998).




Figure 6. General average of the E.C. in the soils of rural DRACH.

Table 3. E.C. Ranges for DRACH soils.

E.C. Range (dS m <sup>-1</sup> )	Area (ha)	Percentage (%)
< 1.0	3848.84	48.01
≥ 1.0 < 1.5	1838.09	22.93
≥ 1.5 < 2.0	1181.34	14.74
≥ 2.00	1148.51	14.33
Total	8016.78	100





Figure 7. Spatial distribution of E.C. in the DRACH soils.

The pH presented an average value of 5.52. In general, it was between the range of 5-6 that corresponds to moderately acid soils, with 46.64% of the total area of the

district (Figure 8). The *Resguardo* area had a mean value of 7.28, that may be associated with the presence of salts such as sodium chloride ( $\text{NaCl}_2$ ) (Mercado *et al.*, 2011).



**Figure 8.** Spatial distribution of pH in the DRACH soils.

For the cations, it was evidenced that the element with the greatest presence in DRACH soils was sodium, followed by Ca in less quantity. Considering the ranges provided by Castro and Gómez (2010), for Na greater than  $0.5 \text{ cmol kg}^{-1}$  and for Ca higher than  $6 \text{ cmol kg}^{-1}$ , there is a high grade of salinization due to very high concentration levels. In Figure 9, the behavior of Na within the district is observed, 73.93% of the soils have low concentrations with levels of  $0.1 \text{ cmol kg}^{-1}$ , 21.49% are in the middle range with values between  $0.1\text{-}0.5 \text{ cmol kg}^{-1}$  and 4.57% have high levels with values greater than  $0.5 \text{ cmol kg}^{-1}$ . High values were found in the Salitre area with  $4.29 \text{ cmol kg}^{-1}$ , followed by El Cebadero with  $4.085 \text{ cmol kg}^{-1}$  that may be related to the saline deposits ( $\text{Na}_2\text{SO}_4$  and  $\text{NaCl}_2$ ), which are of the characteristics of the thermal sources of Paipa (Moreno and Fечи, 2006). Taiz *et al.* (2006), mention that the Na concentration of a sodium soil can not only directly damage the plants, but also degrade the structure of the soil, reducing the porosity and being a highly hygroscopic element, it traps the water molecules of the soil, which causes the water of hydration to decrease for other nutrients, also affecting the structure of the soil by breaking up its particles (Madueño *et al.*, 2006). Besides that the hydrolysis of the sodium clays, leads to the alkalinization of the profile, and these cause an intense mineral alteration (Mata *et al.*, 2014). While if  $\text{Ca}^{2+}$  and

soluble salts are very abundant in the soil, it is likely that the profile is very poorly differentiated, but its structure tends to be stable, as a result of the flocculating action of  $\text{Ca}^{2+}$  so the high osmotic pressure of the soil solution is responsible for low productivity (Mata *et al.*, 2014).

$\text{Ca}^{2+}$  reported high values in the Salitre rural zone with  $42.01 \text{ cmol kg}^{-1}$ , followed by las Vueltas with  $30 \text{ cmol kg}^{-1}$  while the lowest values were found in the village of Rincon de Vargas with  $4.236 \text{ cmol kg}^{-1}$ . Mercado *et al.* (2011) obtained Ca values higher than  $6 \text{ cmol kg}^{-1}$  in an average of 89.5% of the total area studied.

The sulphates ( $\text{SO}_4^{=}$ ) were the dominant anions, mainly due to their high solubility, the area of Cebadero presented an average of  $93.298 \text{ mg kg}^{-1}$  being the highest value, followed by the area of Salitre with  $54.238 \text{ mg kg}^{-1}$ ; they are high values considering the ranges established by Castro and Gómez (2010). In Figure 10, the behavior of the sulphates in the district is observed, 48.90% of the area is at a low level (11 ppm), 48.86% of the area presents an optimum range with 38 ppm and the 0.30% of the district presents problems due to high levels (64 ppm).

For chlorides (Cl) the rural area *el Resguardo* reported the value of the highest average with  $42.80 \text{ mg kg}^{-1}$






Figure 9. Spatial distribution of  $\text{Na}^{2+}$  in the DRACH soils.




Figure 10. Spatial distribution of  $\text{SO}_4^{=}$  in the DRACH soils.

followed by Siatame in Sogamoso with  $30.76 \text{ mg kg}^{-1}$  the behavior of these is noted in Table 3. These latter are at a low level according to Castro and Gómez (2010), where the high ranges are  $142 \text{ mg kg}^{-1}$  and above. Qadir *et al.* (2007) state that the predominant soluble salts in saline soils are sulfates, chlorides, bicarbonates of  $\text{Na}^{2+}$ ,  $\text{Ca}^{+2}$  and  $\text{Mg}^{+2}$ . Na is the cation that most binds to sulfates and chlorides to form salts, with less frequency found potassium and bicarbonates, carbonates and nitrates (Dorronsoro, 2011; Ramírez, 2011).

Fernández *et al.* (2007) indicate that the high variability of the parameters mentioned above may be due to the

fact that soil properties such as interchangeable cations, sulphates and others not related to soil morphology are affected by the use and / or management. Similarly, Narváez *et al.* (2014) indicate that in saline soils, the depth of the water table, the evapotranspiration rate, and other geohydrological factors impact on the chemical properties, resulting also in a high spatial and temporal variability of the same. According to this, the high variability obtained for the parameters is generated by the interaction of the different properties of the soil with its immediate environment, this means, agricultural practices, variations in climate among others, agricultural practices, variations in climate among others, cause

changes at different scales (Mallarino and Wittry, 2004). For the waters, it was found that the water tables oscillated between 0.54 and 2.99 m in the irrigation units San Rafael in Nobsa and Cucho in Santa Rosa de Viterbo; these values coincide with those reported by Castro and Gómez (2015) who argue that in the dry season about 30% of the area, the reading and analysis levels of the District's aeration profile zone are between 0.0 and 1.0 m, while in the critical rainy season there is a rise in the water table, reaching times of flooding, which can affect up to 45% of the area. It is worth noting that the waters are frequented by sales of soluble, the upper basins of the Chicamocha River have geological units composed of a very important lithological record, mostly composed of sedimentary rocks with some outcrops of igneous rocks in the Paipa and Iza surroundings (Moreno and Fechi, 2006). The phreatic mantles are found on calcareous materials can cause salinization, if these are located in climates, they are characterized by values of evapotranspiration superior to those of precipitation, the phreatic levels

can ascend by the capillarity to the surface of the soils (IGAC, 2012).

Water quality analysis for irrigation was carried out in 31 wells (Figure 11). According to the classification of waters by to the Riverside standards (US Salinity Laboratory - USLS), all of them have problems of salinity and sodium (Olías *et al.*, 2005). The average value of the E.C was  $3.12 \text{ dS m}^{-1}$ , for pH was 6.34. In this regard, Taiz *et al.* (2006) mention that the higher the concentration of salts in water, the greater the electric conductivity and the lower the osmotic potential (the higher the osmotic pressure). The above allows to relate the E.C. of the soil with the water used for irrigation, in some units such as Ministry where E.C of  $4.29 \text{ dS m}^{-1}$  was observed in the soil and  $9.79 \text{ dS m}^{-1}$  in the water; in Cucho with  $4.66 \text{ dS m}^{-1}$  in water and  $5.64 \text{ dS m}^{-1}$  in soil, and finally in the irrigation unit of Vueltas with  $8.54 \text{ dS m}^{-1}$  in water and  $2.6 \text{ dS m}^{-1}$  on soil, indicating that water used for irrigation may be contributing to the salinization of soils in the district.




Figure 11. Spatial location of the water sampling points in the DRACH.

According to investigations by Alfaro and Ingeominas (2010), in Paipa springs there are problems with sulphated waters, as well as high concentrations of  $\text{Na}^{2+}$ ,  $\text{K}$ ,  $\text{Ca}^{+2}$ ,  $\text{Mg}^{+2}$ ,  $\text{Cl}^{-1}$ ,  $\text{SO}_4^{-}$ ,  $\text{HCO}_3^{-}$ . For Porras (2010), in the present thermal manifestations, alkaline-chlorinated waters appear, probably these are related to deep hydraulic circuits (reservoir), sulphate-alkaline waters

that owe their origin to the addition of alkaline sulphates to the preceding waters, the area is characterized by the predominance of essentially Cretaceous and Tertiary sedimentary rocks, and the presence of possible volcanic necks with an approximate age of 2.5 million years, in addition there are thermal manifestations of high temperature, probably due to the presence of an

acidic magmatic intrusion, located at a depth of 5 km. There is possible a presence of two thermal aquifers in granular and silicic sedimentary rocks with predominant secondary permeability is highlighted, in addition, the Honda or Río Salitre stream carries a high quantity of salts, supplied by the thermal springs that appear in the middle and lower area of the sub-basin (POT Paipa, 2012). Moreno and Fечи (2006), affirm that the presence of thermomineral waters in the Paipa sector are affecting the DRACH soil, due to the fact that at present they are drained to a great extent to the Vargas canal, generating: disaggregation of soil particles caused by sodium sulphates, ionic contamination by chlorine and sodium, mainly due to inadequate handling of thermal or sulfuric waters (IGAC, 2012).

## CONCLUSIONS

The main soluble salt present in the soil was sodium sulphate ( $\text{Na}_2\text{SO}_4$ ), followed by calcium sulfate ( $\text{Ca}_2\text{SO}_4$ ), sodium chloride ( $\text{NaCl}_2$ ) and calcium chloride ( $\text{CaCl}_2$ ), which are highly soluble and toxic for the plants.

It was found that the areas with the highest problem of salinization are in Tibasosa in Patrocinio, Ucaca, Las Vueltas zones; in Santa Rosa de Viterbo in the Salitre zone and finally in Duitama in Cebadero and Higuera zones, as they have electrical conductivities greater than  $2 \text{ dS m}^{-1}$ .

The groundwater of the sector plays an important role in the dynamics of salinization of the DRACH, due to the connection with the upper basin of the Chicamocha River, which presents geological units composed of a very important lithological record.

The presence of the Paipa springs with sulfated water problems and high concentrations of cations and anions strongly influence the salinization of DRACH soils.

## REFERENCES

- Abbas A, Khan S, Hussain N, Hanjra M, Akbar S. 2013. Characterizing Soil Salinity in Irrigated Agriculture Using a Remote Sensing Approach. *Physics and Chemistry of the Earth* 55-57(201): 43-52. doi: 10.1016/j.pce.2010.12.004.
- Alfaro C and Ingeominas. 2010. Geothermal resources in Colombia. International Congress in Environmental Sciences and Technologies. IV session: Energy, production and sustainable consumption. Univalle, Cali.
- AQUASTAT-FAO. 2014. Irrigation, Deforested Crops, Environment. At: <http://www.fao.org/nr/water/aquastat/didyouknow/indexesp3.stm>. Consultation: June, 2016.
- Aragüés R, Urdanoz V, Cetin M, Kirdac C, Daghari H, Ltifi W, Lahlou M, Douaik A. 2011. Soil salinity related to physical soil characteristics and irrigation management in four Mediterranean irrigation districts. *Agricultural Water Management* 98(6): 959-966. doi: 10.1016/j.agwat.2011.01.004.
- Cambardella CA, Moorman TM, Novak JM, Parkin TB, Karlen DL, Turco RF and Konopka AE. 1994. Field scale variability of soil in central Iowa soils. *Soil Science Society of America Journal* 58: 1501-1511.
- Castro H and Gómez M. 2010. Fertility of soils and fertilizers. Pp 238-256. In: Burbano, H. Silva, F. Soil science, basic principles. First edition. Published by Sociedad Colombiana de la Ciencia del Suelo. 562p.
- Chedlia B, Ahmeda B, Salwa A, Magdich A, Bechir B, Rouina B, Boukhris A, Abdullah F. 2012. Effects of irrigation with saline water on the distribution of soil salinity and some physiological responses of cultivated field of olive chemlali. *Journal of Environmental Management* 113: 538-44. doi: 10.1016/j.jenvman.2012.03.016
- Dorransoro C. 2011. Environmental Impacts of Soil Pollution - Soluble Salts - Part 02. In: [www.estrucplan.com.ar](http://www.estrucplan.com.ar). Check November 2017.
- Duncan A, Bethune G, Thayalakumaran T, Christen E, McMahon T. 2008. Management of salt mobilisation in the irrigated landscape a review of selected irrigation regions. *Journal of Hydrology* 351: 238-252. doi: 10.1016/j.jhydrol.2007.12.002.
- FAO. 2000. Irrigation in Latin America and the Caribbean in figures. United Nations Organization for Agriculture and Food, 345p.
- FAO. 2011. Proceedings of the Global Forum on Salinization and Climate Change. Valencia, 25-29 October 2010. En: [http://www.fao.org/uploads/media/BOOK\\_printing.pdf](http://www.fao.org/uploads/media/BOOK_printing.pdf). Consulta octubre 2017.
- Fernández O, Pérez R, Gubinelli A and Matheus J. 2007. Influence of soil spatial variability on parameters of sugarcane (*Saccharum officinarum*) in Monay, Trujillo State. *Andean Agriculture* 13: 39-52.
- Giraldo HR. 2002. Introducción a la geoestadística: Teoría y aplicación. Universidad Nacional de Colombia. Bogotá. 94p.
- Girón-Ríos Y, Oleschko K, Jeanfrancois P, Hernández M, Camarillo E and Velázquez J. 2009. Fractal analysis of the reflectance of saline soils. *Agro-science* 43(4): 403-416.
- IDEAM. 2017. Mapa nacional de degradación de suelos por salinización 2017.
- Inter-institutional Research Group on Sulphated Soils Tropical Acids. GISSAT - UPTC. 2012. Impact assessment and management plan for the soils affected by the floods in the valleys of Tundama and Sugamuxi (Boyacá). Tunja.
- IGAC. 2012. Protocol of Degradation of soils and lands by salinization. 323p.
- Jaramillo JD, González SH and Álvarez MF. 2008. Spatial variability of some physico-mechanical properties of heavy textured soils. *CES Magazine. Veterinary Medicine* 3(2).
- Kurtulus C, Canbay M, Demir N and Gider D, 2009. Salinity investigation of the region east to the Izmit Gulf in Izmit-Kocaeli, *Journal of Food, Agriculture and Environment* 7(2): 755-758.
- Madueño A, García D, Martínez J and Rubio C. 2006. Germination and development of bean seedlings (*Rhynchosia Minima* L.) Dc in salinity conditions. *Idem* 24(1): 47-54.

- Mallarino A and Wittry D. 2004. Efficacy of grid and zone soil sampling approaches for site-specific assessment of phosphorus, potassium, pH, and organic matter. *Precision Agriculture* 5: 131-144.
- Martínez E, Rodríguez L and Medina N. 2008. Feasibility study of a packing plant for bulb onion (*Allium cepa* L.) in the Irrigation District of Alto Chicamocha (Boyacá). *Colombian Journal of Horticultural Sciences* 8(2): 287-301.
- Mata I, Rodríguez M, López J and Vela G. 2014. Dynamics of salinity in soils. *Digital Magazine of the Man and its Environment Department* 1(5): 26-35.
- Mercado T, Ortega E, Arenas A and Combatt E. 2011. Salt dynamics in the irrigation district La Doctrina, Colombia. *IDESIA (Chile)* 29(1): 83-90. doi: 10.4067/S0718-34292011000100011
- Moreno M and Fечи Y. 2006. *Geology and Geomorphology. Formulation of the Management Plan and Environmental Management of the Alta Chicamocha River Basin*. Regional Autonomous Corporation of Boyacá, Universidad Pedagógica y Tecnológica de Colombia for Economic Studies, National University of Colombia-Institute of Environmental Research. 108 p.
- Narváez H, Bustamante I and Combatt E. 2014. Estimation of salinity in delta soils of the Sinú River in Colombia, using multiple linear regression models. *Idesia (Arica)* 32: 81-90. doi: 10.4067/S0718-34292014000300011
- Olías J, Cerón C and Fernández I. 2005. On the use of the classification of irrigation waters of the U.S. Laboratory Salinity (USLS Geogaceta, 37).
- Pacheco M, Cusba M and Torres S. 2004. Strategic marketing plan for the horticultural chain in the Irrigation District of Alto Chicamocha.
- Porras C. 2010. Clean development mechanism applied to the generation of electric power, with the use of geothermal energy of low enthalpy in oil production fields of Apiay, Suria and Campo, Castilla (Departamento del Meta, Colombia). Research Master in Design and Project Management. Universidad de La Sabana, Escuela de Engineering. Bogotá.
- POT 2012. Land Management Plan Municipality of Paipa. Biophysical subsystem (natural resources and the environment). 97p. In: [http://cdim.esap.edu.co/BancoMedios/Documentos%20PDF/pot%20-%20paipa%20-%202\\_biofisico%20\(97%20pag%20-%202230%20kb\).pdf](http://cdim.esap.edu.co/BancoMedios/Documentos%20PDF/pot%20-%20paipa%20-%202_biofisico%20(97%20pag%20-%202230%20kb).pdf). Query: April, 2017.
- Qadir M, Oster J, Schubert S, Noble A and Sahrawat K. 2007. Phytoremediation of sodic and saline-sodic soils. *Advances in Agronomy* 96: 197-247. doi: 10.1016/S0065-2113(07)96006-X
- Ramírez J. 2011. General assessment of salinity and modeling of salinization risks in Valle del Cauca soils, practical use of the PLA model to evaluate the risk of salinization in irrigated lands in the flat part of the Valle del Cauca (Colombia). Universidad Nacional de Colombia Facultad de Ciencias Agropecuarias Coordinación General de Postgrados Palmira 145 p.
- Taiz L and Zeiger E. 2006. Stress by salinity. *Vegetal Physiology*, Edited by Universitat Jaume I, 1338.
- Ulset A, Ruiz M, Herrera J and Ponce De Leon D. 1998. A geostatistical method of soil salinity spacing site sample. *Geoderma* 86: 143-151.
- Rhoades J, Kandiah A, Mashali A, FAO. 1992. The use of saline waters for crop production. 145p. <http://www.fao.org/3/a-t0667e.pdf>
- 
-