

Androgenic studies in the production of haploids and doubled haploids in *Capsicum* spp.

Estudios androgénicos en la producción de haploides y doble haploides en *Capsicum* spp.

doi: 10.15446/rfnam.v73n1.76044

Manuel Alejandro Sánchez¹, Yacenia Morillo Coronado^{2*} and Ana Cruz Morillo Coronado³

ABSTRACT

Keywords:

Anther
Chromosomal duplication
In vitro
Microspore
Pepper

Capsicum spp. is a horticultural crop of agronomic interest and is considered the fourth most important vegetable in the world. It is an important nutritional and medicinal source, and its production generates employment in the tropics. In this species, the genetic variability is wide and with great potential, which has been exploited to generate outstanding varieties. Breeding programs seek different alternatives to accelerate the production of improved varieties with desirable agronomic characteristics. These objectives can be achieved with the production of haploid and doubled haploid plants via androgenesis or gynogenesis, being androgenesis the approach most used for paprika cultures. The purpose of this review is to present the results of different researches in obtaining haploids and doubled haploids in cultivars of *Capsicum* spp. and its impact on the genetic improvement of this crop.

RESUMEN

Palabras clave:

Antera
Duplicación cromosómica
In vitro
Microspora
Pimiento

Capsicum spp es un cultivo hortícola de interés agronómico y es considerado como el cuarto vegetal de mayor importancia en el mundo, al ser una importante fuente nutricional, medicinal y generadora de empleo en los trópicos. En esta especie la variabilidad genética es amplia y con gran potencial el cual ha sido aprovechado para generar variedades sobresalientes. En este sentido, los programas de mejoramiento buscan diferentes alternativas que conduzcan a la integración de nuevas técnicas para acelerar la producción de variedades mejoradas con características agronómicas deseables, tales objetivos se pueden lograr con la producción de plantas haploides y doble haploides, por vía androgénica o ginogenética, siendo el enfoque más utilizado la androgenesis para cultivos de pimentón. El propósito de esta revisión es dar a conocer los resultados de diferentes investigaciones en la obtención de haploides y doble haploides en cultivares de *Capsicum* spp. y su impacto en el mejoramiento genético de este cultivo.

¹ Biotechnological Engineer. Independent Professional. Manizales, Colombia.

² Universidad de Caldas. AA 275. Manizales, Colombia.

³ Universidad Pedagógica y Tecnológica de Colombia. CP 15003. Tunja, Colombia.

* Corresponding author: <ymorillo@agrosavia.co>

The genus *Capsicum* spp. ($2n=2x=24$) belongs to the Solanaceae family, concentrated mainly in tropical and subtropical regions of America (López-España *et al.*, 2016). It comprises a set of more than thirty species. *Capsicum annum*, *C. chinense*, and *C. frutescens* stand out (Toquica *et al.*, 2003) as the most cultivated because of their high content of vitamins A, C, and Calcium (Nuez Viñals *et al.*, 1996). In Colombia, there is a growing demand for *Capsicum* species due to their good profitability, production, and export possibilities (Medina *et al.*, 2006). The main producing departments of this cultivar are Valle del Cauca, Santander, Antioquia, Córdoba, and Bolívar – being the latter the largest producer with 1.8 million t in an area of 24,700 ha (Agronet, 2010).

The genetic improvement of this cultivar occurs mainly through conventional techniques, thus generating genotypes with valuable genes related to production and resistance to biotic and abiotic stresses (Regner, 1996). However, there are inherent limitations to sexual reproduction: incompatibility and imprecision. Besides, obtaining pure or homozygous lines is time-consuming since they require at least six cycles of self-fertilization (Cravero *et al.*, 2011). In this context, the production of haploid and doubled haploid plants *in vitro* is a useful tool since pure lines can be obtained in one generation, thus reducing time and production costs (Maraschin *et al.*, 2005). As an advantage, doubled haploid plants have a great application in genetic and cytogenetic studies (Dwivedi *et al.*, 2015).

Haploid plants are those that have the number of gametophytic chromosomes of their progenitors (Das *et al.*, 2018). Doubled haploid plants are produced through the duplicated chromosome of the haploids (Dwivedi *et al.*, 2015), either spontaneously or artificially induced by antimitotic agents (Badu *et al.*, 2017). After chromosomal duplication, the haploids become 100% homozygous and can be used directly as parental lines in the production of F_1 lines (Bal and Abak, 2007).

Given the importance of haploids and doubled haploids in the improvement of crops, different artificial strategies have been developed for their production, such as: chromosomal elimination (bulbosum method), hormonal treatment, heat shock methods, induction of gynogenesis

(culture of ovaries), and androgenesis involving anthers and microspores, the latter being the most commonly used in pepper cultivars (Das *et al.*, 2018).

Several studies have been focused on the conventional improvement of *Capsicum* spp. However, new research has emerged using *in vitro* techniques in the induction of whole plants from androgenesis to generate doubled haploids, which is used to support breeding programs. Therefore, the objective of the present review is to present some scientific achievements in the production of haploids and doubled haploids via androgenesis in pepper crops. The information is useful for those genetic improvement programs that seek to apply techniques that increase the efficiency in obtaining cultivars, shortening the time required to obtain them, and maximizing the use of resources.

GENERALITIES OF HAPLOIDS AND DOUBLED HAPLOIDS INBREEDING

Importance of haploids and doubled haploids

The constant agricultural evolution and the high demand from the growing population for food with nutritional value force improvement programs to accelerate the production of improved varieties with desirable agronomic properties. The main purpose is to seek alternatives that lead to the integration of new methodologies and technologies, such as the production of doubled haploid plants (Barro *et al.*, 2001; Achar, 2002).

The doubled haploid plants make up a very useful tool for the breeder in the generation of new and improved varieties (Tuvesson *et al.*, 2000). Currently, one of the main drawbacks of conventional breeding programs for autogamous species is the long period needed to produce an improved line. Due to several generations of self-pollination are required to achieve homozygous genotypes. Once the new genotypes are obtained, the agronomic and industrial traits of interest are consecutively evaluated until they can finally be delivered to farmers (Jacquard *et al.*, 2009).

Biotechnological techniques involving *in vitro* culture of anthers, isolated microspores, ovaries, etc., can be used with the same efficacy as conventional methods of self-fertilization or backcrossing (Guzy-Wrobelka and Szarejko, 2003). With the *in vitro* techniques, it

is possible to develop resistant cultivars to various plant diseases; saving time, labor, and space in the experimental field (Wędzony *et al.*, 2009). The evaluation of these methodologies will constitute an important tool for the genetic improvement and for obtaining new varieties with excellent agronomic attributes so that they can be distributed to the farmers in less time.

Brief history of anther culture for haploid and doubled haploid production

Several reports are concluding that duplicate haploids can accelerate the production of new plant varieties of agronomic interest, being them applied in different species (Thomas *et al.*, 2003). Among the most outstanding crops are barley (*Hordeum vulgare* L.), pepper (*Capsicum annuum* L.), rice (*Oryza sativa* L.), tobacco (*Nicotiana tabacum* L.), and wheat (*Triticum aestivum* L) (Ferrie, 2007).

The first report on the use of haploid lines in pepper was published by Christensen and Bamford (1943), finding spontaneous haploid plants among the evaluated material. Similarly, in 1945, Tool and Bamford obtained for the first time doubled haploid plants in cultivars of *Capsicum* L. using colchicine. Subsequently, Campos and Morgan (1958) succeeded in inducing haploids by intraspecific crossing (Dumas de Vaulx and Pochard, 1974). The yield of these haploid lines was similar to their parents, although with a much lower percentage of fertility; these investigations opened the way to use haploids in plant breeding. The next step in the use of haploids was the development of biotechnological techniques such as the cultivation of anthers, developed in the early seventies. Wang *et al.* (1973) and George and Narayanaswamy (1973) were the first to report the successful regeneration of haploids in pepper from anther culture (Harn *et al.*, 1975). These researches showed embryonic induction, but a successful regeneration of plants was rarely obtained.

In the works carried out by Sibi *et al.* (1979) and Dumas de Vaulx *et al.* (1981) in *Capsicum*, using the cultivation of anthers, the purpose was to make the technique more efficient in terms of obtaining regenerated plants by making certain modifications. Thus, for instance, the changes made by Sibi *et al.* (1979) consisted of pre-treatment of anthers in cold (4 °C, 48 h) and

determination of the state of optimal development of the microspore (medium or late uninucleate), which allowed them to obtain seedlings for each isolated anther. On the other hand, Dumas de Vaulx *et al.* (1981) exposed the anthers to thermal shock of 35 °C during eight days of darkness, obtaining between 5-10% of regenerated plants of pepper genotypes.

Dolcet-Sanjuan *et al.* (1997) describe a new protocol using a biphasic system of semi-solid and liquid medium, replacing sucrose with maltose and enriching the atmosphere of the anther culture medium with CO₂ for androgenesis in pepper hybrids, where they managed to increase the number of embryos (up to 3,561 per 100 flowers) and plants (up to 23 per 100 flowers) with 65% of plants duplicated spontaneously. Time later, Yin *et al.* (2010) confirm that pretreatment at low temperatures, the combinations of growth regulators, the concentrations of activated carbon, and pre-culture temperatures are critical factors that affect the formation of androgenic embryos.

Over the years, research aimed at the use and production of doubled haploids has focused on the refinement of *in vitro* culture. It has been seen that the optimization of *in vitro* factors such as quality of the medium, light intensity, temperature, nutrient replacement, among others, improve the androgenic response of the obtained *in vitro*-plants. However, it has been shown that this response is highly influenced by its different genotypes (Morrison *et al.*, 1986).

Despite the advantages offered by the aforementioned processes, there are numerous reports that argue the recalcitrance of the genus *Capsicum* to *in vitro* morphogenesis, which manifests itself in various ways such as the low efficiency of the regeneration systems as well as the low reproducibility of the protocols of regeneration –the high index of deformed somatic embryos, the low rate of germination and/or conversion of somatic embryos into plants (Steinitz *et al.*, 2003). Besides the above, a protocol for a specific genotype for the production of doubled haploids has not been fully established (Irikova *et al.*, 2011). The most critical steps are the production of structures similar to the embryos and the subsequent generation of the plant, the collection of flower buds in the optimal state, stress

treatments, in addition to the environment and the culture conditions (Irikova *et al.*, 2011).

COMPOSITION OF CULTURE MEDIA AND CULTURE CONDITIONS

Basal media are the main culture components since these contain the nutritional requirements for an embryogenic response (Koleva-Gudeva *et al.*, 2007). The most used media for haploid and doubled haploid induction are N6, MS (Murashige and Skoog, 1962), NN (Nitsch and Nitsch), B5, CP (Dumas de Vaulx *et al.*, 1981), among others, including the modified versions. Moreover, sucrose, maltose, and mannitol are the main sources of carbohydrates in these media, due to their nutritional and osmotic effects (Powell, 1990; Thompson *et al.*, 1986). Taşkin *et al.* (2011) found a higher embryo production and formation in MS media after comparing four types of culture using pepper species. However, other studies using MS and CP basal media for embryogenic formation from anthers found that CP was the adequate medium since normal embryos of *Capsicum annuum* L. were formed in all cases (Luitel and Kang 2013). Interestingly, although basal media are important for achieving a positive androgenic response, most researchers conclude that the frequency of callus and embryo formation depends on the genotype.

On the other hand, growth regulators (e.g., auxins, cytokines, among others) play a fundamental role by greatly affecting the development of the microspore and promoting cell division in embryos and calli (Chen, 1986). Olszewska *et al.* (2014) explored the effect of anther age and two concentrations of kinetin in the regeneration medium. The modifications assessed by the authors led to a greater androgenic response in *Capsicum* genotypes compared to the traditionally used protocols for this species. Similar findings were reported by Cheng *et al.* (2013) while studying the effect of growth regulators and activated carbon on embryogenesis induction in *C. annuum* L. The authors concluded that the concentration of activated carbon could act as a promoter of embryogenesis in microspore culture. In particular, the addition of antioxidants and activated carbon is often useful with some genotypes since it reduces tissue blackening caused by phenols. However, Vélez Torres *et al.* (2010) assessed the effect of different concentrations of growth regulators

on pepper cultivars and concluded that exogenous regulators determine the successful formation of embryogenesis, while the endogenous hormone levels define the response *in vitro*.

It has been shown that stress pre-treatments (which vary considerably among species) applied to anthers and microspores induce the sporophytic pathway and inhibit the gametophytic pathway (development of fertile pollen) (Germanà, 2011) and promote the embryogenic potential. Pre-treatments, such as thermal shock (cold, hot), high humidity, water stress, nitrogen deprivation, ethanol, among others, are the most used for the androgenic pathways (Shariatpanahi *et al.*, 2006). Ercan and Ayar Şensoy (2011) studied the androgenic response in pepper cultivars after a pre-treatment with cold (4 °C) and darkness for 24 hours and obtained 44 embryos from 2398 anthers in *in vitro* culture. Later, Popova *et al.* (2016) assessed the embryogenic response of *C. annuum* L. to different lengths of a pre-treatment of low temperature and darkness (during the first eight days). The results indicated that embryogenic efficiency decreases at low temperatures and under photoperiod inhibition. Several studies have shown that the exposure of androgenic cultures to alternating light periods (14/10 h; light/dark; 2,000 lux m⁻²) generates an artificial environmental signal that regulates pollen morphogenesis *in vitro* (Reynolds and Crawford, 1997).

MAIN METHODS FOR HAPLOID AND DOUBLED HAPLOID INDUCTION IN *Capsicum* spp.

There are several techniques for obtaining haploids - *in vivo* and *in vitro* methodologies (Figure 1). The most used methods are gynogenesis and androgenesis (Irikova *et al.*, 2016). Gynogenesis involves female gametophytes (ovules and ovaries), and androgenesis uses male gametes (anthers and microspores) to generate haploids. Gynogenesis is the least favored technique due to its low efficiency (Forster *et al.*, 2007). On the other hand, male gametogenesis is one of the most used methods for haploid and doubled haploid production (Kasha and Maluszynski, 2003), and it has been successfully used in more than 200 species, mainly annual plants, including pepper (Mitykó *et al.*, 2006).




Figure 1. Haploid and doubled haploid production methods.

Anther culture

There is no specific protocol or culture condition for *Capsicum* spp. to induce androgenesis *in vitro*. However, some common guidelines are provided for plant formation from anther culture (Figure 2). In general, most studies report an optimal floral bud size between 3 and 6 cm in diameter. The buds are washed with distilled water, and surface disinfection is performed with calcium hypochlorite, 70% ethanol, Tween 20, among others. After determining the developmental stage of the microspore through staining with acetocarmine, Schiff reagent or DAPI, among others, those in early or late uninucleate phases are cultured in induction media with plant growth regulators (2,4D, dichlorophenoxyacetic acid; KIN, kinetin, among others) and maintained under a pre-treatment at approximately 28-35 °C with light (12-18 h) and dark (6-12 h) periods. Following embryo formation, these cultures are transferred to regeneration media (free of growth hormones) to stimulate the formation of roots and shoots. The entire process is conducted under aseptic conditions in a laminar flow cabinet (Reinert and Bajaj, 1977). The most used methods for ploidy analysis of *in vitro* plants are cytological techniques involving chloroplast counts in guard cells, chromosome counts in

somatic cells from root apices, as well as flow cytometry. Besides, isoenzyme and DNA marker-based techniques are also used (Germanà, 2011).

Microspore culture

Similar to the protocol for anther culture, there is no standard condition to induce plant formation from microspores. The explant can come from microspores or immature pollen to obtain haploid plants, with the advantage that embryos can be obtained from heterozygote somatic tissue (Regner, 1996). This methodology is similar to the one for anther culture, which begins with the mechanic isolation of microspores from a single anther using a glass rod. The microspores are then cultured in an isolation medium containing sucrose, ascorbic acid, L-proline, biotin, and nicotinic acid, according to most studies. The microspore is generally suspended using a nylon net and then transferred to liquid or solid culture media without growth hormones (regeneration or multiplication media). Finally, a haploid plant is obtained and, subsequently, a double haploid plant by treatment with colchicine. This protocol is performed under sterile conditions (Das *et al.*, 2018).




Figure 2. General scheme for the production of haploid and doubled haploid plants from anther/microspore cultures.

The anther culture technique is often the method of choice by most researchers for the production of doubled haploid. The advantage of this procedure is that the nutritional requirements of the anthers are much simpler than those of the isolated microspores (Bajaj, 1990), and it is not as laborious as microspore culture. Despite these advantages, there are several limitations, such as the regeneration of anther wall somatic tissue, the production of mixoploids (in somatic and gametic tissue), and the formation of asynchronous microspores (Kott *et al.*, 1988). Besides, the use of grains in advanced developmental stages is challenging since these suppress the androgenic capacity of the younger grains by releasing toxic substances (Bhojwani and Razdan, 1996).

Another determining factor for the success of anther culture is the selection of the floral buds since pollen in the early uninucleate phase is more sensitive to

external stress, and this could hinder embryogenesis (Irikova *et al.*, 2011). According to Kim *et al.* (2004), the optimal anther phase for embryo production is the early binucleate phase. Likewise, Supena *et al.* (2006a) determined that a critical factor for *C. annuum* L. cultivars is the selection of microspores in the late unicellular phase.

The difficulties discussed can be overcome by culturing isolated microspores. This technique allows not only to discard the use of somatic tissue (i.e., the unknown effect of the anther wall) (Mishra and Goswami, 2014) but also to isolate and culture microspores in an adequate phase; thus, contributing to an efficient regeneration. In particular, there are thousands of microspores per anther, which allows obtaining a greater number of haploid plants, and this can be applied at a large scale on a variety of genotypes (Taşkin *et al.*, 2011). Supena *et al.* (2006b) found a greater rate of spontaneous re-diploidization from

isolated microspores compared to cultured anther. These results prompted studies such as the one by Lantos *et al.* (2009), in which isolated microspore cultures in pepper genotypes were improved by applying wheat ovary co-culture. Similarly, Heidari-Zefreh *et al.* (2018) improved the efficacy of microspore embryogenesis and, subsequently, the regeneration frequency of *C. annuum* L. seedlings by applying different concentrations of ascorbic acid in the medium along with a thermal shock. Despite the success of these studies, there is limited available research regarding isolated microspore culture in *C. annuum* L., mainly since this technique requires more suitable equipment and technical-scientific skills.

Chromosomal duplication of haploid plants

The research focused on the rapid development of improved cultivars using *in vitro* techniques usually requires first obtaining haploid plants for them to be later diploidized (duplication of chromosomes). Seedlings derived by androgenesis lead to the spontaneous formation of haploids and doubled haploids; however, the percentage of the latter is very low. Due to this fact, it has become necessary to resort to chemical substances, including colchicine ($C_{22}H_{25}NO_6$), widely used for the induction of polyploidy in plants (Urwin, 2014). This substance is an alkaloid extracted from *Colchicum autumnale*, which inhibits the self-assembly of tubulin, preventing the formation of the microtubules of the spindle, affecting only cells that are in the division, therefore acting as a "mitotic poison" (Badu *et al.*, 2017). The most important advantage of polyploidy is that plants tend to have better performance and morphological characteristics, such as the height and size of plant organs (Hannweg *et al.*, 2016), and the increase in biomass in general (Urwin, 2014).

This mutagenic can be applied to plants in different ways: by aqueous solution submerging the roots (systemic absorption) or by microinjection, in which the solution is incorporated into the *in vitro* plant by mechanical inoculation. On the other hand, calli can also be imbibed in colchicine before they are planted in the regeneration medium (Dhooghe *et al.*, 2011).

A large number of polyploid induction processes have been reported with colchicine, being a very effective mechanism, but it has also been seen that its use can

generate chimerization and death of the tissue with which it has been in contact, so the production of viable polyploid plants decreases in a large proportion (do Régo *et al.*, 2009). Because colchicine has negative effects, the use of Oryzalin and other methods that affect mitotic processes, have begun to be used, generating higher rates of viable polyploid seedlings.

Techniques for the chromosomal duplication of haploids in pepper cultivars are generally based on treatment with colchicine. Gémesné Juhász *et al.* (2001a, 2001b) showed that they could increase the efficiency in 95% of doubled haploid production in pepper cultivars (*Capsicum annuum* L.) by applying colchicine (0.04%) to the regeneration medium. In subsequent investigations, Gémesné Juhász *et al.* (2001a) indicated that the medium supplemented with colchicine resulted in an effective diploidization of 50-95% explants.

FINAL CONSIDERATIONS

The biotechnology techniques and advancements in molecular biology have enabled the development of useful tools for plant improvement. Among these, the production of haploid and doubled haploid plants is relevant to conventional improvement since it reduces the time required to obtain new varieties. Doubled haploid plants constitute genetically pure lines, which do not exist in conventional cultures and allow to rapidly fix new gene combinations resulting in individuals with greater production, resistance to pests and diseases, and higher tolerance to biotic and abiotic factors. However, the application of these tools must still overcome several key factors, including recalcitrant plants, genotype dependence, among others, that challenge the production of a suitable percentage of endogamic lines. Besides, a better understanding of the cellular, biochemical, and molecular basis of embryo induction through the androgenic pathway is required. Consequently, the current protocols need improvement, and new genotype-independent strategies should be implemented. The processes involved in microspore-induced embryogenesis should be further investigated, mainly, the cell division processes for obtaining haploid plants. The haploid induction technique can be efficiently combined with other plant biotechnology techniques, resulting in important achievements in genetic studies, gene mapping, QTL localization,

genomics, and mutagenesis and genetic transformation research. Overall, there is a clear interest in this tool for plant improvement and its future application in several agriculturally important crops.

REFERENCES

Achar PN. 2002. A study of factors affecting embryo yields from anther culture of cabbage. *Plant Cell, Tissue and Organ Culture* 69(2): 183-188. doi: 10.1023/A:1015232008585

AGRONET. 2010. Análisis-Estadísticas, producción nacional por producto. En: <https://www.agronet.gov.co/Paginas/ProduccionNacionalProducto.aspx>; consulta: octubre 2018.

Badu M, Tripathy B, Sahu GS and Jena AK. 2017. Role of doubled haploids in vegetable crop improvement. *Journal of Pharmacognosy and Phytochemistry* 6(6): 384-389.

Bajaj YPS. 1990. *In vitro* production of haploids and their use in cell genetics and plant breeding. pp 1-44. In: Bajaj YPS (ed.). *Biotechnology in agriculture and forestry. Part I. Haploids in crop improvement*. Vol. 12. Springer, Berlin.

Bal U and Abak K. 2007. Haploidy in tomato (*Lycopersicon esculentum* Mill.): a critical review. *Euphytica* 158(1-2): 1-9. doi: 10.1007/s10681-007-9427-1

Barro F, Fernandez-Escobar J, De La Vega M and Martin A. 2001. Doubled haploid lines of *Brassica carinata* with modified erucic acid content through mutagenesis by EMS treatment of isolated microspores. *Plant Breeding* 120(3): 262-264. doi: 10.1046/j.1439-0523.2001.00602.x

Bhojwani SS and Razdan MK. 1996. *Plant tissue culture: Theory and practice, a revised edition*. Vol. 5. First edition. Elsevier, Amsterdam. 776 p.

Campos FF and Morgan DT. 1958. Haploid pepper from a sperm, an androgenetic of *Capsicum frutescens*. *Journal of Heredity* 49(4): 134-137. doi: 10.1093/oxfordjournals.jhered.a106786

Chen Z. 1986. Induction of androgenesis in woody plants. pp 42- 66. In: Han H and Hongyuan Y (eds.). *Haploids in higher plants in vitro*. China Academic Publishers, Springer-Verlag.

Cheng Y, Ma RL, Jiao YS, Qiao N and Li TT. 2013. Impact of genotype, plant growth regulators and activated charcoal on embryogenesis induction in microspore culture of pepper (*Capsicum annuum* L.). *South African Journal of Botany* 88: 306-309. doi: 10.1016/j.sajb.2013.08.012

Christensen H and Bamford R. 1943. Haploids in twin seedlings of pepper *Capsicum annuum* L. *Journal of Heredity*. 34: 99-104.

Cravero VP, López Anido FS, Espósito MA y Cointry EL. 2011. Mejoramiento convencional y no convencional de especies hortícolas. *BAG Journal of Basic and Applied Genetics* 22(1): 1-4.

Das A, Kumar K, Tribhuvan K, Das SS and Mishra M. 2018. Development of haploid and double haploid in fruit crops-A review. *International Journal of Current Microbiology and Applied Sciences* 7(5): 2119-2132. doi: 10.20546/ijcmas.2018.705.247

Dhooghe E, Van Laere K, Eeckhaut T, Leus L and Van Huylenbroeck J. 2011. Mitotic chromosome doubling of plant tissues *in vitro*. *Plant Cell, Tissue and Organ Culture* 104(3): 359-373. doi: 10.1007/s11240-010-9786-5

do Rêgo MM, do Rêgo ER, Otoni WC and Bruckner CH. 2009. Epigenetic effects in induced *in vitro* tetraploids passion fruit (*Passiflora edulis* Sims.). *Acta Horticulturae* 829: 167-175. doi: 10.17660/ActaHortic.2009.829.24

Dolcet-Sanjuan R, Claveria E and Huerta A. 1997. Androgenesis in *Capsicum annuum* L. effects of carbohydrate and carbon dioxide enrichment. *Journal of the American Society for Horticultural Science* 122(4): 468-475.

Dumas de Vaulx R and Pochard E. 1974. Essai d'introduction de la parthénogénèse haploïde par action du protoxyde d'azote sur les fleurs de piments (*Capsicum annuum*). *Annales de l'amélioration des plantes* 24: 283-305.

Dumas de Vaulx R, Chambonnet D and Pochard E. 1981. *In vitro* culture of pepper (*Capsicum annuum* L.) anthers: High rate plant production from different genotypes by +35 °C treatment. *Agronomie* 1: 859-864.

Dwivedi SL, Britt AB, Tripathi L, Sharma S, Upadhyaya HD and Ortiz R. 2015. Haploids: constraints and opportunities in plant breeding. *Biotechnology Advances* 33(6): 812-829. doi: 10.1016/j.biotechadv.2015.07.001

Ercan N and Ayar Şensoy F. 2011. Androgenic responses of different pepper (*Capsicum annuum* L.) cultivars. *Biyoloji Bilimleri Araştırma Dergisi* 4(2): 59-61.

Ferrie AMR. 2007. Doubled haploid production in nutraceutical species: a review. *Euphytica* 158(3): 347-357. doi: 10.1007/s10681-006-9242-0

Forster BP, Herberle-Bors E, Kasha KJ and Touraev A. 2007. The resurgence of haploids in higher plants. *Trends in Plant Science*. 12(8):368-375. doi: 10.1016/j.tplants.2007.06.007

Gémesné Juhász A, Petus M, Venczel G, Zatykó L, Gyulai G and Csepő M. 2001a. Genetic variability of anther donor versus spontaneous doubled haploid descendants and colchicine induced doubled haploid sweet pepper (*Capsicum annuum* L.) lines. *Acta Horticulturae* 560: 149-152. doi: 10.17660/ActaHortic.2001.560.24

Gémesné Juhász J, Petu M, Venczel G, Sagi Z and Zatyko L. 2001b. Colchicine, an efficient genome doubling agent for anther derived haploid paprika (*Capsicum annuum* L.) plants. In: 11th EUCARPIA Meeting Genetics and Breeding of *Capsicum* and Eggplant. Antalya, Turkey. 146 p.

George L and Narayanaswamy S. 1973. Haploid *Capsicum* through experimental androgenesis. *Protoplasma* 78(4): 467-470. doi: 10.1007/BF01275781

Germaná MA. 2011. Anther culture for haploid and doubled haploid production. *Plant Cell, Tissue and Organ Culture* 104(3): 283-300. doi: 10.1007/s11240-010-9852-z

Guzy-Wroblewska J and Szarejko I. 2003. Molecular and agronomic evaluation of wheat doubled haploid lines obtained through maize pollination and anther culture methods. *Plant Breeding* 122(4): 305-313. doi: 10.1046/j.1439-0523.2003.00858.x

Hannweg K, Visser G, de Jager K and Bertling I. 2016. *In vitro*-induced polyploidy and its effect on horticultural characteristics, essential oil composition and bioactivity of *Tetradenia riparia*. *South African Journal of Botany* 106: 186-191. doi: 10.1016/j.sajb.2016.07.013

Harn C, Kim MZ, Choi KT and Lee YI. 1975. Production of haploid callus and embryoid from the cultured anther of *Capsicum annuum*. *Sabrao Journal* 7: 71-77.

Heidari-Zefreh A, Shariatpanahi M, Mousavi A and Kalatejari S. 2018. Enhancement of microspore embryogenesis induction and plantlet regeneration of sweet pepper (*Capsicum annuum* L.)

using putrescine and ascorbic acid. *Protoplasma* 256(1): 13-24. doi: 10.1007/s00709-018-1268-3

Irikova T, Grozeva S and Rodeva V. 2011. Anther culture in pepper (*Capsicum annuum* L.) *in vitro*. *Acta Physiologiae Plantarum* 33(5): 1559-1570. doi: 10.1007/s11738-011-0736-6

Irikova T, Kintzios S, Grozeva S and Rodeva V. 2016. Pepper (*Capsicum annuum* L.) anther culture: fundamental research and practical applications. *Turkish Journal of Biology* 40(4): 719-726. doi: 10.3906/biy-1506-79

Jacquard C, Mazeyrat-Gourbeyre F, Devaux P, Boutilier K, Baillieul F and Clément C. 2009. Microspore embryogenesis in barley: anther pretreatment stimulates plant defence gene expression. *Planta* 229(2): 393-402. doi: 10.1007/s00425-008-0838-6

Kasha KJ and Maluszynski M. 2003. Production of doubled haploids in crop plants. An introduction. pp. 1-4. In: Maluszynski M, Kasha KJ, Forster BP and Szarejko I (eds.). *Doubled haploid production in crop plants: A Manual*. Springer, Dordrecht. 428 p. doi: 10.1007/978-94-017-1293-4_1.

Kim M, Kim J, Yoon M, Choi DI and Lee KM. 2004. Origin of multicellular pollen and pollen embryos in cultured anthers of pepper (*Capsicum annuum*). *Plant Cell, Tissue and Organ Culture* 77(1): 63-72. doi: 10.1023/B:TICU.000016506.02796.6a

Koleva-Gudeva LR, Spasenoski M and Trajkova F. 2007. Somatic embryogenesis in pepper anther culture: The effect of incubation treatments and different media. *Scientia Horticulturae* 111(2): 114-119. doi: 10.1016/j.scienta.2006.10.013

Kott LS, Polsoni L, Ellis B and Beversdorf WD. 1988. Autotoxicity in isolated microspore cultures of *Brassica napus*. *Canadian Journal of Botany* 66(8): 1665-1670. doi: 10.1139/b88-227

Lantos C, Juhász AG, Somogyi G, Ötvös K, Vági P, Mihály R, Kristóf Z, Somogyi N and Pauk J. 2009. Improvement of isolated microspore culture of pepper (*Capsicum annuum* L.) via co-culture with ovary tissues of pepper or wheat. *Plant Cell, Tissue and Organ Culture* 97(3): 285-293. doi: 10.1007/s11240-009-9527-9

López-España RG, Hernández-Verdugo S, Parra-Terraza S, Porras F, Pacheco-Olvera A, Valdez-Ortiz A and Muy-Rangel MD. 2016. Diferenciación geográfica de poblaciones de chile silvestre (*Capsicum annuum* L. var. *glabriusculum*) del norte de México. *Phyton* 85(1): 131-141.

Luitel BP and Kang WH. 2013. *In vitro* androgenic response of minipaprika (*Capsicum annuum* L.) genotypes in different culture media. *Horticulture, Environment, and Biotechnology* 54(2): 162-171. doi: 10.1007/s13580-013-0110-2

Maraschin SF, de Priester W, Spaink HP and Wang M. 2005. Androgenic switch: an example of plant embryogenesis from male gametophyte perspective. *Journal of Experimental Botany* 56(417): 1711-1726. doi: 10.1093/jxb/eri190

Medina CI, Lobo M y Gómez A. 2006. Variabilidad fenotípica en poblaciones de ají y pimentón de la colección colombiana del género *Capsicum*. *Corpoica Ciencia y Tecnología Agropecuaria* 7(2): 25-39.

Mishra VK and Goswami R. 2014. Haploid production in higher plant. *International Journal of Biological and Chemical Sciences* 1(1): 26-45.

Mitykó J and Gémes Juhász A. 2006. Improvement in the haploid technique routinely used for breeding sweet and spice peppers in Hungary. *Acta Agronomica Hungarica* 54(2): 203-219. doi: 10.1556/AAgr.54.2006.2.8

Morrison RA, Koning RE and Evans DA. 1986. Anther culture of an interspecific hybrid of *Capsicum*. *Journal of Plant Physiology* 126(1): 1-9. doi: 10.1016/S0017-1617(86)80210-3

Murashige T and Skoog F. 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiologia Plantarum* 15(3): 473-497. doi: 10.1111/j.1399-3054.1962.tb08052.x

Nuez Viñals F, Gil Ortega R and Costa García JC. 1996. *El cultivo de pimientos, chiles y ajíes*. Mundiprensa Libros SA, Madrid. 607 p.

Olszewska D, Kisiala A, Niklas-Nowak A and Nowaczyk P. 2014. Study of *in vitro* anther culture in selected genotypes of genus *Capsicum*. *Turkish Journal of Biology* 38(1): 118-124. doi: 10.3906/biy-1307-50

Popova T, Grozeva S, Todorova V, Stankova G, Anachkov N and Rodeva V. 2016. Effects of low temperature, genotype and culture media on *in vitro* androgenic answer of pepper (*Capsicum annuum* L.). *Acta Physiologiae Plantarum* 38(11): 1-11. doi: 10.1007/s11738-016-2294-4

Powell W. 1990. Environmental and genetic aspects of pollen embryogenesis. pp. 44-65. In: Bajaj YPS (ed) *Biotechnology in agriculture and forestry. Part I. Haploids in crop improvement*. Vol. 12. Springer, Berlin.

Regner F. 1996. Anther and microspore culture in *Capsicum*. pp. 77-89. In: Jain SM, Sopory SK and Veilleux RE (eds). *In vitro haploid production in higher plants*. Springer, Dordrecht. 414 p. doi: 10.1007/978-94-017-1858-5_5

Reinert J. 1977. Chapter II. Haploid. pp. 249-340. In: Reinert J and Bajaj YPS (eds.). *Applied and fundamental aspects of plant cell, tissue and organ culture*. Springer, Berlin-Heidelberg-New York. 803 p.

Reynolds TL and Crawford RL. 1997. Effects of light on the accumulation of abscisic acid and expression of an early cysteine-labeled metallothionein gene in microspores of *Triticum aestivum* during induced embryogenic development. *Plant Cell Reports* 16(7): 458-463. doi: 10.1007/BF01092766

Shariatpanahi ME, Belogradova K, Hessamvaziri L, Heberle-Bors E and Touraev A. 2006. Efficient embryogenesis and regeneration in freshly isolated and cultured wheat (*Triticum aestivum* L.) microspores without stress pre-treatment. *Plant Cell Reports* 25(12): 1294-1299. doi: 10.1007/s00299-006-0205-7

Sibi M, Dumas de Vaulx R and Chambonnet D. 1979. Obtention de plantes haploïdes par androgenèse *in vitro* chez le Piment (*Capsicum annuum* L.). *Annales de l'amélioration des plantes* 29(5): 583-606.

Steinitz B, Küsek M, Tabib Y, Paran I and Zelcer A. 2003. Pepper (*Capsicum annuum* L.) regenerants obtained by direct somatic embryogenesis fail to develop a shoot. *In vitro Cellular & Developmental Biology-Plant* 39(3): 296-303. doi: 10.1079/IVP2002405

Supena EDJ, Muswita W, Suharsono S and Custers JBM. 2006a. Evaluation of crucial factors for implementing shed-microspore culture of Indonesian hot pepper (*Capsicum annuum* L.) cultivars. *Scientia Horticulturae* 107(3): 226-232. doi: 10.1016/j.scienta.2005.08.006

Supena EDJ, Suharsono S, Jacobsen E and Custers JBM. 2006b. Successful development of a shed-microspore culture protocol for doubled haploid production in Indonesian hot pepper

(*Capsicum annuum* L.). *Plant Cell Reports* 25(1): 1-10. doi: 10.1007/s00299-005-0028-y

Taşkin H, Büyükalaca S, Keleş D and Ekbiç E. 2011. Induction of microspore-derived embryos by anther culture in selected pepper genotypes. *African Journal of Biotechnology* 10(75):17116-17121. doi: 10.5897/AJB11.2023

Thomas WTB, Forster BP and Gertsson B. 2003. Doubled haploids in breeding. pp 337-349. In: Maluszynski M, Kasha KJ, Forster BP and Szarejko I (eds.). *Doubled haploid production in crop plants, a manual*. Kluwer Academic Publishers, Dordrecht.

Thompson MR, Douglas TJ, Obata-Sasamoto H and Thorpe TA. 1986. Mannitol metabolism in cultured plant cells. *Physiologia Plantarum* 67(3): 365-369. doi: 10.1111/j.1399-3054.1986.tb05749.x

Toole M and Bamford R. 1945. The formation of diploid plants from haploid peppers. *Journal of Heredity* 36: 67-70.

Toquica SP, Rodríguez F, Martínez E, Duque MC and Tohme J. 2003. Molecular characterization by AFLPs of *Capsicum* germplasm from the Amazon department in Colombia. *Genetic Resources and Crop Evolution* 50(6): 639-647. doi: 10.1023/A:1024429320771

Tuvesson S, Ljungberg A, Johansson N, Karlsson KE, Suijs LW and Josset JP. 2000. Large-scale production of wheat and triticale doubled haploids through the use of a single anther culture method. *Plant Breeding* 119(6): 455-4594. doi: 10.1046/j.1439-0523.2000.00536.x

Unwin NAR. 2014. Generation and characterization of colchicine-induced polyploid *Lavandula x intermedia*. *Euphytica* 197(3): 331-339. doi: 10.1007/s10681-014-1069-5

Vélez Torres M, Robledo Paz A, Corona Torres T, Aguilar Rincón V, Ramírez Vallejo P and Suárez Espinosa J. 2010. Obtención de plantas haploides en chile miahuateco (*Capsicum annuum* L.). *Revista Mexicana de Ciencias Agrícola* 1(2): 187-199.

Wang YY, Sun CS, Wang CC and Chien NF. 1973. The induction of the pollen plants of *Triticale* and *Capsicum annuum* from anther culture. *Scientia Sinica* 16: 147-151. doi: 10.1360/ya1973-16-1-147

Wędzony M, Forster BP, Żur I, Golemic E, Szechyńska-Hebda M, Dubas E and Gotębiowska G. 2009. Progress in doubled haploid technology in higher plants. pp. 1-33. In: Touraev A, Forster BP and Jain SM (eds.). *Advances in Haploid production in Higher Plants*. Springer, Dordrecht. 348 p. doi: 10.1007/978-1-4020-8854-4_1

Yin T, Tian S, Luo S, Chen X, Wang Y and Shen S. 2010. Studies on isolated microspore embryoid induction of *C. annuum* L. *Frontiers of Agriculture in China* 4(4): 438-442. doi: 10.1007/s11703-010-1028-4