

Molecular prevalence of Bovine Leukemia Virus in specialized dairies in the department of Antioquia, Colombia

Prevalencia molecular del Virus de Leucosis Bovina en lecherías especializadas del departamento de Antioquia, Colombia

<https://doi.org/10.15446/rfnam.v76n2.104722>

Daniela Castillo Rey^{1*}, Albeiro López-Herrera¹ and Cristina Úsuga-Monroy¹

ABSTRACT

Keywords:

DNA
Enzootic bovine leukemia
Lymphocytes
Prevalence
Polymerase chain reaction
Retrovirus

Dairy production systems are a sensitive sector of the primary economy frequently affected by pathogens that negatively impact production parameters, the bovine leukemia virus (BLV) one of these. In this study, the molecular prevalence of BLV was determined in the specialized dairy sector of Antioquia using the viral marker of the envelope gene (*env*). Blood samples were taken from 575 bovines from specialized dairies in Antioquia distributed in 53 herds and located in the three specialized milk production areas of Antioquia (north, east, and Valle de Aburrá). DNA extraction was performed by salting out, and a nested PCR was performed to detect the *env* gene. The products were visualized on a 2% agarose gel with GelRed as an intercalator. A molecular prevalence of BLV of 17.0% in animals and 71.7% in herds were found, being Valle de Aburrá the area where the highest rate of positive animals was obtained (21.1%), unlike the northern area with the lowest rate (15.6%). The molecular prevalence of BLV in this study is lower than that of previous studies in the department, which ranged between 47 and 73%, and this may be associated with factors of breed resistance, the age of the animals, or management practices in the herds. These results can contribute to creating BLV control strategies and optimizing milk production in the department of Antioquia, being relevant to paying attention to the behavior of this pathogen under different production system conditions.

RESUMEN

Palabras clave:

ADN
Leucosis Bovina Enzoótica
Linfocitos
Prevalencia
Reacción en cadena de la polimerasa
Retrovirus

Los sistemas de producción de leche son un sector sensible de la economía primaria, viéndose frecuentemente afectados por patógenos que impactan negativamente parámetros productivos, siendo el virus de la leucosis bovina (BLV) uno de estos. En este estudio, se determinó la prevalencia molecular a BLV en la lechería especializada de Antioquia mediante el marcador viral del gen de la envoltura (*env*). Se tomó muestra de sangre de 575 bovinos de lecherías especializadas de Antioquia distribuidos en 53 hatos, ubicados en las tres zonas de producción de leche especializada de Antioquia (norte, oriente y Valle de Aburrá). Se realizó extracción de ADN por Salting out y se realizó una PCR anidada para detección del gen *env*. Los productos se visualizaron en un gel de agarosa al 2% con GelRed como intercalante. Se encontró una prevalencia molecular a BLV del 17,0% en animales y 71,7% en hatos, siendo el Valle de Aburrá la zona donde se obtuvo la mayor tasa de animales positivos (21,1%), a diferencia del norte que tuvo la más baja (15,6%). Se halló que la prevalencia molecular a BLV de este estudio es menor a la de investigaciones previas en el departamento, la cual se encontraba entre el 47 y 73%, y esto puede estar asociado a factores de resistencia raciales, edad de los animales o las prácticas de manejo en los hatos. Estos resultados pueden contribuir a crear estrategias de control del BLV y optimizar la producción lechera en el departamento de Antioquia, siendo relevante poner atención al comportamiento de este patógeno bajo las diferentes condiciones que tienen los sistemas de producción.

¹Grupo de Investigación Biodiversidad y Genética Molecular "BIOGEM", Universidad Nacional de Colombia sede Medellín, Colombia.
dcastillor@unal.edu.co , alherrera@unal.edu.co , cusugam@unal.edu.co 

*Corresponding author

The specialized milk production systems with bovines constantly face the presence of pathogens that affect the sanitary conditions and the economy of herds. One of these pathogens is bovine leukemia virus (BLV), whose target cells are B lymphocytes, causing enzootic bovine leukemia (EBL), which affects the immune system of animals causing immunosuppression and allowing entry to other pathogens (Bartlett et al. 2020; Marawan et al. 2021). Most animals are asymptomatic to the disease; only 30% of infected animals develop persistent lymphocytosis and only 1-5% develop lymphomas (Barez et al. 2015). In addition to affecting the health of animals, it generates a detriment in the quantity of milk produced and the compositional and sanitary quality of the milk, having adverse effects on the economy of dairy herds (Marawan et al. 2021; Úsuga-Monroy et al. 2018a).

The BLV belongs to the Retroviridae family; these viruses are characterized by transcribing their genome from RNA to DNA through the reverse transcriptase enzyme and inserting themselves into the host cell genome (Polat et al. 2017). When this occurs, the virus is renamed provirus; infections of this type are permanent. BLV transmission occurs through contact with fluids that have the presence of infected cells (blood, colostrum, and milk) from an infected animal to another healthy one, for which good management practices in production systems are vital to prevent virus transmission (Bartlett et al. 2014; Ruiz et al. 2018). It has also been found that it can be transmitted in the womb and at the time of delivery (Kuczewski et al. 2021).

The viral genome consists of four primordial genes that code for structural proteins and enzymes (*env*, *gag*, *pol*, and *pro*). In addition, it has two regulatory genes (*tax* and *rex*) and the R3 and R4 genes that code for accessory proteins, and a part that encodes microRNAs. The envelope gene (*env*) encodes proteins *gp30* and *gp51*, which play a fundamental role in the infection process, are the recognition center of B lymphocytes and modulate the immune response, so changes in their structure can affect the infective capacity of the virus (Canova et al. 2021; de Brogniez et al. 2016). Therefore, it is a highly conserved genome region and is the most used worldwide for virus detection and genetic research (Blazhko et al. 2020).

In 2016, the molecular prevalence of BLV in animals of the Holstein breed of the department of Antioquia was evaluated, finding 47.6% of positivity (Úsuga-Monroy et al. 2018c). Two years later, in a study carried out in six regions of the country, 73% of positivity was obtained in Antioquia, but only one municipality was evaluated (Corredor-Figueroa et al. 2020). Thus, this is the first extensive study that covers different breeds and geographical regions of specialized dairy in Antioquia. Although various studies have shown the implications of BLV in milk production systems (Bartlett et al. 2014; Úsuga-Monroy et al. 2018a), the virus is not included among the mandatory control diseases in the country by the Colombian Agricultural Institute (Instituto Colombiano Agropecuario - ICA), although, since 2015 and according to Resolution No. 003714 of 2015 (Instituto Colombiano Agropecuario 2015), it is mandatory to declare if its presence is found.

The lack of control mechanisms and knowledge regarding this pathogen by producers have contributed to the increase in the number of positive animals, facilitating the circulation of BLV within dairy herds. Milk production is a sensitive sector of the national primary economy, and the department of Antioquia is one of the leading milk producers in the country, with three regions (north, east, and Valle de Aburrá) where this activity prevails for the peasant economy. With this context in mind, the aim of this study is to evaluate the prevalence of BLV in the main dairy breeds and their crosses in the specialized dairy herds of this department.

MATERIALS AND METHODS

Experimental design

The study population consisted of 224,714 mechanical and traditional specialized dairy milking cows from Valle de Aburrá and the north, and east regions of Antioquia, Colombia, the largest milk-producing regions in the department (Secretaría de Agricultura y Desarrollo Rural, 2019). The sample size was calculated from the study population, with an expected molecular prevalence of 47% (Úsuga-Monroy et al. 2018c), a relative error of 10% (i.e., an absolute error of 4.4%), and a design effect of 1.2, obtaining a sample size of 575 bovines that were selected in 16 municipalities of the three regions mentioned above. The number of bovines sampled per herd was proportional to the sample size and the total

number of bovines in the herd; these were selected randomly. Each herd was given a list of bovines that were selected.

Ethical aspects: Ethics Committee

This research is part of the macroproject "Bovine leukemia virus in dairies of Antioquia: Evaluation of the zoonotic potential and the effect on reproductive performance", endorsed by the Institutional Committee for the Care and Use of Animals (CICUA) of Universidad Nacional de Colombia "020-2020". For the development of the project, no permits or licenses were required. However, in the execution of the project, the legislation and other current regulations pertaining to it in terms of ethics, environmental regulations, or access to genetic resources were complied with. The permits for taking blood samples from the animals and obtaining the required information from each herd were processed with the owners of the herds included in the research, who signed a consent.

Blood sampling

Blood samples were taken from the middle coccygeal vein with an 18G needle with a vacutainer vacuum system (VACUETTE®) and tubes with EDTA as an anticoagulant to obtain the DNA of the nucleated peripheral blood cells. All samples were homogenized by inversion and transferred to the Animal Biotechnology Laboratory of Universidad Nacional de Colombia, Medellín, under refrigeration conditions (4 °C).

DNA extraction

Blood samples with anticoagulant were centrifuged to obtain the layer of white cells from which the DNA was extracted using the salting out technique (Miller et al. 1988). The DNA obtained was resuspended in TE 1X pH=8.0 buffer (1 M Tris HCl and 0.5 M EDTA) and stored

at 4 °C until analysis. The sample was run on a 1% agarose gel stained with SYBR-safe to verify the integrity of the extracted DNA. Extracted DNA was quantified in a spectrophotometer (NanoDrop2000®, Massachusetts, United States) at 260 and 280 nm optical densities to verify its concentration as well as the quality through the 260:280 ratio. In addition, a PCR for a bovine constitutive gene (Toll-like receptor 6) was performed with random samples to verify that there were no inhibitors in the samples.

Molecular analysis for the detection of BLV in cattle Controls

As a positive control for the PCR tests, a mixture of DNA from bovines previously diagnosed positive serologically by the ELISA technique and by PCR for the *env* gene by the BIOGEM group was used. Negative controls were performed with a reaction in the absence of DNA and with DNA from a bovine previously diagnosed by serology and molecularly as negative for BLV infection.

Nested PCR of the *env* gene

The PCR to detect the BLV *env* gene was performed in a final volume of 25 µL with 150 ng of DNA, 0.2 mM of dNTPs, 1 IU of Taq polymerase, 1X of buffer, and 1.5 µL of 10 mM of each envFw and envRv oligonucleotide (Fechner et al. 1996) for the first reaction, and envFW2 and envRV2 (Fechner et al. 1996) for the second reaction (Table 1). The reaction conditions for the two PCR cycles were 5 min at 94 °C, 40 cycles of 30 s at 94 °C, 30 s at 60 °C, 1 min at 72 °C, and 5 min at 72 °C. At the end of the second reaction, a 444 bp fragment of the BLV provirus *env* gene was obtained in the positive animals. Agarose gel electrophoresis was performed to observe the PCR results in which the positive animals for the BLV exhibited a band at the height of 444 pb.

Table 1. Primers used for diagnostic PCR to BLV.

Name	Nucleotide sequences	Expected band size (pb)
envFw	TCTGTGCCAAGTCTCCCAGATA	598
envRv	AACAACAACCTCTGGGGAGGGT	
envFw2	CCCACAAGGGCGGGCGCCGGTTT	444
envRv2	GCGAGGCCGGGTCCAGAGCTGG	

Statistical analysis

The results of the PCR were integrated into an Excel database with the information of the sampled animals (breed, age, parity, region and herd), and descriptive statistics were used to analyze the results. The qualitative traits were summarized in absolute and relative frequencies, estimating 95% confidence intervals. Quantitative variables were summarized using the mean and standard deviation or median with quartiles, depending on data normality. The Chi-square test (X^2) was performed to determine the possible association between breed and the presence or absence of the virus. To establish the association between breed and the presence of BLV, breeds and breed types were grouped into: Holstein x Crossbred: where crossbred can be Viking Red, Ayrshire, BON, Brown Swiss, Normande, Angus; Other purebred: where it can be: Jersey, Ayrshire, BON, Brown Swiss, Viking Red and Other crossbreeds: where there were more than three crossbreds. And they were analyzed through an Odds ratio (OR) by means of a 2x2 contingency table. It was determined that if the OR is more than one, the breed or crossbred has a low risk of being positive for BLV and if the OR is less than one, the breed or crossbred has a high risk of being positive for BLV. The significance of the data was accepted as $P < 0.05$. OR analysis was performed using Epi Info™ v7.2 CDC software.

RESULTS AND DISCUSSION

The fragment amplified in the PCR corresponds to the region that codes for the surface protein *gp51*, vital in the infectivity cycle of the virus since it is responsible for the recognition of the receptor and the binding of the virus to the host cell. Therefore, it is decisive to know the range of virus hosts (Gao et al. 2020). The primers selected to carry out the nested PCR of the *env* gene have been used in various diagnostic and genotyping studies of the virus since 1996 due to the low genetic variability of this region concerning other genome regions (Fechner et al. 1996; Polat et al. 2017).

The results obtained in this study for the 575 bovines belonging to the specialized dairy in Antioquia show that the prevalence of bovine leukemia virus in the department is 17.04% (Table 2), i.e., lower than the results obtained in previous studies (Corredor-Figueroa et al. 2020; Ortiz et al. 2016; Úsuga-Monroy et al. 2018c). This result may be due to the fact that this is the first extensive study

carried out in the region where cattle of different racial components (Holstein, Jersey, Viking Red, Ayrshire, and their Jerhol crosses and trihybrids, among others), age groups (calves, heifers, and cows in production) and geographical regions of the department were sampled. Furthermore, this research contrasts with previous studies where only the population of one breed (Úsuga-Monroy et al. 2018c), a municipality, or an age group were sampled (Corredor-Figueroa et al. 2020; Ortiz et al. 2016).

Table 2. Molecular prevalence to BLV of the *env* gene.

Result	Number of animals	Percentage (%)
Negative	477	82.96
Positive	98	17.04

The low prevalence may also be the result of improved sanitary measures in specialized dairy herds in the department, the implementation of good livestock practices, and biosecurity protocols in the herds that prevent the transmission of the virus, such as the use of one needle per animal, effective as a control strategy for the BLV (Kuczewski et al. 2019). In Colombia, since 2012, the payment of raw milk to producers dependent on compositional and hygienic factors has been implemented, finding that this measure has been effective in stimulating progress in production systems (Carulla and Ortega 2016) and is directly related to the improvement of the sanitary scheme of the production systems and the general management of the animals in it. Nevertheless, the prevalence in herds was relatively high (71.7%) in this study, indicating that in more than half of the productive systems, there is at least one infected animal, existing the risk of an increase in the number of infected animals by contagion. Therefore, it is necessary to evaluate what factors can increase or decrease the risks of spreading the disease in the dairy systems of the department. It is well known that practices such as using shared needles for the application of medicines are one of the main risk factors for the infection of bovines in herds (Kuczewski et al. 2021; Ortiz et al. 2016). Moreover, it is essential to transmit information to producers, creating control strategies to contribute to the progress of this productive sector.

In the department of Antioquia, the main breeds found in specialized dairies are Holstein and Jersey. Crosses

between them and with other reference breeds are also widely used when analyzing prevalence data according to racial components (Table 3). It can be inferred that this factor may be influencing the results obtained since the categories Holstein x (Cross) and other crossbreeds are associated with an increased risk of being infected with BLV. There is evidence that genetic factors can make animals susceptible or resistant to diseases. An example is the Colombian Blanco Orejinegro (BON) breed, where some studies have determined the resistance/susceptibility of these individuals to some viral infections such as foot-and-mouth disease and vesicular stomatitis (Lopez-Herrera et al. 2000, 2002).

In the case of EBL, it has been found that the molecular prevalence of the virus is lower in BON compared to data from pure breeds (Herrera et al. 2011; Úsuga-Monroy et al. 2018b). The heritability of this susceptibility factor to the bovine leukemia virus infection among Holstein and Jersey cattle populations has also been estimated at 8%, indicating that genetics plays an important role in the incidence of BLV in different cattle breeds (Marawan et al. 2021). In

order to analyze the breed factor, several categories were combined to determine whether or not the racial factor has an influence on BLV infection. These results can be a precedent for later studies where this hypothesis was evaluated in depth. It should be noted that the results obtained on the prevalence of BLV in the specialized dairy in Antioquia for the different racial components are not sufficient to make comparative statements between susceptibility/resistance to BLV infection due to the number of individuals for some breeds. However, to establish the association between the breed and the presence of BLV, the following groupings were made: Holstein x Cross, others purebreds and other crossbreeds and compared with the Holstein breed (Table 3). It was identified that the racial component (Holstein x Cross and other crossbreeds) does influence the probability of not having BLV infection ($P < 0.05$) compared to the Holstein breed. Therefore, these results show a trend that generates the hypothesis that there may be breeds with greater resistance to infection by the virus in the Antioquia dairy, however, our data do not allow us to establish individually which breed or breeds are more resistant to BLV infection.

Table 3. Odds Ratio (OR) for other breeds and crosses with respect to the Holstein breed.

Category	OR ¹	95% Confidence interval		Pr>Chisq
Holstein x (Cross ²)	2.4528	1.2739	4.7229	0.006*
Others purebreds ³	1.766	0.8145	3.8292	0.146
Other crossbreeds ⁴	2.4528	1.4608	4.1187	0.0005*

¹OR>1, $P < 0.05$: probability of resistance to BLV; OR>1, $P < 0.05$: probability of susceptibility to BLV. ²Viking Red, Ayrshire, BON, Brown Swiss, Normande, Angus. ³Jersey, Ayrshire, BON, Brown Swiss, Viking Red. ⁴More than three crossbreeds.

It has been found that in the bovine major histocompatibility complex (BoLA), with a fundamental role in the presentation of antigens and the immune response of the organism against infection, there is a genetic region (*BoLA-DRB3*) that influences the development of BLV infection (Lo et al. 2020; Notsu et al. 2021). Most cattle with specific mutations in the allele (*DRB3*009:02*) maintain undetectable proviral loads and are positively related to the development of lymphomas (Lo et al. 2020). Identifying these genetic variations associated with resistance to BLV infection helps develop control programs based on the genetic selection of animals with these genes (Andoh et al. 2021). Studies recommend the evaluation of this factor in BLV positive and negative bovines from dairies in Antioquia

to determine if it is the determining factor for the low molecular prevalence in the population studied.

It has been found that the association between *BoLA-DRB3* gene polymorphisms and BLV infection varies between breeds (Hernandez et al. 2018; Lo et al. 2021a), supporting the hypothesis that some breeds are more susceptible than others to infection with BLV. Moreover, environmental factors also relate to the susceptibility/resistance that cattle may have (Andoh et al. 2021). Therefore, more in-depth studies are needed to test this hypothesis as, so far, only animals heterozygous for *BoLA-DRB3* have been shown to have advantages against BLV infection regardless of their breed (Hernandez et al. 2018; Lo et al. 2021b).

Another factor that can be decisive in the prevalence results of BLV in this study is the age of the sampled animals. Other studies have found that the number of infected animals increases as age increases (Murakami et al. 2011). Calves are less likely to be infected, as the main transmission route of the virus is iatrogenic in production systems (Ruiz et al. 2018). It is until after the first calving that prevalence rates increase due to increased exposure to the virus. Nevertheless, the presence of the virus has been found in calves because there is a risk of transmission during pregnancy and birth and later with milk consumption (Marawan et al. 2021; Ruiz et al. 2018). It should also be considered that the onset of the persistent lymphocytosis stage in infected cattle generally occurs before 5 years of age, and after that, between 5 and 8 years of age, it passes to the lymphoma stage (Zyrianova and Kovalchuk 2019). Therefore, many adult cattle individuals may be positive for viral infection and show positive serological tests but are negative for the molecular detection test by PCR since BLV-positive cells may be sequestered in lymphomas and not be circulating in the blood.

When detailing the seropositivity of the study samples by the number of calvings (Figure 1), it is observed that as the number of calvings increases, the prevalence rate also increases, reaching the highest percentage in the group of cows from four to six calvings (20.7%). Further, the positivity rate decreases in the interval of more than seven calvings, possibly due to the sequestration of BLV-positive cells in internal tissues. On the other hand, it is known that most of the cattle in specialized dairies throughout the country are discarded before reaching seven calvings due to the reduction in productive traits. It is then possible that the lower prevalence in this range of calvings is because the animals that continue in the system after seven calvings have good health conditions, and their productivity has been efficient throughout their lives. Knowing the effects of BLV infection on productive and reproductive characteristics, the possibility that these animals were not infected with BLV during their productive life and, therefore, have not been discarded should be considered.




Figure 1. Results of molecular prevalence to BLV for bovines with a different number of calvings in the specialized dairy in the department of Antioquia. Group 0<2 calvings includes animals from three months of age until the first calving.

Detailing the analysis of seropositivity in the three geographical areas where the study was carried out (Figure 2), the area with the highest prevalence was Valle de Aburrá, with 21.05% positivity to BLV. This area has a lower percentage of livestock production systems dedicated to dairy than the north region of Antioquia, in which the lowest molecular prevalence was found (15.65%).

The analysis of the results obtained in the 16 municipalities evaluated in three regions (Figure 3) shows that there is variability in these, finding values from 7.1% (Carmen de Viboral from the eastern region) to 61.5% (Girardota from the Valle de Aburrá region). Nonetheless, it should be noted that these values cannot be compared because the experimental design was based on the population of each region and not on the one at the municipality




Figure 2. Results of molecular prevalence to BLV for the north (15.6%), east (17.9%), and Valle de Aburrá (21.1%) regions in the department of Antioquia, Colombia

level. However, it is a result that must be considered when proposing future studies. In Valle de Aburrá, the municipality with the highest prevalence was Girardota (Figure 3A). Of the three municipalities evaluated in the region, Medellín has the lowest prevalence despite being the one with the largest bovine population included in the study for this region. The second is the east region

(Figure 3B), with the municipality of Santuario exhibiting the highest prevalence (50%) with respect to the other four evaluated. Conversely, the municipality with the lowest BLV seropositivity, Carmen de Viboral (7.1%), is also found in this municipality. The north region (Figure 3C) was the one where a higher number of municipalities were evaluated (eight) since it is the one with the most specialized dairy,




Figure 3. Molecular prevalence of bovine leukemia virus in municipalities of the department of Antioquia A. Molecular prevalence in the municipalities of Valle de Aburrá, B. Molecular prevalence in the municipalities of the east region, and C. Molecular prevalence in the municipalities of the north region.

having most a low prevalence rate (between 7.7 and 33.3%) compared to the municipalities of other regions. On average, seropositivity for this region is 17%, which is the same value that the molecular prevalence of BLV had for the entire department. These results show a variation between the different geographical areas of the department, and future studies would be necessary to evaluate this and the possible reasons for these differences.

CONCLUSIONS

The bovine leukemia virus is a pathogen found in most of the specialized dairy systems of the department of Antioquia, generating economic losses to producers. Thus, it is necessary to implement measures to control and prevent this disease in the territory. The results of this study contribute to the creation of BLV control strategies in the specialized dairy of Antioquia since differences in molecular positivity to BLV were detected by the number of calvings (age of the cows), racial component, herd, region, and municipality. It is necessary to identify factors contributing to reducing the presence of the virus in the territory to optimize this agricultural sector of the economy in the department. Thus, future studies are necessary to identify in greater detail the behavior of this pathogen under the different conditions that exist in the dairy production systems of Antioquia, as well as the influence that the different racial components may have.

ACKNOWLEDGMENTS

To the G8+1 group of universities and the Government of Antioquia, Colombia, who, in the 2020 joint call for R+D+i projects, financed the “Bovine leukemia virus in dairies of Antioquia: Evaluation of the zoonotic potential and the effect on reproductive performance” project. Furthermore, thanks to the specialized dairy producers, who allowed the collection of samples and the use of data from their herds to carry out this research. Finally, to the Dairy Control Program of Colanta.

REFERENCES

- Andoh K, Akagami M, Nishimori A et al (2021) Novel single nucleotide polymorphisms in the bovine leukemia virus genome are associated with proviral load and affect the expression profile of viral non-coding transcripts. *Veterinary Microbiology* 261. <https://doi.org/10.1016/J.VETMIC.2021.109200>
- Barez PY, de Brogniez A, Carpentier A et al (2015) Recent Advances in BLV Research. *Viruses* 7(11):6080–6088. <https://doi.org/10.3390/V7112929>
- Bartlett PC, Sordillo LM, Byrem TM et al (2014) Options for the control of bovine leukemia virus in dairy cattle. *Journal of the American Veterinary Medical Association* 244(8):914–922. <https://doi.org/10.2460/javma.244.8.914>
- Bartlett PC, Ruggiero VJ, Hutchinson HC et al (2020) Current developments in the epidemiology and control of Enzootic Bovine Leukemia as caused by Bovine Leukemia Virus. *Pathogens* (Basel, Switzerland) 9(12). <https://doi.org/10.3390/pathogens9121058>
- Blazhko N, Vyshegurov S, Donchenko A et al (2020) Genotypes diversity of env gene of Bovine leukemia virus in Western Siberia. *BMC Genetics* 21(Suppl 1): 70. <https://doi.org/10.1186/s12863-020-00874-y>
- Canova R, Weber MN, Budaszewski RF et al (2021) Bovine leukemia viral DNA found on human breast tissue is genetically related to the cattle virus. *One Health* 13:100252. <https://doi.org/10.1016/J.ONEHLT.2021.100252>
- Carulla JE and Ortega E (2016) Dairy production systems of Colombia: challenges and opportunities. *Latin American Archives of Animal Production* 24(2). Retrieved from https://ojs.alpa.uy/index.php/ojs_files/article/view/2526
- Corredor-Figueroa AP, Salas S, Olaya-Galán NN et al (2020) Prevalence and molecular epidemiology of bovine leukemia virus in Colombian cattle. *Infection, Genetics and Evolution*:80. <https://doi.org/10.1016/j.meegid.2020.104171>
- de Brogniez A, Mast J and Willems L (2016) Determinants of the bovine leukemia virus envelope glycoproteins involved in infectivity, replication and pathogenesis. In *Viruses* (Vol. 8, Issue 4). Multidisciplinary Digital Publishing Institute (MDPI). <https://doi.org/10.3390/v8040088>
- Fechner H, Kurg A, Geue L et al (1996) Evaluation of polymerase chain reaction (PCR) application in diagnosis of bovine leukaemia virus (BLV) infection in naturally infected cattle. *Zentralblatt Fur Veterinarmedizin. Reihe B. Journal of Veterinary Medicine. Series B* 43(10):621–630. <https://doi.org/10.1111/j.1439-0450.1996.tb00361.x>
- Gao A, Kouznetsova VL and Tsigelny IF (2020) Bovine leukemia virus relation to human breast cancer: Meta-analysis. *Microbial Pathogenesis* 149. <https://doi.org/10.1016/J.MICPATH.2020.104417>
- Hernandez D, Montes D and Alvarez LA (2018) Association of BoLA-DRB3.2 alleles with enzootic bovine leukosis: profiles BLV infection, persistent lymphocytosis and antibody production in Harton del Valle Cattle. *Indian Journal of Science and Technology* 11(24): 1–14. <https://doi.org/10.17485/ijst/2018/v11i24/128164>
- Herrera Hernandez D, Terranova Posso A and Hernandez Herrera D (2011) Detección del virus de la leucosis bovina en ganado criollo colombiano mediante PCR-anidado. *Acta Agronómica* 60(4): 312–318. ISSN 0120-2812.
- Instituto Colombiano Agropecuario (2015) Resolución No. 003714. Disponible en: <https://www.ica.gov.co/getattachment/3188abb6-2297-44e2-89e6-3a5dbd4db210/2015R3714.aspx>
- Kuczewski A, Hogeveen H, Orsel K et al (2019) Economic evaluation of 4 bovine leukemia virus control strategies for Alberta dairy farms. *Journal of Dairy Science* 102(3):2578–2592. <https://doi.org/10.3168/jds.2018-15341>
- Kuczewski A, Orsel K, Barkema HW et al (2021) Invited review: Bovine leukemia virus—Transmission, control, and eradication. *Journal of Dairy Science* 104(6):6358–6375. <https://doi.org/10.3168/jds.2020-18925>

- Lo CW, Borjigin L, Saito S et al (2020) BoLA-DRB3 Polymorphism is Associated with differential susceptibility to bovine leukemia virus-induced lymphoma and proviral load. *Viruses* 12(3):352. <https://doi.org/10.3390/V12030352>
- Lo CW, Takeshima SN, Okada K et al (2021a) Association of bovine leukemia virus-induced lymphoma with BoLA-DRB3 polymorphisms at DNA, amino acid, and binding pocket property levels. *Pathogens* 10(4):437. <https://doi.org/10.3390/PATHOGENS10040437>
- Lo CW, Takeshima SN, Wada S et al (2021b) Bovine major histocompatibility complex (BoLA) heterozygote advantage against the outcome of bovine leukemia virus infection. *HLA* 98(2):132–139. <https://doi.org/10.1111/tan.14285>
- Lopez-Herrera A, Arango A, Zuluaga F et al (2000) Fenotipificación de la resistencia a la infección por el virus de la fiebre aftosa en ganado blanco orejinegro (BON). *Iatreia* 13.
- Lopez-Herrera A, Salazar A, Restrepo G et al (2002) Resistencia natural, in vitro, a los virus de estomatitis vesicular y de rinotraqueitis infecciosa en ganado Blanco Orejinegro. *Revista Colombiana de Ciencias Pecuarias* 15(1):100-106. ISSN: 0120-0690.
- Marawan MA, Alouffi A, El Tokhy S et al (2021) Bovine leukaemia virus: Current epidemiological circumstance and future prospective. *Viruses* 13(11). Multidisciplinary Digital Publishing Institute (MDPI). <https://doi.org/10.3390/v13112167>
- Miller SA, Dykes DD and Polesky HF (1988) A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Research* 16(3):1215. <https://doi.org/10.1093/nar/16.3.1215>
- Murakami K, Kobayashi S, Konishi M et al (2011) The recent prevalence of bovine leukemia virus (BLV) infection among Japanese cattle. *Veterinary Microbiology* 148(1):84–88. <https://doi.org/10.1016/J.VETMIC.2010.08.001>
- Notsu K, El Daous H, Mitoma S et al (2021) A pooled testing system to rapidly identify cattle carrying the elite controller BoLA-DRB3*009:02 haplotype against bovine leukemia virus infection. *HLA* 99(1):12–24. <https://doi.org/10.1111/TAN.14502>
- Ortiz, D, Sanchez A, Tobon J et al (2016) Seroprevalence and risk factors associated with bovine leukemia virus in Colombia. *Journal of Veterinary Medicine and Animal Health* 8(5):35–43. <https://doi.org/10.5897/JVMAH2016.0457>
- Polat M, Takeshima SN and Aida Y (2017) Epidemiology and genetic diversity of bovine leukemia virus. *Virology Journal* 14(1). <https://doi.org/10.1186/s12985-017-0876-4>
- Ruiz V, Porta NG, Lomónaco M et al (2018) Bovine Leukemia virus infection in neonatal calves. risk factors and control measures. *Frontiers in Veterinary Science* 5(Oct). <https://doi.org/10.3389/fvets.2018.00267>
- Secretaría de Agricultura y Desarrollo Rural (2019) Anuario Estadístico del Sector Agropecuario en el Departamento de Antioquia 2018. In Gobernación de Antioquia.
- Úsuga-Monroy C, Echeverri-Zuluaga JJ and López-Herrera A (2018a) Bovine leukemia virus decreases milk production and quality in Holstein cattle. *Archivos de Zootecnia* 67(258):254–259. <https://doi.org/10.21071/az.v67i258.3661>
- Úsuga-Monroy C, Echeverri-Zuluaga JJ and López-Herrera A (2018b) El componente racial influencia la resistencia a la infección con el virus de la leucosis bovina. *Revista de La Facultad de Medicina Veterinaria y de Zootecnia* 65(2). <https://doi.org/10.15446/rfmvz.v65n2.75632>
- Úsuga-Monroy C, Echeverri-Zuluaga JJ and López-Herrera A (2018c) Molecular and serological detection of bovine leukemia virus in a population of Holstein cows, from Colombia. *Revista Mexicana De Ciencias Pecuarias* 9(2):387–399. <https://doi.org/10.22319/rmcp.v9i2.4232>
- Zyrianova IM and Kovalchuk SN (2019) Bovine leukemia virus tax gene/Tax protein polymorphism and its relation to Enzootic Bovine Leukosis. *Virulence* 11(1):80–87. <https://doi.org/10.1080/21505594.2019.1708051>

