

Seroprevalence and Risk Factors of Brucellosis in Dogs in Enugu and Anambra States, Nigeria*

Chidiebere O. Anyaoha¹ / Lynda O. Majesty-Alukagberie² / Iniobong C I. Ugochukwu³ / John A. Nwanta⁴ / Boniface M. Anene⁵ / Steve I. Oboegbulam⁶

Abstract

Seroprevalence of the antibodies of *Brucella canis* and *Brucella abortus* in dogs was assessed using a cross-sectional survey in Anambra and Enugu States, Nigeria. A total of 123 Companion dogs made up of 65 clinic dogs, 34 slaughter dogs and 24 household dogs were screened. For *B. abortus* antibody assay, the collected serum was used for Rose Bengal plate test (RBPT), Serum agglutination test (SAT) and Solid Phase Immunoassay technique with Immunocomb® Canine Brucellosis Antibody Test Kit was used. Out of the 123 dogs screened, none was positive for *Brucella abortus* antibodies while 34 (27.7%) of the dogs screened were positive for *B. canis* antibodies. There was a significant association ($P < 0.05$) between infection and sex, the infection was significantly higher ($P < 0.05$) in female than male dogs. Prevalence was significantly higher ($P < 0.05$) in Exotic breeds than in mixed and local dog breeds. There was no association ($P > 0.05$) between infection and antibody titre levels in the different categories of dogs. However, there was significant association ($P < 0.05$) between the presence of *Brucella canis* antibodies and free roaming of dogs. This study provides the first serological evidence of *B. canis* infection in dogs in Enugu and Anambra States. This shows that *B. canis* is endemic in both states, underscoring the need for further studies. Female dogs, exotic breeds and freely roaming dogs are at a higher risk of *Brucella* infection in the study area; therefore, preventive and control measures are strongly recommended.

Keywords: antibody titre, *Brucella abortus*, *Brucella canis*, Canine brucellosis, risk factors, seroprevalence, zoonosis

* Investigation article.

1 Department of Veterinary Public Health and Preventive Medicine, University of Nigeria, Nsukka.

<https://orcid.org/0000-0001-7883-8063>

2 Department of Veterinary Public Health and Preventive Medicine, University of Nigeria, Nsukka.

<https://orcid.org/0000-0001-7294-3487>

3 Department of Veterinary Pathology and Microbiology, University of Nigeria, Nsukka

✉ iniobongugochukwu@unn.edu.ng

<https://orcid.org/0000-0001-7410-8311>

4 Department of Veterinary Public Health and Preventive Medicine, University of Nigeria, Nsukka.

<https://orcid.org/0000-0002-9923-8005>

5 Department of Veterinary Medicine, University of Nigeria, Nsukka.

6 Department of Veterinary Public Health and Preventive Medicine, University of Nigeria, Nsukka.

<https://orcid.org/0000-0003-4229-2307>

How to cite this article: Anyaoha CO, Majesty-Alukagberie LO, Ugochukwu ICI, Nwanta JA, Anene BM, Oboegbulam SI. Seroprevalence and Risk Factors of Brucellosis in Dogs in Enugu and Anambra States, Nigeria. Rev Med Vet. 2020;(40): 45-59. <https://doi.org/10.19052/mv.vol1.iss40.5>

Seroprevalencia y factores de riesgo de la brucelosis en perros de los Estados Enugu y Anambra, Nigeria

Resumen

Se evaluó la seroprevalencia de los anticuerpos de *Brucella canis* y *Brucella abortus* en perros usando un sondeo transversal en los Estados Anambra and Enugu, Nigeria. Se examinó un total de 123 perros de compañía, de los cuales 65 eran perros de clínica, 34 perros de matadero y 24 perros caseros. Para el ensayo de anticuerpos de *B. abortus*, el suero muestreado se usó para la prueba de Rosa de Bengala (RBPT), prueba de aglutinación del suero (SAT) y se usó la técnica de inmunoensayo en fase sólida con el kit de prueba de anticuerpos para brucelosis canina Immunocomb®. De los 123 perros analizados, ninguno dio positivo para los anticuerpos de *Brucella abortus* mientras que 34 (27.7%) de los perros analizados dieron positivo para los anticuerpos de *B. canis*. Hubo una asociación significativa ($P < 0.05$) entre infección y género; la infección fue significativamente más alta ($P < 0.05$) en las hembras que en los machos. La prevalencia fue significativamente

más alta ($P < 0.05$) en las razas exóticas que en las razas cruzadas y las razas locales. No hubo ninguna relación ($P > 0.05$) entre la infección y los niveles de titulación de anticuerpos en las diferentes categorías de perros. Sin embargo, hubo una relación significativa ($P < 0.05$) entre la presencia de anticuerpos *Brucella canis* y los perros que andan libremente por doquier. Este estudio provee la primera evidencia serológica de infección con *B. canis* en perros de los Estados Enugu y Anambra. Esto muestra que la *B. canis* es endémica en ambos estados, enfatizando la necesidad de hacer más estudios. Las hembras, las razas exóticas y los animales que deambulan libremente se encuentran en el riesgo más alto de infección con *Brucella* en el área de estudio; por consiguiente, se recomienda enormemente tomar medidas preventivas y de control.

Palabras clave: titulación de anticuerpo, *Brucella abortus*, *Brucella canis*, brucelosis canina, factores de riesgo, seroprevalencia, zoonosis

INTRODUCTION

Nigeria with its estimated 180,000,000 population, is the most populous country in the African continent. It has a great proportion of the world's poor livestock keepers and provides a focal point for neglected zoonosis (1). Brucellosis is a devastating zoonotic disease causing huge economic losses for livestock farms around the globe (2, 3). It also poses serious human health hazards worldwide (4, 5, 6).

The genus *Brucella* comprises various species with both veterinary and human medical importance and contains a group of closely related species (7). This group includes *Brucella melitensis* affecting primarily small ruminants while *B. abortus* affects primarily cattle, *B. suis* affects pigs, *B. ovis* affects sheep, *B. canis* affects dogs while the other members include, *B. neotomae*, *B. microti*, *B. ceti* and *B. pinnideipialis* (8, 9, 10, 11).

Brucella spp. infect not only their specific hosts but also other domestic and wild animal species, which in turn can serve as disease reservoirs for other animal species and humans (2). Brucellosis, a major neglected zoonotic disease, is transmitted by direct contact with infected animals, their secretions, or by ingesting their dairy products, which requires intensive attention in many communities around the globe (2, 12). It could also be transmitted by consumption of contaminated food of animal origin, and through aerosol (13). Infection can also result through contact with infected aborted materials such as aborted fetuses, placenta membranes or fluids and other vaginal exudates (2,11).

A report of brucellosis outbreak was firstly reported in a government cattle farm located in Zaria in 1934. Serum plate agglutination test showed positive reaction in about 15% of the total animals in the farm (14).

Brucellosis is a reportable disease in Nigeria, but its incidence, prevalence and distribution is difficult to determine as the system of disease surveillance and reporting is incomplete and inept (15,16). Serological prevalence rates between 0.20% and 79.70% have been reported

in animals and humans in various parts of the country (17-22) emphasis is laid mostly on cattle, sheep and goat; but the same cannot be said of dogs.

Canine brucellosis can lead to reproductive losses both in dog and human infections due to contact with infected urine or other genital secretions (23). *Brucella canis*, the aetiology of Canine brucellosis, is considered a rare cause of human brucellosis but the clinical importance of this infection may have been underrated so far because of frustrations and complications associated with making a diagnosis (24).

The rise in pet ownership in Nigeria, especially dogs, is linked with some risk factors that render the Nigerian human population vulnerable to this disease. It is worth mentioning that many imported foreign dog breeds are not screened before entering into our country (25), thus providing a source of infection. In turn, dogs from Nigeria can be found elsewhere around the world.

As a zoonosis, it is also regarded as a significant public health issue, thereby making susceptible some persons in occupational risk like butchers, abattoir workers, veterinarians and livestock owners as well as other humans (26, 27, 6).

This study will provide information on the current seroprevalence of *Brucella* antibodies in dogs in Enugu and Anambra States, Nigeria as well as the prevalent species. It will also provide information on sex, breed, and age distribution as well as some risk factors that may influence *Brucella* infection in dogs, thus contributing to the epidemiology of Canine Brucellosis in Enugu State and Anambra State, Nigeria.

MATERIALS AND METHODS

Study design

The study was conducted as a cross-sectional survey using purposive sampling technique to screen dogs for *Brucella* antibodies.

Study Area

Enugu State is located between latitude 5°55"N and 7°55"N and longitude 6°53"E and 7°55"E while Anambra State is located between latitude 6°20"N and longitude 7°00"E. Enugu state covers a total land area about 802,295 Km² and has a population of 2,500,000 with a population density of 240 persons per square kilometers (28). On the other hand, Anambra State covers a total land area about 4,844 Km² and according to the National Population Commission (2006), it has a population about 3,902,051 with a population density of about 840 persons per square kilometre (28). Dogs are kept as part of the people's culture in these 2 States for breeding, hunting and protecting their homes. Dog meat is eaten by some parts of the population.

Study Population

The study purposively targeted 3 groups: (a) Dogs taken to veterinary clinics around the states; (b) Household dogs with history of infertility or abortion; and (c) Dogs at slaughter points in the markets as shown in Figure 1.

Figure 1. Dogs at slaughter points.



Sample Size Determination

The required sample size was determined using the following formula:

$$n = \frac{Z^2 pq}{d^2} \quad (29)$$

Where n = Desired sample size, $Z^2 = 1.96$ (normal distribution) from table, p = Prevalence rate from the average of previous studies, d = Desired absolute precision of $\pm 5\%$ with 95% CI, $q = 1 - p$. In this study, according to Adedoyin (30), a prevalence rate of 7.94% was used for sample size determination. Using the formula above, a sample size of 112 was calculated. However, a sample size of 123 dogs was screened in this study.

Sampling Technique

Five major veterinary clinics, 2 in each Enugu and Onitsha metropolis, one in Nsukka Urban; 2 major slaughter points in each state; and households with dogs having history of infertility or abortion were selected by purposive sampling method. Visits were made to the purposively selected veterinary clinics, households, and slaughter points, once every other week for 6 months. A total of 123 dogs made up of 65 Clinic dogs, 34 Slaughter dogs and 24 Household dogs were screened. Profiles of the dogs taken to the clinics and household dogs were also collected. Gathered data include: sex, age, and breed, history of infertility / abortion in female dogs, and some possible risk factors / management practices to *Brucella* infection in dogs.

Collection of Blood Samples

Five milliliters of blood were aseptically collected from the cephalic vein of each animal, after proper restraint. The blood was kept in a slanted position for about 30 minutes to allow for proper clotting. It was then centrifuged at 3,000 rpm for 5 minutes. The sera was then harvested one after the other using separate syringes into well labelled bijoux bottles and stored at -20°C until they were analysed.

Serological Sample Analysis

For *B. abortus* identification, 2 tests were used, namely, Rose Bengal Plate Test (RBPT) and Serum Agglutination Test (SAT). The Antigen was procured from Central Veterinary Lab., New Haw, Weybridge Surrey, England. Both of the tests were done as described by Alton *et al.*, (31) and Morgan *et al.* (32). Titres of 1:40 (50 IU/ml) and above were taken as diagnostic for *B. abortus* as determined by Morgan *et al.*, (32) and Sati, (33).

Identification of *B. canis* was done using Immuno-Comb® Canine Brucellosis Antibody Test Kit specific for *B. canis* antigen (Biogal Galed Labo., Kibbatz Galed 19240, Israel) and this was done as previously described by Muhairwa *et al.* (34). Furthermore, Blood bacterial culture was used for confirmation of the seropositive sampled dogs. The bacteria were cultured aerobically and the identity confirmed by standard gram staining, microscopic and biochemical tests characteristics as described by Chessbrough, (35).

Ethical Considerations

Ethical considerations guiding the use and behavior in experiments on animals were strictly observed and the experimental protocol was approved by the University of Nigeria Nsukka Senate Committee on Medical Re-

search Ethics. Proper permit and consent were obtained from the Veterinary Clinics before obtaining the data for this experiment.

Statistical Analysis

Using Statistical Package for Social Sciences (SPSS) 17.0, Chi-square (χ^2) statistic, odds ratio and 95% confidence interval were used to determine whether there were significant association between *Brucella* antibody prevalence and sex, age, breed, history of infertility/abortion in female dogs, and some possible risk factors/management practices to *Brucella* infection in dogs.

RESULTS

Distribution of Dogs based on States and Sources

Over the study term, a total of 123 dogs were screened. Table 1 summarizes the sources/categories of the dogs screened. Enugu State accounted for 68 of the total dogs while Anambra accounted for 55 of the total dogs screened. Veterinary clinics accounted for 65 of the total dogs screened while Slaughter points and households for 34 and 24 dogs screened, respectively.

Table 1. Distribution of dogs screened for Brucellosis based on sources

Source	Number of screened dogs		
	Anambra State	Enugu State	Total
Veterinary Clinics	37	28	65
Slaughter House/ Market	6	28	34
Household	12	12	24
Grand Total	55	68	123

Source: own work

Prevalence of *Brucella abortus* Antibodies Based on the Sources of the Dogs

Out of the 123 sera samples screened for *Brucella abortus* antibodies, none (0%) was positive using both the Rose Bengal Plate Test (RBPT) and the Serum Agglutination Test (SAT) (Table 2).

Prevalence of *Brucella canis* Antibodies Based on the Sources of the Dogs

Out of the total 123 dogs from the different sources screened, 34 were positive for *B. canis* antibody using the Immunocomb[®] Canine *Brucella* Antibody Test Kit as shown in Figure 2. This gave an overall seroprevalence rate of 27.7%. Out of 65 dogs screened in Veterinary Clinics (clinic dogs) in Anambra and Enugu States, 22 or 18% were positive for *B. canis* (Table 3). Four (3.3%) out of 34. slaughter dogs screened were positive. Out of the 24 household dogs screened, 8 or 6.5% were posi-

tive. There was no association between infection rate and sources/categories of the dogs screened ($\chi^2 = 5.925$, $P > 0.05$).

Figure 2. Immuno-Comb showing positive serum samples * represents the positive serum samples.



Source: own work

Table 2. Prevalence of *Brucella abortus* Antibodies Based on the Sources of the Dogs

Source	Number screened	Number positive	
		RBPT	SAT
Vet. Clinics (Clinical cases)	65	0 (0.00)	0 (0.00)
Slaughter points	34	0 (0.00)	0 (0.00)
Households	24	0 (0.00)	0 (0.00)
Total	123	0 (0.00)	0 (0.00)

Legend: RBPT=Rose Bengal Plate Test SAT= Serum Agglutination Test.

Source: own work

Table 3. Prevalence of *Brucella canis* Antibodies Based on the Sources of the Dogs

Source	Number screened	Number positive Immunocomb [®] B. canis Antibody Test Kit
Vet. Clinics (Clinical dogs)	65	22 (18)
Slaughter points	34	4 (3.3)
Households	34	8 (6.5)
Total	123	34 (27.7)

($\chi^2 = 5.925$, $P > 0.05$)

Source: own work

Antibody Titre Levels of *B. canis* Positive Dogs Screened in Anambra and Enugu States

Out of the 34 *B. canis* positive dogs as shown in Figure 1, 16 (47.1%) had a titre level of 1:200 (IFA Titre); 10 (29.4%) had titre level of 1:400 (IFA Titre), 4 (11.8%) had a titre level of 1:600 (IFA Titre); while the remaining 4 (11.8%) had a titre level of 1:800 (IFA Titre) (Table 4).

Antibody Titre Levels of *Brucella canis* Positive Dogs Based on the Sources of Sogs

The antibody titre levels of *B. canis* positive dogs based on sources/categories of screened dogs are shown in Table 4. Ten (29.4%) of 16 positive dogs with antibody titre 1:200 (IFA Titre) are clinic dogs; 4 (11.8%) are

slaughter dogs and 2 (5.9%) are household dogs. Seven (20.6%) of 10 positive dogs with titre 1:400 (IFA Titre) are clinical dogs; 3 (8.8%) are slaughter dogs and none from the households. Five (14.7%) of 8 dogs with titre 1:600 (IFA Titre) and above are clinic dogs; 2 (5.8%) were household dogs and only one (2.9%) was a slaughter dog. Chi- Square analysis showed no association between antibody titre levels and sources of the dogs ($\chi^2 = 3.767$; $p > 0.05$). Two household dogs and 2 of the 5 clinic dogs with titre 1:600 (IFA Titre) and above had history of recent abortion or infertility.

Sex Distribution of *Brucella canis* Antibody Positive Dogs

Female dogs had a seroprevalence of 22% (Table 6) while male dogs had a seroprevalence of 6%. There was a strong association ($p < 0.05$) between the infection of *Brucella canis* and sex of the dogs screened (Table 5).

Table 4. Antibody Titre Levels of *Brucella canis* Positive Dogs Based on the Sources of Dogs

Titre level	Total	Clinical dogs (%)	Slaughter Dogs (%)	Household Dogs (%)
1 : 200	16	10 (29.4)	4 (11.8)	2 (5.9)
1 : 400	10	7 (20.6)	3 (8.8)	0 (0)
1 : 600	4	3 (8.8)	0 (0)	1 (2.9)
1 : 800	4	2 (5.9)	1 (2.9)	1 (2.9)
Total	34	22 (64.7)	8 (23.5)	4 (11.8)

($\chi^2 = 3.767$; $p > 0.05$)

Source: own work

Table 5. Sex Distribution of *Brucella canis* Antibodies of Dogs Sampled in Southeast Nigeria

Sex	Number sampled	Number positive (%)
Male	45 (36.6)	7 (5.9) ^a
Female	78 (63.4)	27 (21.9) ^b
Total	123 (100.0)	34 (27.6)

($\chi^2 = 5.174$, $p < 0.023$, $df = 1$)

Source: own work

Antibody Titre Levels of *B. canis* Positive Dogs according to Sex

Out of the 27 positive female dogs, 13 (38.2%), 8 (23.5%), 3 (8.8%) and another 3 (8.8%) had antibody titre levels of 1:200 (IFA Titre), 1:400 (IFA Titre), 1:600 (IFA Titre) and 1:800 (IFA Titre), respectively. In males out of the 7 positive dogs, 3 (8.8%), 2 (5.9%), 1 (2.9%) and 1 (2.9%) had titre levels of 1:200 (IFA Titre), 1:400 (IFA Titre), 1:600 (IFA Titre) and 1:800 (IFA Titre), respectively. Chi-square analysis showed that there is no association ($P>0.05$) between titre level and sex (Table 6).

Age Distribution of *Brucella canis* Antibody Positive Dogs (Clinic and Household Dogs)

Dogs below 1 year old had a seroprevalence of 3.4%, dogs 1-<3 years of age had seroprevalence of 10.1%, while those 3-<5 years and 5 years and above, had seroprevalence of 15.7% and 4.5%, respectively. Thus, 23 out of 66 dogs aged between 1 year and 5 years were positive, having a seroprevalence of 34%. There was no association ($p>0.05$) between *Brucella canis* infection and age in the dogs screened (Table 7).

Table 6. Antibody Titre Levels of *B. canis* Positive Dogs according to Sex

Titre level	Total (%)	Female (%)	Male (%)
1:200	16	13 (38.2)	3 (8.8)
1:400	10	8 (23.5)	2 (5.9)
1:600	4	3 (8.8)	1 (2.9)
1:800	4	3 (8.8)	1 (2.9)
Total	34 (100)	27 (79.4)	7 (20.6)

($P>0.05$; $\chi^2 = 0.130$)

Source: own work

Table 7. Age Distribution of *Brucella canis* Antibody Positive Dogs (Clinic and Household Dogs)

Age (years)	Number sampled	Number positive (%)
<1year	11	3 (3.4)
1-<3 years	32	9 (10.1)
3-<5 years	34	14 (15.7)
5 & above	12	4 (4.5)
Total	89	30 (33.7)

($\chi^2 = 1.500$, $P>0.05$)

Source: own work

Breed Distribution of *Brucella canis* Antibody Positive Dogs

Twenty-six out of 76 exotic breeds of dog were positive for *B. canis* antibody, which provides a seroprevalence of 21.1% (Table 8). Mixed and local breeds each had seroprevalence of 3.3%. Chi-square analysis revealed that there was a strong association ($P < 0.05$) between infec-

tion and the breeds of dogs, being the infection higher in exotic breeds of dogs.

Table 9 shows the risk factors associated with the presence of *Brucella canis* antibodies in dogs sampled in Enugu and Anambra States of Southeast Nigeria. Female dogs, exotic breeds and freely roaming are at a higher risk of *Brucella* infection in the study area.

Table 8. Breed Distribution of *Brucella canis* Antibody Positive Dogs

Breed	Number sampled	Number Positive (%)
Exotic*	76	26 (21.1)a
Mixed	13	4 (3.3)b
Mongrel	34	4 (3.3)b
Total	123	34 (27.6)

($X^2 = 0.04$; $p < 0.05$)

*Total no of dogs sampled= Rottweiler 28 (36.8%); Mastiff 20 (26.3%); Caucasian 13 (17.1%); Alsatian 13 (17.1%); Boar bull 1 (1.3%); Persian 1 (1.3%).

Source: own work

Table 9. Risk Factors associated with the presence of *Brucella canis* antibodies in dogs presented at the clinics and household dogs sampled in Enugu and Anambra States

Parameter	Total no sample	P. value
SEX Male	45	0.023*
Female	78	
AGE Less than 1 year	11	0.682
1 - < 3years	32	
3 - < 5years	34	
5 and above	12	
Breed Exotic	76	0.050*
Mixed Mongrel	13	
Local	34	
Other animals in the household		
Sheep	1	0.072
Others	6	
None	82	

Continúa

Parameter	Total no sample	P. value
Animal move freely in the neighbourhood		
Yes	35	0.000*
No	54	
Feeding habits of dog		
Scavenging all the time	-	-
Both scavenging and household	27	0.354
Household food only	62	
Purpose of keeping dogs		
Companion for children/security	-	-
Breeding	87	0.308
Hunting	2	
Dog sharing rooms with the household		
Yes	22	0.462
No	67	
History of infertility/abortion in females		
Yes	46	0.898
No	22	
Male dog used for mating		
Yes	16	0.627
No	5	-

Source: own work

DISCUSSION

Brucellosis is endemic in Nigeria and risk factors enhancing its transmission are prevalent (36, 16). *Brucella abortus* antibody was not detected in any of the 123 dogs screened. The zero prevalence suggests that *B. abortus*, though important in livestock ruminants, is not significant in the epidemiology of canine brucellosis in the study area. The apparent absence of *B. abortus* antibody may be attributed to the type of management system practiced in Southeast Nigeria. Dogs are either housed or caged and though they may roam freely in the neighbourhood, they rarely come in contact with livestock ruminants. The dogs medical profiles also showed that most clients feed their dogs with cooked household

foods only. Therefore, the likelihood of being infected with *B. abortus* is rare. The findings of this study on *B. abortus* is in contrast to the works of Cadmus *et al.*, (25) and Adedoyin *et al.*, (30) who reported seroprevalence of 5.46% and 2.48%, respectively, in household dogs in Ibadan, Southwest Nigeria.

The result of the present study suggests a high seroprevalence (27.7%) of *B. canis* in dogs taken to veterinary clinics, apparently healthy slaughter dogs and those in households which were screened. None of the dogs screened in the study were vaccinated, as vaccination of dogs against brucellosis is not routinely carried out in Nigeria because there is no information of any vaccine against *B. canis* (37). Therefore, appearance of anti-

bodies to *Brucella* in dogs in the study area is suggestive of natural exposure to the organism. It also indicates the lack of brucellosis control programme. The seroprevalence of *B. canis* in this study was relatively high when compared to other studies in Nigeria. This may be attributed to the higher sensitivity of the diagnostic technique (Biogals Immunocomb[®] Canine *Brucella* Antibody Test Kit) used. Cadmus *et al.*, (25) and Adedoyin *et al.*, (30) reported a seroprevalence of 0.27% in household dogs and 3.11% in household and hunting dogs, respectively, using *B. canis* Rapid Side Agglutination Test (RSAT). However, this result was lower than the prevalence of 59.43% reported by Cadmus *et al.*, (25) in dogs used for hunting and a seroprevalence of (29.2%) which was reported by Momoh *et al.*, (38). The hunting dogs may have higher exposure probability to *Brucella spp* according to the studies conducted by Cadmus *et al.*, (25).

Female dogs had a higher seroprevalence percentage (21.3%) than male dogs (6.0%). A major contributing factor to higher rates in females could be that a single male dog, if infected, is used in mating different females, it can transmit the infection through infected semen (25). Also it may be due to the fact that most dog owners in our study area preferred to keep more female dogs than males for the purpose of additional income through the sale of their puppies. This increases the chances of more females getting infected during mating. However, Radostits *et al.*, (39) have shown that erythritol, a polyhydric acid found in higher concentration in the placenta and foetal fluids of females than in seminal vesicles and testis of males, can be responsible for females being more susceptible than males. This result was in agreement with the findings by Cadmus *et al.* (25) who reported a prevalence of 6.17% in females and 4.9% in males and Momoh *et al.*, (38) who reported a prevalence of 29.3% in female dogs and 28.6% in male dogs. However, it disagrees with findings in a previous study where a slightly higher rate in males (29.6%) than in females (26.7%) was recorded by Adesiyun *et al.*, (40).

The decrease in the positive samples as titre level increases may be attributed to the disposing or selling off of non-producing dogs by humans as the infection may

have entered into a chronic phase, thereby making these infected dogs not to be productive while those with lower titre levels are more in number because the infection may not have started manifesting its clinical signs.

Dogs taken to the clinics had the highest numbers of positive samples in the different titre levels and this may be attributed to the fact that more samples were gotten from clinical cases more than the other categories of dogs sampled.

The different titre levels were also higher in females more than in males and this may be attributed to the fact that female dogs were sampled more in the study. In addition, due to the asymptomatic nature of *B. canis* in dogs, most male dogs used for breeding are unscreened, and carry the infection for a long period of time, shedding it in the environment through urine and semen; this may result in bitches being mated many times by the same infected male dog as a result of repeated unsuccessful breeding attempt, thereby increasing the infection load in females.

Prevalence was lower among the young animals screened as compared to the older ones. Usually young animals are protected by maternal immunity and thus they are less susceptible to infections. This shows that the infection increases with age. The high prevalence seen in older animals shows the chronic nature of brucellosis as it has been shown to increase with age, and most affected animals carry the infection throughout their lives (39). The reason for the increase in prevalence as the animal age increases may be due to the fact that the bacteria localizes mainly in the reproductive tracts, especially in gravid animals. There is also evidence that the mammary gland may be even a more probable area of localization than the reproductive tract (41). Age-wise prevalence studied by Aulakh *et al.* (42), Abubakar *et al.*, (43) and Momoh *et al.* (38), showed that the incidence is higher in sexually mature animals. Therefore, the increase in age, increases probability of exposure to infection in dogs. However, the results in this study do not agree with previous study by Cadmus *et al.* (25), as they reported more prevalence in dogs below one year old than in adult dogs.

There is a strong association between the infection rate and breeds of dogs screened; with infection occurring more in the exotic breeds than the mixed and local (mongrel) breeds. This may be related to the fact that exotic breeds are the dogs of choice among owners in both states and, therefore, they were the prevalent breeds (61.8%) in the population sampled (76/123). This way the probabilities of occurrence is higher in these breeds. Behzadi and Mogheseh (26), argued that the detection of canine brucellosis in exotic dogs may indicates a new source of infection from abroad as these dogs may be imported from countries and regions where the disease is endemic. The higher prevalence among the exotic breeds is in agreement with the findings by Behzadi and Mogheseh (26); they recorded a prevalence of 19.35% in exotic breeds. It is also in agreement with the findings of Cadmus *et al.*, (25) who got 50.55% in Alsatian breeds of dogs.

There is a strong association ($P < 0.05$) between *B. canis* infection and some risk factors such as sex, breed and dogs moving freely in the neighbourhood; This was further supported by the multivariate logistic regression. Notwithstanding, some of the other risk factors such as age, dogs being used for breeding, dogs sharing rooms with humans, female dogs with history of infertility and abortion, and male dogs used for mating, all provided a considerable higher positive numbers and increased the chances of *Brucella* infection in dogs in the study area, even though the numbers are not significant.

Clinical diagnosis of canine brucellosis just based on the clinical signs is not sensitive enough and a negative blood culture cannot rule out the disease (44; 45). Apart from the use of clinical signs and microbiological cultures, serological tests such as Rapid Slide Agglutination Test (RSAT), 2-mercaptoethanol Rapid Slide Agglutination Test (2ME-RSAT), Agar Gel Immunodiffusion (AGID), Indirect Enzyme Linked Immunoassay (IELISA) and Polymerase Chain Reaction (PCR) can be used for routine diagnosis (46, 44).

Based on the findings of this study, we deduced that the zero seroprevalence of *B. abortus* antibodies in dogs in

this study rules out the possibilities of mixed infections but does not imply the absolute absence of the disease in the studied localities. However, it may be inferred that *B. abortus* in dogs does rarely occur in this region, except when the dogs are in close contact with infected farm animals, as proven by previous researchers.

This study has also shown a *Brucella canis* antibodies seroprevalence of 27.6 % in the study area. It means that the general population is at risk and it calls for serious interventions considering the zoonotic implications of the disease as infected dogs can be a source of infection not only to animals but also to humans, especially in close contacts with these animals due to their occupation. Some factors such as sex, breed, age, dogs moving freely in the neighbourhood, dogs being used for breeding and male dogs used for mating all of them increase the chances of *Brucella* infection in dogs in the study area and, therefore, should be considered in the epidemiology of canine brucellosis in the study area.

The Immunocomb® Canine Brucellosis Antibody Test Kit used for this study has proven to be a reliable test method (99% sensitive and specific) in the diagnosis of *Brucella canis* in dogs, which agrees with the findings by Muhairwa *et al.*, (34) and Chinyoka *et al.*, (47).

We therefore make the following recommendations: free roaming of dogs in dog care management practices is a risk factor for transmission of *Brucella canis* infection in the study area; thus, dogs should be restricted within cages in a fenced home. Feeding dogs with fresh carcasses from abattoirs is associated with higher seropositive of *Brucella canis* infection; then this habit should be avoided, unless carcasses are cooked. Exotic breeds of dogs imported into Nigeria should be properly screened for Brucellosis. Control measures should also be enforced in the study area to remove or eradicate the infections among dogs and avoid the spread to other uninfected dogs as well as the possible transmission to humans. We also recommend that appropriate hygienic measures such as proper disposal of aborted fetuses, placenta and other contaminated materials and disinfection of kennels, premises should be strictly observed

by the dog owners, especially those having dogs as pets and allow them into their homes. Finally, Butchers, animal health workers, meat handlers and laboratory workers, including veterinarians, should wear protective garments to avoid direct contact and inhalation of contaminated aerosols in the clinics and kennels. State and federal health agencies should create a veterinary public health unit that will be handling zoonotic diseases as in the developed countries, for the prevention, surveillance and control of zoonosis in general and brucellosis in particular.

ACKNOWLEDGEMENT

The authors are grateful to the staff of Department of Veterinary Public Health and Preventive Medicine Laboratory, University of Nigeria, Nsukka for their expert assistance throughout this study.

CONFLICT OF INTERESTS

The authors of this manuscript declare not to have any financial or personal conflicting interests.

ETHICAL STATEMENT

Ethical considerations governing the use and behavior in experiments with live animals were observed carefully. The experimental protocol was approved by the University of Nigeria, Nsukka Senate Committee on Medical and Research Ethics.

REFERENCES

- Ducrottoy, MJ, Bertu WJ, Ocholi RA, Gusi AM, Brysinckx W, Welburn S, *et al.* Brucellosis as an emerging threat in developing economies: Lessons from Nigeria. *PloS Negl Trop Dis* 2014; 24, 8(7):e3008. <https://doi.org/10.1371/journal.pntd.0003008>
- Díaz Aparicio E. Epidemiology of brucellosis in domestic animals caused by *Brucella melitensis*, *Brucella suis* and *Brucella abortus*. *Rev Sci Tech* 2013; 32(1): 43-51.
- Njeru J, Wareth G, Melzer F, Henning K, Pletz MW, Heller R, *et al.* Systematic review of brucellosis in Kenya: disease frequency in humans and animals and risk factors for human infection. *BMC Pub Hlth* 2016; 16: 853. <https://doi.org/10.1186/s12889-016-3532-9>
- Hamidy MER, Amin AS. Detection of *Brucella spp.* in the milk of infected cattle, sheep, goats and camels by PCR. *Vet J* 2002; 163: 299-305.
- Cadmus SIB, Ijagbone IF, Oputa HE, Adesokan HK, Stack JA 2006. Serological survey of brucellosis in livestock animals and workers in Ibadan, Nigeria. *Afr J Biomed Res* 2006; 9:163-8.
- Zhang N, Zhou H, Huang DS, Guan P. Brucellosis awareness and knowledge in communities worldwide: A systematic review and meta-analysis of 79 observational studies. *PLoS neglect trop dis* 2019; 13(5): e0007366. <https://doi.org/10.1371/journal.pntd.0007366>
- Scholz HC, Mühldorfer K, Shilton C, Benedict S, Whatmore AM, Blom J, *et al.* The Change of a Medically Important Genus: Worldwide Occurrence of Genetically Diverse Novel *Brucella* Species in Exotic Frogs. *PLoS ONE* 2016; 11(12): e0168872. <https://doi.org/10.1371/journal.pone.0168872>
- Scholz HC, Hubalek Z, Sedlacek Vergnaud G, Tomaso H, Aldahouk S, Melzer F, *et al.* *Brucella macroti sp.* Isolated from the common vole *Microtus arvalis*. *Int J Syst Evol Microb* 2008; 58: 375-82.
- Olsen SC, Palmer MV. Advancement of knowledge of *Brucella* over the past 50 years. *Vet Pathol.* 2014;51(6):1076–89.
- Franc, K.A., Krecek, R.C., Häsler, B.N. *et al.* Brucellosis remains a neglected disease in the developing world: A call for interdisciplinary action. *BMC Public Health* 18, 125 (2018). <https://doi.org/10.1186/s12889-017-5016-y>
- Tekle M, Legesse M, Edao BM, Ameni G, Mamo G. Isolation and identification of *Brucella melitensis* using bacteriological and molecular tools from aborted goats in the Afar region of north-eastern Ethiopia. *BMC*

- Microbiol 2019; 19: 108. Available from: <https://doi.org/10.1186/s12866-019-1474-y>
12. De Massis F, Zilli K, Di Donato G, Nuvoloni R, Pelini S, Sacchini L, D'Alterio N, Di Giannatale E. Distribution of *Brucella* field strains isolated from livestock, wildlife populations, and humans in Italy from 2007 to 2015. *PloS one* 2019; 14(3): e0213689. <https://doi.org/10.1371/journal.pone.0213689>.
 13. Gul ST, Khan A. Epidemiology and epizootology of brucellosis: A review. *Pak Vet J* 2007; 27: 145-51.
 14. Banerjee K, Bhalty MA. A survey of bovine brucellosis in Northern Nigeria (A preliminary communication). *Bull Epiz Dis Afr* 1970; 18: 333-335.
 15. Ocholi RA, Kwaga JP, Ajogi I, Bale J, Bertu WJ. Epidemiology, Problems and prospect for control of brucellosis in Nigeria. *Vom J Vet Sci* 2004; 1: 78-86.
 16. Agada CA, Mohammed J, Okoh AEJ, Ogunja JA (2018) Prevalence and risk factors associated with brucellosis among high-risk individuals in Lafia, Nasarawa state, Nigeria. *Int J One Hlth* 2018;4: 45-51.
 17. Esuruoso GO 1974. Bovine brucellosis in Nigeria. *Vet. Rec* 1974; 95(3): 54-58.
 18. Chukwu CC. 1987. Studies on seroprevalence of bovine brucellosis in Enugu and Nsukka, Nigeria. *Zariya Vet* 1987; 2: 25-30.
 19. Ocholi RA, Kalejaiye JO, Okewole PA. Brucellosis in Nigeria. A review. *Trop Vet* 1993; 11: 15-26.
 20. Junaidu AU, Oboegbulem SI, Sharubutu GH, Daneji AI. Brucellosis in camels (*Camelus dromedaries*) slaughtered in Sokoto, Northwestern Nigeria. *Anim Prod Res Adv* 2006; 2: 158-60.
 21. Junaidu AU, Oboegbulem SI, Salihu MD. Seroprevalence of brucellosis in prison farm in Sokoto State, Nigeria. *Asian J Epidemiol* 2008; 1(1): 24-8.
 22. Onunkwo JI, Njoga EO, Nwanta JA, Shoyinka SVO, Onyenwe IW, Eze JI. Serological survey of Porcine *Brucella* infection in South East, Nigeria. *Nig Vet J* 2011; 32: 60-62.
 23. Kauffman LK, Bjork JK, Gallup JM, Boggiatto PM, Bellaire BH, Petersen CA_2014. Early detection of *Brucella canis* via quantitative polymerase chain reaction analysis. *Zoon Pub Hlth* 2014; 61(1): 48-54.
 24. Lucero NE, Escobar GI, Ayala SM, Jacob N 2005. Diagnosis of human brucellosis caused by *Brucella canis*. *J Med Microbiol* 2005; 54: 457-61.
 25. Cadmus SIB, Adesokan HK, Ajala OO, Odetokun WO, Perrett LL, Stack JA. Seroprevalence of *Brucella abortus* and *B. canis* in household dogs in southwestern Nigeria: A preliminary report. *J South Afr Vet Assoc* 2011; 82: 56-57.
 26. Behzadi MA, Mogheiseh A. Outbreak investigation of brucellosis at a Kennel in Iran. *Pak Vet J* 2011; 31: 379-380.
 27. Awah-Ndukum J, Mouiche MMM, Kouonmo-Ngonyum L, Bayang HN, Manchang TK, Poueme RSN, Kouamo J, Ngu-Ngwa V, Assana E, Feussom KJM, Zoli AP. Seroprevalence and risk factors of brucellosis among slaughtered indigenous cattle, abattoir personnel and pregnant women in Ngaoundéré, Cameroon. *BMC Infect Dis* 2018; 18(1):611. <https://doi.org/10.1186/s12879-018-3522-x>.
 28. National Population Commission. Federal Republic of Nigeria Official Gazette, 2009, Abuja; 2006.
 29. Thursfield, M. *Veterinary epidemiology*. 3rd ed. Oxford: Blackwell; 2007.
 30. Adedoyin FJ, Adesokan HK, Otuh PI, Omobowale TO, Abiola JO, Stack JA *et al.* Sero-Prevalence of Brucellosis in Household and Stray Dogs in Ibadan, South Western Nigeria. Document taken from Proceedings of the 49th annual congress of the Nigerian Veterinary Medical Association; 2012 Nov 5-9; Ado- Ekiti, Ekiti State, Nigeria.
 31. Alton GG, Jones LM, Pietx DE 1975. Laboratory techniques in Brucellosis (Monograph Series No. 55). 2nd Ed. Geneva: FAO/WHO; 1975.
 32. Morgan WJB, Mackinon DJ, Gill KPW, Gover SGM, Norris PW. Standard laboratory techniques for the diagnosis of brucellosis. 2nd Ed. MAF. Weybridge, England: Central Veterinary Laboratory; 1978.
 33. Sati SN 2002. Seroepidemiological survey of *Brucella abortus* infection in Fulani breeding herds and trade cattle in the middle belt and southeast. (Ph.D thesis). Nsukka: University of Nigeria, Nsukka; 2002.
 34. Muhairwa AP, Mwakijungu EO, Msoffe PLM, Mtambo MMA. Seroprevalence and factors affecting canine mo-

- nocytic ehrlichiosis and canine brucellosis in Tanzania. *Res Opin Anim Vet Sci* 2012; 2: 181–188.
35. Chessbrough M. *District Laboratory Practice in Tropical Countries, Part 2*. Cambridge Low Price Edition, Cambridge: University Press; 2000.
 36. Adesokan HK, Alabi PI, Ogundipe MA 2016. Prevalence and predictors of risk factors for Brucellosis transmission by meat handlers and traditional healers' risk practices in Ibadan, Nigeria. *J Prev Med Hyg* 2016; 57(3): E164-71.
 37. Palomarez – Ressendiiz E, Arellano – Reynoso B, Hernandez – Castro R, Tenorio – Gutierrez V, Salas–Tellez E, Suarez–Guemes F, Diaz-Aparicio E. Immunogenic Response of *Brucella canis* virB10 and virB11 Mutant in Murine Model. *Front Cell Infect Microbiol* 2012; 2:35. <https://doi.org/10.3389/fcimb.2012.00035>.
 38. Momoh HA, Ijale GO, Ajogi I, Okolocha EC. Seroprevalence of Canine Brucellosis in Jos, Plateau State, Nigeria. *Asian J Epidemiol* 2014; 7(2): 36-42.
 39. Radostits OM, Gay CC, Hinchcliff KW, Constable PD. *Veterinary Medicine. A textbook of the diseases of cattle, horses, sheep, pigs, and goats*. Brucellosis. 10th ed., Edinburgh, London, New York, Oxford, Philadelphia, St. Louis, Sydney, Toronto: Saunders Elsevier; 2007.
 40. Adesiyun AA, Adbullahi SU, Adeyanju JB. Prevalence of *Brucella abortus* and *brucella canis* antibodies in dogs in Nigeria. *J Small Anim Pract* 1986; 27: 31-7.
 41. Abubakar M, Mansoor M, Arshed MJ. Bovine brucellosis: old and new concepts with Pakistan perspective. *Pak Vet J* 2012; 32: 147-55.
 42. Aulakh HK, Patil PK, Sharma S, Kumar H, Mahajan V, Sandhu KS. A Study on the Epidemiology of Bovine Brucellosis in Punjab (India) Using Milk-ELISA. *Acta Vet Brno* 2008; 77: 393-9.
 43. Abukakar M, Arshed MJ, Hussain M, Ehtisham-Ul-Haq, Ali Q. Serological evidence of *Brucella abortus* prevalence in Punjab province, Pakistan-a cross-sectional study. *Transbound Emerg Dis* 2010; 57: 443-7.
 44. Escobar GI, Boeri EJ, Ayala SM, Lucero NE. The feasibility of using antigens prepared with rough *Brucella* strains for diagnosis of canine brucellosis. *Rev Argent Microbiol* 2010; 42(1): 35-40.
 45. Johnson CA, Carter TD, Dunn JR, Baer SR, Schallow MM, Bellay YM, Guerra MA, Frank NA. Investigation and characterization of *Brucella canis* infections in pet-quality dogs and associated human exposures during a 2007–2016 outbreak in Michigan. *J Amer Vet Med Assoc* 2018; 253(3): 322-36.
 46. Keid LB, Soares RM, Vasconcellos SA, Megid J, Salgado VR, Richtzenhain LJ. Comparison of agar gel immunodiffusion test, rapid slide agglutination test, microbiological culture and PCR for the diagnosis of canine brucellosis. *Res Vet Sci* 2009; 86(1):22-6.
 47. Chinyoka S, Dhliwayo S, Marabini L, Dutlow K, Matope G, Pfukenyi DM. Serological survey of *Brucella canis* in dogs in urban Harare and selected rural communities in Zimbabwe. *J South Afr Vet Assoc* 2014; 85(1): <https://doi.org/10.4102/jsava.v85i1.1087>