

PRESENCE OF BOVINE LEUKEMIA VIRUS GENOTYPES 1 AND 3 IN ANTIOQUIA, COLOMBIA

PRESENCIA DE LOS GENOTIPOS 1 Y 3 DEL VIRUS DE LA LEUCOSIS BOVINA EN ANTIOQUIA, COLOMBIA

Cristina Úsuga-Monroy¹, Francisco J. Díaz², José Julián Echeverri-Zuluaga³,
Luis Gabriel González-Herrera⁴, Albeiro López-Herrera⁵

¹Zoot, M.Sc., Ph.D.(c), Faculty of Agricultural Sciences, Biodiversity and Molecular Genetics Group. Universidad Nacional de Colombia, Street 59A number 63-20, Medellín, Colombia, e-mail: cusugam@unal.edu.co; ²MD, Ph.D., Faculty of Medicine, Immunovirology Group. Universidad de Antioquia, Street 62 number 52-59, Medellín, Colombia, e-mail: francisco.diaz@udea.edu.co; ³Zoot, M.Sc, Ph.D., Faculty of Agricultural Sciences, Biodiversity and Molecular Genetics Group, Universidad Nacional de Colombia, Street 59A number 63-20, Medellín, Colombia, e-mail: jecheve@unal.edu.co; ⁴MVZ, M.Sc. Ph.D., Faculty of Agricultural Sciences, Biodiversity and Molecular Genetics Group, Universidad Nacional de Colombia, Street 59A number 63-20, Medellín, Colombia, e-mail: luggonzalezhe@unal.edu.co; ⁵Zoot, MV, M.Sc., Ph.D., Faculty of Agricultural Sciences, Biodiversity and Molecular Genetics Group. Universidad Nacional de Colombia, Street 59A number 63-20, Medellín, Colombia, e-mail: alherrera@unal.edu.co

Rev. U.D.C.A Act. & Div. Cient. 21(1): 119-126, Enero Junio 2018 - DOI:10.31910/rudca.v21.n1.2018.670

SUMMARY

Bovine leukemia virus (BLV) is an immunosuppressant retrovirus that primarily affects dairy livestock, its target cells are B lymphocytes in which it integrates its genome infecting cattle for life. It is important to identify the distribution of the BLV in the region and to reconstruct its evolutionary history through phylogenetic trees, for the province of Antioquia this is the first report of the BLV genotypes. The aim of this study was to identify the genotype of BLV circulating in dairy cattle of different regions of the province of Antioquia, Colombia. DNA was extracted from 8 Holstein cows. Nested PCR was performed to amplify a fragment of 444 pb of the *env* gene. The *env* viral gene codes for surface protein gp51, gene is highly conserved and it is used for phylogenetic analysis. Obtained amplicons were sequenced, manually aligned in MEGA V7 program, and compared to 53 viral *env* gene sequences registered in GenBank. Phylogenetic analysis was performed by Maximum Likelihood and Bayesian methods. Two circulating genotypes were found: the most common genotype was 1, found in seven samples; they grouped with sequences from EE. UU, Argentina and Japan; only one sample was classified as genotype 3 and was grouped with samples from EE. UU and Japan. At least two genotypes (1 and 3) of BLV are circulating in Antioquia; however, more cattle and herds should be evaluated to elucidate the diversity and distribution of BLV in Colombia.

Key words: Antioquia, BLV, Colombia, *Env* gene, Genotype, Phylogenetic analysis.

RESUMEN

El virus de la leucosis bovina (BLV) es un retrovirus inmunosupresor que afecta, principalmente, al ganado lechero; sus células diana son los linfocitos B, en los cuales, integra su genoma, infectando al ganado de por vida. Es importante identificar la distribución del BLV en la región y reconstruir su historia evolutiva, a través de árboles filogenéticos; para el departamento de Antioquia, este es el primer reporte de los genotipos del BLV. El objetivo de este estudio fue identificar el genotipo de BLV que circula en ganado lechero de diferentes regiones del departamento de Antioquia, Colombia. Se extrajo ADN de 8 vacas Holstein. Se realizó una PCR anidada, para amplificar un fragmento del gen *env* de 444 pb. El gen *env* viral codifica la proteína de superficie gp51, altamente conservado y es usado en análisis filogenéticos. Los amplicones obtenidos se secuenciaron, se alinearon manualmente en el programa MEGA V7 y se compararon con 53 secuencias del gen *env* viral, registradas en GenBank. El análisis filogenético, se realizó por métodos de Máxima Verosimilitud y Bayesianos. Se encontraron dos genotipos circulantes: el genotipo más común fue 1, hallado en siete muestras, agrupadas con secuencias de EE. UU, Argentina y Japón; solo una muestra se clasificó como genotipo 3 y se agrupó con muestras de EE. UU y Japón. Al menos dos genotipos (1 y 3) de BLV están circulando en Antioquia; sin embargo, se deben evaluar más bovinos y hatos para elucidar la diversidad y la distribución de BLV en Colombia.

Palabras clave: Análisis filogenético, Antioquia, BLV, Colombia, Genotipo, Gen *env*.

INTRODUCTION

Enzootic bovine leukemia (EBL) is a disease that primarily affects dairy cattle. Infected cattle show no clinical symptoms, but in most cases, they develop persistent lymphocytosis due to proliferation of B cells; in others, the infected bovine present the tumor form or lymphosarcomas (Aida *et al.* 2013). This disease causes losses in yield and reproductive levels in infected cattle; studies have shown that herds infected with BLV had a decrease in milk production from 2,5 to 28,14% compared to the production of the uninfected herd (Emanuelsson *et al.* 1992; Nekouei *et al.* 2016) and increased cull rate, as well as susceptibility to other infectious diseases as mastitis (Kakinuma *et al.* 2014), diarrhea and pneumonia (Emanuelsson *et al.* 1992). The etiological agent of EBL is the bovine leukemia virus (BLV), a retrovirus belonging to the *Deltaretrovirus* genus (Wu *et al.* 2003). BLV is related, from structural and functional point of view, with human T-cell lymphotropic viruses (HTLV-1, HTLV-2 and HTLV-3) and simian T-cell lymphotropic virus (STLV-1) (Hulo *et al.* 2011). BLV has become a very important tool for HTLV-1 study (Gillet *et al.* 2007; Aida *et al.* 2013), since the infection produced by both viruses is similar, affecting either T-cells (HTLV) or B-cells (BLV), none of them produce detectable viremia and their genomic organization is the same. BLV genome is a single stranded RNA of positive sense, consists of three genes: *gag*, *pol* and *env*, encoding different proteins: p15, p12 and p24 proteins are codified by *gag* gene; integrase enzyme (IN) and reverse transcriptase (RT) enzyme are codified by *pol* gene; gp51 and gp30 glycoproteins are codified by *env* gene (Gillet *et al.* 2007); moreover, BLV has a *pX* region that codifies for different regulatory and incidental proteins (tax, rex, G4, G3). Among them viral transcription activators (tax), viral mRNA exporters, and structural protein expression (rex), high viral load maintenance proteins (G4), and low viral load proteins (R3) (Gillet *et al.* 2007). *Env* gene is highly conserved and the antigen gp51 is generally present in all animals during different stages of infection (Baruta *et al.* 2011). Variation in *env* gene sequence have allowed to identify nine genotypes of BLV (Polat *et al.* 2016). The most prevalent genotypes worldwide are 1, 2 and 3 (Inoue *et al.* 2011). In America BLV is widely distributed and these genotypes have been found: in the United States, 1 and 2; in Costa Rica, 1 and 5 (Zhao *et al.* 2007); in Brazil, 1, 2 and 6 (Camargos *et al.* 2002); in Chile, 7 (Felmer *et al.* 2005); in Uruguay, 1 (Moratorio *et al.* 2010), and in Bolivia, 9 (Polat *et al.* 2016). The aim of this study was to identify the genotypes of BLV in different municipalities of Antioquia, Colombia.

MATERIALS AND METHODS

Animals and samples: This research is descriptive and only the population sample was evaluated in a moment of time. The sequenced amplicons were selected according to the

municipality of origin. This investigation was of exploratory type and therefore it tries to give a first approximation to the possible genotypes that circulate in the dairy herds of the province of Antioquia. Blood samples were drawn from 500 Holstein cows, which were located in herds of six municipalities of Antioquia: Bello, Belmira, Entreríos, Medellín, Rionegro and San Pedro de los Milagros. Sampling was approved by the ethics committee of the National University of Colombia (CEMED-022, July 13, 2015).

DNA extraction: Sampling was done with an 18G needle with vacuum vacutainer system (DBvacutainer®) and EDTA as an anticoagulant; samples were homogenized by inversion and transferred to the laboratory under refrigeration for DNA extraction. DNA from samples was obtained by the *salting out* technique (Miller *et al.* 1988), re-suspended in TE 1X pH 8.0 (Tris HCl 1M and EDTA 0.5M) and stored at 4°C until analysis. The quality and quantity of DNA obtained was evaluated in a spectrophotometer (NanoDrop 2000®, Massachusetts-USA) and agarose gel 1%.

Nested PCR: A region of the viral *env* gene (gp51) was amplified to obtain a 444bp fragment with the primers reported by Beier *et al.* (2001). The first round of PCR was performed in a final volume of 25µl with 150ng DNA, 3.0µl of 10mM of each primer *env* 5023 (5'-TCTGTGCCAAGTCTCCAGATA-3') and *env* 5608 (5'-AACAACAACCTCTGGGGAGGGT-3'), 0.4mM dNTPs, 1X buffer PCR (ThermoScientific®), 3mM MgCl₂ and 1U Taq Polymerase DNA. In the second PCR reaction 5µl of the first amplification PCR product was used as template DNA, with same concentrations of other reagents and *env* primers 5099 (5'-CCCACAAGGGCGGCGCCGTTT-3') and *env* 5521 (5'-GCGAGGCCGCGGTCCAGAGCTGG-3') in a final volume of 30µL. Reactions for both PCR were identical, and were performed in a T3 thermocycler (Biometra®) with the following protocol: initial denaturation at 94°C for 5 minutes, followed by 40 cycles of 94°C for 30 seconds, 60°C for 30 seconds, and 72°C for 1 minute; the final extension was run at 72°C for 5 minutes. As a negative and positive control, we run PCR reactions without template DNA and with DNA of a cow that tested positive in a previous study (Úsuga-Monroy, *et al.* 2015), respectively. The product of the second reaction was checked on an agarose gel stained with 2% EZ-VISION (Amresco®) in a Gel Doc (BioRad, California-USA).

Sequencing and phylogenetic analysis: Eight amplicons sent for sequencing in a commercial facility (Macrogen Inc. Korea); these amplicons were from different herds located at these municipalities in the province of Antioquia: Bello (1), Belmira (2), Entreríos (1), Medellín (2) Rionegro (1), and San Pedro de Los Milagros (1). The nucleotide sequences of the samples were compared with 53 partial sequences of the viral *env* gene registered in GenBank representative of the 9 BLV

genotypes and representing different geographic regions including several South American countries. Sequences were manually aligned in the software MEGA V7. Phylogenetic analysis was performed using two methods. First, a Maximum Likelihood analysis using MEGA V7 was performed. The Kimura 2-parameters with Gamma distribution (K2+G) was best substitution model based on the Bayesian Information Criterion (BIC). Bootstrap values were determined with 1000 repetitions and only significant values over 70% were considered as significant. A second phylogenetic tree was built by Bayesian methods using MrBayes V3.2.2. The evolution model used was Kimura 2-parameters with Gamma (K2+G) distribution. Convergence (average standard deviation of split sequences) of 0.005 was obtained after 1.000.000 generations. Trees were sampled every 1000 generations. The consensus tree was obtained after discarding (burn in) the initial 25% of the generations. The final tree was edited in FigTree V1.4.0.

RESULTS AND DISCUSSION

PCR is highly sensitive methodology of direct diagnosis for infection with BLV (Rola *et al.* 2002). PCR helps to rationally direct diagnostic tests for early diagnosis and disease control plans. The BLV is widely distributed in the world, probably migrated from Europe to America through infected cattle. It is considered that in America the disease is enzootic. Molecular prevalence for bovine leukemia virus in this study was 44% of infection, other South American countries show lower molecular prevalences as for example Chile (27,9%), Venezuela (33,3%), Bolivia (30,7%) and Peru (42,3%); however, countries such as Paraguay (54,7%), Brazil (60,8%) and Argentina (77,4%) have higher molecular prevalences (Polat *et al.* 2017).

The most studied BLV gene is the envelope gene, which translates for the gp51 glycoprotein. The gp51 has an essential role in the life cycle of the virus, is required to enter the cell and as a target of neutralizing antibodies. The BLV clusters have been carried out through the phylogenetic analysis of a 444pb fragment of the envelope gene (*env*) of the virus (Beier *et al.* 2001), these studies allow to identify the distribution of BLV in the region and to reconstruct its evolutionary history. In addition, small differences have been demonstrated in many ways for the gp51 mutations which are related to the geographic location of the sample, allowing to group the BLV in ten genotypes (Polat *et al.* 2016). Proviral BLV DNA was detected in 219 samples and 281 specimens tested negative by nested PCR. Eight positive PCR products were selected for sequencing. Their sequences were aligned to 53 sequences of the *env* gene from the following countries: United States, Costa Rica (North and Central America), Argentina, Brazil, Uruguay, Paraguay, Peru, Bolivia, Chile (South America), Germany, Russia, Belgium, Poland, Ukraine,

Croatia (Europe), Japan, (Asia) and Australia (Oceania). In the Maximum Likelihood phylogenetic the sequences from Bello, Rionegro, San Pedro, Entreríos, Medellín and one sequence from Belmira were grouped to isolates from United States, Argentina and Japan in genotype 1, with a branch support of 79%. One out of the eight sequences analyzed from the Belmira was cluster within genotype 3, with sequences from United States and Japan; the branch support for this clade was 97%. Thus, the municipality of Belmira showed both genotypes, 1 and 3 (Figure 1). A phylogenetic analysis was also performed by Bayesian methods in which 7 out of the 8 sequences analyzed were also clustered in genotype 1, and only one sequence from the municipality of Belmira was grouped into genotype 3; branch support values, expressed in posterior probabilities, were 100% in both clades (Figure 2). The results of both phylogenetic reconstructions obtained by Maximum Likelihood and Bayesian methods were similar.

BLV sequences BELMIRA_466, MEDELLÍN_0407, MEDELLÍN_722, RIONEGRO_512 are the identical using p-distance model, although they are geographically remote municipalities. BELMIRA_847 has a higher proportion of nucleotide sites different than other samples and is farther from BELLO_364 sequence (Table 1). ENTRERRÍOS_026 sequence formed a clade with M35242 sequence from the United States, with a posterior probability value of 83% for Bayesian phylogenetic tree. BELMIRA_847 sequence was the only one grouped in genotype 3 along with sequences from the United States and Japan with a posterior probability value of 100% for Bayesian phylogenetic tree (Figure 2) and branch supporting value of 97% for the phylogenetic tree by Maximum Likelihood (Figure 1).

Maximum Likelihood analysis and Bayesian analysis grouped one sample from the municipality of Belmira in genotype 3, while the remaining 7 samples were grouped in genotype 1. The municipality of Belmira was the only one presenting both genotypes 1 and 3 in samples from the same municipality. The presence of the two genotypes (1 and 3) in the municipality of Belmira indicates that the BLV has probably entered the country through the purchase of livestock and import of artificial insemination straws on several occasions, other studies have also reported the presence of two or more genotypes in one herd (Ababneh *et al.* 2012; Mekata *et al.* 2015). Genotypes 1 and 3 are circulating in specialized dairy farms in Antioquia, since there is no limitation for viral transmission. Recently the viral load has been linked to horizontal transmission because a cow infected with less than 3 copies of the virus per 100 cells cannot spread BLV to other cattle of more than 30 months old (Mekata *et al.* 2015). Transmission of the virus can be vertical and horizontal. In vertical transmission colostrum is a vehicle for the virus (Nagy *et al.* 2007), also exosomes in milk from infected cows contain BLV proteins, they are transmitted by an alternative




Figure 1. Phylogenetic tree constructed by Maximum Likelihood using as substitution model Kimura 2-parameters with Gamma distribution (K2+G). Bootstrap values are indicated as data from 1000 replicas (Bar 0.005 substitutions per site) based on partial sequences of 400 nucleotides of BLV *env* gene. Colombian sequences are indicated as (●) and were compared with 53 reported in GenBank sequences for the *env* gene. Genotype, access numbers and country of origin of the sequences indicated.




Figure 2. Bayesian phylogenetic tree using Kimura 2-parameters with Gamma distribution (K2+G). Posterior probability values are indicated as numbers on the bar. The analysis is based on partial sequences of 400 nucleotides of BLV *env* gene. Colombian sequences are indicated as (●) and were compared with 53 reported in GenBank for *env* gene. Genotype, access numbers and country of origin of the sequences indicated.

Table 1. Estimates of Evolutionary Divergence between Sequences. The number of base differences per site from between sequences are shown. The analysis involved 8 nucleotide sequences. Codon positions included were 1st+2nd+3rd+Noncoding. All positions with less than 95% site coverage were eliminated. That is, fewer than 5% alignment gaps, missing data, and ambiguous bases were allowed at any position. There were a total of 400 positions in the final dataset. Evolutionary analyses were conducted in MEGA V7.

Sequences	1	2	3	4	5	6	7	8
1. BELLO_364								
2. BELMIRA_466	0.005							
3. BELMIRA_847	0.030	0.025						
4. ENTRERRIOS_026	0.008	0.003	0.028					
5. MEDELLIN_0407	0.005	0.000	0.025	0.003				
6. MEDELLIN_722	0.005	0.000	0.025	0.003	0.000			
7. RIONEGRO_512	0.005	0.000	0.025	0.003	0.000	0.000		
8. SANPEDRO_102	0.008	0.003	0.028	0.005	0.003	0.003	0.003	

route different from the viral infection and have an important role in removing proteins through BLV infected cells (Yamada *et al.* 2013). In the horizontal transmission reusing needles for vaccination and reuse of gloves during palpation or dehorning (Gillet *et al.* 2007; Ortega *et al.* 2016) help to spread the virus. Infection through semen is also included in horizontal transmission (Dus Santos *et al.* 2007).

The most prevalent genotypes BLV worldwide are 1, 2 and 3 (Inoue *et al.* 2011). In America, the following genotypes are found: genotypes 1, 3 and 4 are found in the United States, 1 and 5 are in Costa Rica (Zhao *et al.* 2007), 1, 2, 5, 6 and 7 are in Brazil (Camargos *et al.* 2002; Moratorio *et al.* 2010), 4 and 7 in Chile (Felmer *et al.* 2005), 1 in Uruguay (Moratorio *et al.* 2010), 1, 2, 4 and 6 in Argentina (Rodriguez *et al.* 2009), 1, 2 and 6 in Perú (Polat *et al.* 2016), 1, 2 and 6 in Paraguay (Polat *et al.* 2016) and 1, 2, 6 and 9 in Bolivia (Polat *et al.* 2016). Genotypes 1 and 2 were reported in dairy cattle in the province of Nariño, Colombia (Benavides *et al.* 2017), while in our study in the province of Antioquia, genotypes 1 and 3 were found in Holstein cows. According to Inoue *et al.* (2011) the most frequent BLV genotypes are 1, 2 and 3. These data suggest that genotype 1 is the most widespread, it is also the most frequent detected in samples from Antioquia, Colombia in this research. For now, no relationship has been reported between genotype and virulence or resistance, this may be due to the few changes at the nucleotide level that the virus presents. The low mutation rate of BLV is due to the long generation time of *Deltaretrovirus*. BLV mutation rate is 1.7×10^{-5} s/n/c is similar to the human T-cell lymphotropic virus (HTLV-1) 1.6×10^{-5} s/n/c mutation rate (Sanjuán *et al.* 2010). Bovine leukemia virus genotypes 1 and 3 are circulating in dairy herds in the province of Antioquia, Colombia. The

two genotypes can be found in the same municipality. It is important to maintain good iatrogenic practices such as: do not reuse needles, palpation gloves, do not use colostrum or milk from infected mothers in healthy calves. In this way, the dispersion of viral particles within and between herds can be controlled. On the other hand, it is important to evaluate more dairy herds and other fragments of the viral genome to establish genetic variability analysis for BLV and to have a better focus about the distribution of the virus in the dairy herds of the department and the country.

Conflict of interest: This manuscript was prepared and reviewed with the participation of all the authors, who declare that they have no conflict of interest that compromise the validity of the results. **Financing:** Research Directorate of the National University of Colombia, Medellín. Project Quipu Code 201010012967.

BIBLIOGRAPHY

1. ABABNEH, M.M.; AL-RUKIBAT, R.K.; HANANEH, W.M.; NASAR, A.T.; AL-ZGHOUL, M.B. 2012. Detection and molecular characterization of bovine leukemia viruses from Jordan. Arch Virol. 157(12):2343-2348.
2. AIDA, Y.; MURAKAMI, H.; TAKAHASHI, M.; TAKESHIMA, S. 2013. Mechanisms of pathogenesis induced by bovine leukemia virus as a model for human T-cell leukemia virus. Front Microbiol. 4(328):1-11.
3. BARJTA, D.A.; ARDOINO, S.M.; BRANDAN, J.L.; SOSA, R.E.; MARIANI, E.L.; ALBRETCH, E.M. 2011. Leucosis Bovina Enzoótica. Ciencia Veterinaria. 13(1):9-14.

4. BEIER, D.; BLANKENSTAIN, P.; MARQUARD, O.; KUZMAK, J. 2001. Identification of different BLV provirus isolates by PCR, RFLPA and DNA sequencing. *Berl. Münch. Tierarztl. Wochenschr.* 114(7-8):252-256.
5. BENAVIDES, B.; MUÑOZ, S.; CERIANI, C. 2017. Molecular analysis of a fragment of bovine leukemia virus env gene by Nested-PCR in dairy cows from Pasto, Nariño. *Rev. Med. Vet.* 33:67-75.
6. CAMARGOS, M.F.; STANCEK, D.; ROCHA, M.A.; LESSA, L.M.; REIS, J.K.; LEITE, R.C. 2002. Partial sequencing of env gene of bovine leukaemia virus from Brazilian samples and phylogenetic analysis. *J. Vet. Med. B Infect. Dis. Vet. Public Health.* 49(7):325-331.
7. DUS SANTOS, M.J.; TRONO, K.; LAGER, I.; WIGDOROVITZ, A. 2007. Development of a PCR to diagnose BLV genome in frozen semen samples. *Vet. Microbiol.* 119(1):10-18.
8. EMANUELSSON, U.; SCHERLING, K.; PETTERSSON, H. 1992. Relationships between herd bovine leukemia virus infection status and reproduction, disease incidence, and productivity in Swedish dairy herds. *Preventive Veterinary Medicine.* 12:121-131.
9. FELMER, R.; MUÑOZ, G.; ZÚÑIGA, J.; RECABAL, M. 2005. Molecular analysis of a 444 bp fragment of the bovine leukaemia virus gp51 env gene reveals a high frequency of non-silent point mutations and suggests the presence of two subgroups of BLV in Chile. *Vet. Microbiol.* 108(1-2):39-47.
10. GILLET, N.; FLORINS, A.; BOXUS, M.; BURTEAU, C.; NIGRO, A.; VANDERMEERS, F.; BALON, H.; BOUZAR, A.; DEFOICHE, J.; BURNY, A.; REICHERT, M.; KETTMANN, R.; WILLEMS, L. 2007 Mechanisms of leukemogenesis induced by bovine leukemia virus: prospects for novel anti-retroviral therapies in human. *Retrovirology.* 4(18):1-32.
11. HUJO, C.; DE CASTRO, E.; MASSON, P.; BOUGUELERET, L.; BAIROCH, A.; XENARIOS, I.; LE MERCIER, P. 2011. ViralZone: a knowledge resource to understand virus diversity. *Nucleic Acids Res.* 39:576-582.
12. INOUE, E.; MATSUMURA, K.; MAEKAWA, K.; NAGATSUKA, K.; NOBUTA, M.; HIRATA, M.; MINAGAWA, A.; OSAWA, Y.; OKAZAKI, K. 2011. Genetic heterogeneity among bovine leukemia viruses in Japan and their relationship to leukemogenicity. *Arch. Virol.* 156(7):1137-1141.
13. KAKINUMA, S.; MAEDA, Y.; OHTSUKA, H.; KONNAI, S.; OIKAWA, M. 2014. Bovine Leukemia virus titer and leukocyte population associated with mastitis in periparturient dairy cows. *Intern. J. Appl. Res. Vet. Med.* 12(3):239-244.
14. MEKATA, H.; SEKIGUCHI, S.; KONNAI, S.; KIRINO, Y.; HORII, Y.; NORIMINE, J. 2015. Horizontal transmission and phylogenetic analysis of bovine leukemia virus in two districts of Miyazaki, Japan. *J. Vet. Med. Sci.* 77(9):1115-1120.
15. MILLER, S.A.; DYKES, D.D.; POLESKY, H.F. 1988. A simple salting out procedure for extracting ADN from human nucleated cells. *Nucleic Acids Res.* 16(3):1215.
16. MORATORIO, G.; OBAL, G.; DUBRA, A.; CORREA, A.; BIANCHI, S.; BUSCHIAZZO, A.; CRISTINA, J.; PRITSCH, O. 2010. Phylogenetic analysis of bovine leukemia viruses isolated in South America reveals diversification in seven distinct genotypes. *Arch. Virol.* 155(4):481-489.
17. NAGY, D.W.; TYLER, J.W.; KLEIBOEKER, S.B. 2007. Decreased Periparturient Transmission of Bovine Leukosis Virus in Colostrum-Fed Calves. *J. Vet. Intern. Med.* 21(5):1104-1107.
18. NEKOUEI, O.; VANLEEUWEN, J.; STRYHN, H.; KELTON, D.; KEEFE, G. 2016. Lifetime effects of infection with bovine leukemia virus on longevity and milk production of dairy cows. *Prev. Vet. Med.* 133:1-9.
19. ORTEGA, D.; SÁNCHEZ, A.; TOBÓN, J.; CHAPARRO, Y.; CORTÉS, S.; GUTIÉRREZ, M. 2016. Seroprevalence and risk factors associated with bovine leukemia virus in Colombia. *J. Vet. Med. Anim. Health.* 8(5):35-43.
20. POLAT, M.; TAKESHIMA, S.N.; HOSOMICHI, K.; KIM, J.; MIYASAKA, T.; YAMADA, K.; ARAINGA, M.; MURAKAMI, T.; MATSUMOTO, Y.; DE LA BARRA DIAZ, V.; PANEI, C.J.; GONZÁLEZ, E.T.; KANEMAKI, M.; ONUMA, M.; GIOVAMBATTISTA, G.; AIDA, Y. 2016. A new genotype of bovine leukemia virus in South America identified by NGS based whole genome sequencing and molecular evolutionary genetic analysis. *Retrovirology.* 13(4):1-23.

21. POLAT, M.; TAKESHIMA, S.N.; AIDA, Y. 2017. Epidemiology and genetic diversity of bovine leukemia virus. *Virol. J.* 14(1): 1-16.
22. RODRIGUEZ, S.M.; GOLEMBA, M.D.; CAMPOS, R.H.; TRONO, K.; JONES, L.R. 2009. Bovine leukemia virus can be classified into seven genotypes: evidence for the existence of two novel clades. *J. Gen. Virol.* 90(11): 2788-2797.
23. ROLA, M.; KUZMAK, J. 2002. The detection of bovine leukemia virus proviral DNA by PCR-ELISA. *J. Virol. Methods.* 99(1-2): 33-40.
24. SANJUÁN, R.; NEBOT, M.R.; CHIRICO, N.; MANSKY, L.M.; BELSHAW, R. 2010. Viral Mutation Rates. *J. Virol.* 84(19):9733-9748.
25. ÚSUGA-MONROY, C.; ECHEVERRI, J; LÓPEZ-HERRERA, H. 2015. Diagnóstico molecular del virus de leucosis bovina en una población de vacas Holstein, Colombia. *Arch. Zootec.* 64(248):383-388.
26. WU, D.; MURAKAMI, K.; MOROOKA, A.; JIN, H.; INOSHIMA, Y.; SENTSU, H. 2003. In vivo transcription of bovine leukemia virus and bovine immunodeficiency-like virus. *Virus Res.* 97(2):81-87.
27. YAMADA, T.; SHIGEMURA, H.; ISHIGURO, N.; INOSHIMA, Y. 2013. Cell Infectivity in Relation to Bovine Leukemia Virus gp51 and p24 in Bovine Milk Exosomes. *PloS One.* 8(10):1-7.
28. ZHAO, X.; BUEHRING, G.C. 2007. Natural genetic variations in bovine leukemia virus envelope gene: possible effects of selection and escape. *Virology.* 366(1):150-165.

Recibido: Noviembre 28 de 2017

Aceptado: Marzo 8 de 2018

Cómo citar:

Úsuga-Monroy, C.; Díaz, F.J.; Echeverri-Zuluaga, J.J.; González-Herrera, L.G.; López-Herrera, A. 2018. Presence of bovine leukemia virus genotypes 1 and 3 in Antioquia, Colombia. *Rev. U.D.C.A Act. & Div. Cient.* 21(1): 119-126. DOI:10.31910/rudca.v21.n1.2018.670