

# On the Activation of Quantum Nonlocality

Andrés Felipe Ducuara<sup>1,2,\*</sup>, Javier Madroñero<sup>1,2</sup>, John Henry Reina<sup>1,2</sup>

Edited by

Juan Carlos Salcedo-Reyes  
(salcedo.juan@javeriana.edu.co)

1. Departamento de Física, Universidad del Valle, A.A. 25360, Cali, Colombia.

2. Centre for Bioinformatics and Photonics-CIBioFi, Calle 13 No. 100-00, Edificio 320, No. 1069, Cali, Colombia

\* andres.ducuara@correounivalle.edu.co

Received: 12-06-2015

Accepted: 18-11-2015

Published on line: 16-05-2016

**Citation:** Ducuara AF, Madroñero J, Reina JH. On the Activation of Quantum Nonlocality, *Universitas Scientiarum*, 21 (2): 129-158, 2016.  
doi: 10.11144/Javeriana.SC21-2.otao

**Funding:** COLCIENCIAS, Fondo CTI-SGR and Universidad del Valle

**Electronic supplementary material:**  
N/A



## Abstract

We report on some quantum properties of physical systems, namely, entanglement, nonlocality,  $k$ -copy nonlocality (superactivation of nonlocality), hidden nonlocality (activation of nonlocality through local filtering) and the activation of nonlocality through tensoring and local filtering. The aim of this work is two-fold. First, we provide a review of the numerical procedures that must be followed in order to calculate the aforementioned properties, in particular, for any two-qubit system, and reproduce the bounds for two-qudit Werner states. Second, we use such numerical tools to calculate new bounds of these properties for two-qudit Isotropic states and two-qubit Hirsch states.

**Keywords:** Qubits; quantum information; quantum nonlocality; entanglement.

## Introduction

The understanding and classification of the properties of quantum states are important subjects from both fundamental and practical points of view (Horodecki et al. 2009, Brunner et al. 2014). *Entanglement* (Horodecki et al. 2009) and *nonlocality* (Brunner et al. 2014) are useful resources for quantum protocols, namely: quantum teleportation (Bennett et al. 1993) and cryptography (Ekert 1991). However, a deeper understanding of the relationship between them is still required (Brunner et al. 2014). Even though entanglement is necessary in order to achieve nonlocality, these two properties are not equivalent, i. e., there exist entangled *local* states (Werner 1989). Since several quantum protocols exclusively make use of nonlocality as a resource (Acín et al. 2006a), it is natural to ask whether it is possible to use these (apparently

useless) entangled local states in order to achieve (activate) nonlocality by means of processes termed *activation scenarios*. There are three of these mechanisms: *local filtering* (Popescu 1995), *tensoring* (Navascués & Vértesi 2011), and *quantum networks*. Additionally, it is possible to consider combinations of them (Brunner et al. 2014).

In this work, we begin by briefly reviewing the aforementioned activation scenarios and pay particular attention to the following: (1) the complete characterisation of *hidden nonlocality* (or activation through local filtering) for two-qubit systems recently derived in Pal & Ghosh (2015), (2) a case of the activation through tensoring called *k-copy nonlocality* (or *superactivation of nonlocality*) (Palazuelos 2012, Cavalcanti et al. 2013), and (3) activation through the combination of *tensoring* and *local filtering* (Masanes et al. 2008, Liang et al. 2012). Moreover, we have focused (though not restricted) on the study of these three scenarios over two-qubit systems. The latter could be of interest for future studies on states coming from the dynamics of open quantum systems where entanglement and nonlocality (restricted to the standard definition) are usually investigated (Batle & Casas 2010, Batle & Casas 2011). We then used the already mentioned tools to perform numerical simulations in order to investigate these quantum properties for some states of interest. We first reproduced the bounds for the so-called two-qudit Werner states. We then reported new bounds of these properties for the so-called two-qudit Isotropic states and two-qubit Hirsch states.

This work is organised as follows. The first two sections deal with entanglement and nonlocality. The third section establishes the motivation underneath the activation of nonlocality and consequently, we present an overview of the currently known activation scenarios. Next, we address hidden nonlocality, followed by *k*-copy nonlocality, and the activation of nonlocality through tensoring and local filtering. In the fourth section, we report results regarding the aforementioned quantum properties for some states of interest. First, we reproduced the bounds of these properties for two-qudit Werner states. Second, we report new bounds on the activation through tensoring and local filtering for two-qudit Isotropic states. Third, we report new bounds regarding hidden nonlocality and activation through tensoring and local filtering for two-qubit Hirsch states. Finally, we discuss the obtained results.

## Entanglement

Quantum entanglement was first implicitly introduced in the seminal 1935 EPR article (Einstein et al. 1935) and subsequent discussions, however, it had to wait until 1989 to be formally defined (Werner 1989). A general finite-dimensional bipartite *AB* system is represented by a density matrix or quantum state

$\rho \in D(\mathbb{C}^{d_A} \otimes \mathbb{C}^{d_B})$ , with  $d_A, d_B \geq 2$ , where  $D(\mathbb{H})$  stands for the set of density matrices of the complex Hilbert space  $\mathbb{H}$  or  $D(\mathbb{H}) := \{\rho \in PSD(\mathbb{H}) \mid \text{Tr}(\rho) = 1\}$ , with  $PSD$  the set of *positive semidefinite* complex matrices, that is the matrices  $\rho$  such that  $\forall |\phi\rangle \in \mathbb{H} : \langle \phi | \rho | \phi \rangle \geq 0$ . A general state there can be written as representing an ensemble of pure quantum states  $\{|\psi_i\rangle, p_i\}$ , with  $p_i > 0 \ \forall i$ , and  $\sum_i p_i = 1$  as:

$$\rho = \sum_i p_i \psi_i, \quad (1)$$

with  $|\psi_i\rangle := |\psi_i\rangle \langle \psi_i|$ . We say  $\rho$  is *separable* if:

$$\rho = \sum_i p_i \rho_A \otimes \rho_B, \quad (2)$$

otherwise it is *entangled*. Given a quantum state, it is not a trivial task to know whether it is possible to decompose it as in Eq. (2): there are criteria for entanglement quantification that work quite well for two-qubit systems, and good measures to quantify this are already in place. This said, we lack a unified generalisation to arbitrary high-dimensional and multipartite systems. We next address the main quantifier for two-qubit systems, namely, the *entanglement of formation* (EoF) (Bennett et al. 1996b). The EoF has a compact analytical expression for two qubits (Wootters 1998) and for a couple of bipartite high-dimensional states we are interested in, namely, the Werner and the Isotropic states (Terhal & Vollbrecht 2001, Vollbrecht & Werner 2001, Wootters 2001).

**Entanglement of Formation (EoF):** For pure bipartite states  $\psi = |\psi\rangle \langle \psi|$ , there is an entanglement measure free of ambiguity (in the sense that it is an if and only if criterion) in terms of the well known *von Neumann entropy*,  $E(\psi) := S(\psi_A) = -\text{Tr}(\psi_A \log_2 \psi_A)$  with  $\text{Tr}_A := \text{Tr}_B(\psi)$  the partial trace (Bennett 1996b). For general mixed states Eq. (1), it is natural to ask about the possible generalisation  $\sum_i p_i E(\psi_i)$ . However, since we have infinite possible ensemble decompositions Eq. (1), this definition will depend on the chosen ensemble. Consequently, the following measure, known as the *Entanglement of Formation* (EoF), has been introduced (Bennett 1996b):

$$E(\rho) := \min_{\{\psi_i, p_i\}} \left[ \sum_i p_i E(\psi_i) \right]. \quad (3)$$

In 1998, an analytical expression of Eq. (3) for the particular case of two-qubit systems was derived, (Wootters 1998). Given  $\rho \in D(\mathbb{C}^2 \otimes \mathbb{C}^2)$ , its EoF is given by:

$$EoF(\rho) = h \left( \frac{1 + \sqrt{1 - C(\rho)^2}}{2} \right), \quad (4)$$


where  $h(x) := -x\log_2 x - (1-x)\log_2(1-x)$  is the so-called binary entropy,  $C$  is the so-called concurrence,  $C(\rho) = \max\{0, \lambda_1 - \lambda_2 - \lambda_3 - \lambda_4\}$ , where the  $\lambda_i$ 's are the square roots of the eigenvalues of the product matrix  $\rho\hat{\rho}$ , in decreasing order, with:  $\hat{\rho} = (\sigma_y \otimes \sigma_y) \rho^* (\sigma_y \otimes \sigma_y)$ ,  $\rho^*$  the complex conjugate of  $\rho$  and,  $\sigma_y$  the Pauli matrix. Both,  $C(\rho)$  and  $EoF(\rho)$  go from 0 to 1. There is no known compact analytical expression for general high-dimensional systems, except for a couple of classes of two-qudit states; the so-called Werner and Isotropic states (Terhal & Vollbrecht 2001, Vollbrecht & Werner 2000, Wootters 2001). Next, we focus on the concept of nonlocality.

## Nonlocality

Quantum nonlocality was first implicitly introduced in the seminal EPR article (Einstein et al. 1935) and corresponding subsequent discussions. However, it had to wait until 1964 in order to be formally defined by J. S. Bell (Bell 1964). We proceed as follows: Given a bipartite system  $\rho \in D(\mathbb{C}^{d_A} \otimes \mathbb{C}^{d_B})$ , the first party of the whole system is sent to experimentalist A (Alice), and the second to experimentalist B (Bob). They want to study the two observables  $A$  and  $B$  respectively. These can be written in their spectral decomposition as  $A = \sum_a o_a P_a^x$ ,  $B = \sum_b o_b P_b^y$ , where:  $M^x := \{P_a^x\}$  and  $M^y := \{P_b^y\}$  are sets of projections, and  $o_a, o_b$  their respective eigenvalue spectra. However, they can also be written more generally as  $A = \sum_a o'_a E_a^x$ ,  $B = \sum_b o'_b E_b^y$ , where:  $M^x := \{E_a^x\}$  and  $M^y := \{E_b^y\}$  are now Positive Operators Valued Measurements (POVM's) with  $o'_a, o'_b$  real numbers though not necessarily eigenvalues of  $A$  and  $B$ . Using this quantum state  $\rho$ , Alice and Bob make measurements  $x$  and  $y$ , obtaining outcomes  $a$  and  $b$ . After repeating the experiment enough times, eventually, they are able to establish the probability of obtaining outcomes  $a, b$  after measuring  $x$  and  $y$ . This can be seen as the joint conditional probability function  $p(a, b|x, y, \rho)$ . According to quantum mechanics, we could reproduce this statistics by means of the Born Rule (Nielsen & Chuang 2000):

$$p_Q(a, b|x, y, \rho) := \text{Tr} [(E_a^x \otimes E_b^y) \rho]. \quad (5)$$

Since quantum mechanics has been proven correct so far, we have:  $p(a, b|x, y, \rho) = p_Q(a, b|x, y, \rho)$ . We can see a sketch of this process in **Figure 1**.



**Fig.1.** Schematics of a standard Bell test (Brunner et al. 2014). Experimentalists Alice and Bob, share a bipartite quantum state  $\rho$ . They make measurements  $x$  and  $y$  and obtain outcomes  $a$  and  $b$  respectively.

Bell introduced the notion of locality (Bell 1964) within a formalism covering theories that could reproduce this very same statistics Eq. (5). In this formalism there are probability functions  $\mu$  and  $\xi$ , such that

$$p_L(a, b|x, y, \rho) = \int_{\Lambda} d\lambda \mu(\lambda|\rho) \xi(a, b|x, y, \lambda), \quad (6)$$

where  $\Lambda$  is the often-called *set of hidden variables* and the triplet  $(\Lambda, \mu, \xi)$  is the *ontological model*. The *locality* condition reads  $\xi(a, b|x, y, \lambda) = \xi_A(a|x, \lambda) \xi_B(b|y, \lambda)$ . Mathematically speaking, it means that the functions  $\xi_A$  and  $\xi_B$  are probabilistically independent on  $x$  and  $y$ . Physically speaking, it means that after receiving inputs  $x$  and  $y$ , Alice cannot influence Bob's output and vice versa. Then, the question that arises is the following: Is it possible to reproduce the statistics given by Eq. (5) by means of Eq. (6)? i. e.,

$$p_Q(a, b|x, y, \rho) \stackrel{?}{=} p_L(a, b|x, y, \rho). \quad (7)$$

If this is indeed possible for any POVM (we could relax the condition to just projections), the quantum state  $\rho$  will be called *local*, otherwise  $\rho$  will be *nonlocal*. For instance, it is not hard to prove that separable states Eq. (2) are local states. From the locality condition Eq. (6), it is possible to derive the so-called Bell inequalities (Bell 1964, Brunner et al. 2014) such that, if a quantum state violates one of those inequalities, then that state is nonlocal. In addition to having introduced this concept, Bell also derived the very first Bell inequality with which he was capable of proving the nonlocality of the two-qubit state  $|\psi\rangle := \frac{1}{\sqrt{2}}(|01\rangle - |10\rangle)$  (Bell 1964). It is well known that nonlocality and entanglement are *equivalent* for multipartite pure states (in the sense that every entangled state is nonlocal and viceversa) (Gisin 1991, Gisin & Peres 1992, Popescu & Rohrlich 1992). However, this is not the case for general mixed states, i. e., there exist mixed entangled local states (Werner 1989, Brunner et al. 2014). For a bipartite system, the aforementioned Bell inequalities can be classified through the parameters  $(m_A, m_B, n_A, n_B)$  where  $x = 1, \dots, m_A$ ,  $y = 1, \dots, m_B$ ,  $a = 1, \dots, n_A$ , and  $b = 1, \dots, n_B$  (Brunner et al. 2014). Here, we shall discuss the simplest non trivial case,  $(2, 2, 2, 2)$ . This

inequality has a very compact form, at least for two-qubit systems, which we detail in what follows.

**CHSH Inequality:** For a bipartite system, the Clauser-Horne-Shimony-Holt (CHSH) inequality (Clauser et al. 1969) considers two dichotomic (eigenvalues  $\pm 1$ ) observables per party, namely,  $(A_1, A_2, B_1, B_2)$ . With the notation already introduced, this would be the inequality (2,2,2,2) and it takes the form:

$$|B_\rho(A_1, A_2, B_1, B_2)| := |E_{11} + E_{12} + E_{21} - E_{22}| \leq 2, \quad (8)$$

where  $E_{ij} := \text{Tr}[(A_i \otimes B_j) \rho]$ ,  $i, j = 1, 2$ . Both, this and more general Bell inequalities can be derived in a systematic way by means of a geometric approach (Brunner et al. 2014). The idea is to maximise the  $B_\rho$  function over those four observables. There is no compact solution for general arbitrary dimensions, except for two qubits (Horodecki et al. 1995). In this case any state  $\rho \in D(\mathbb{C}^2 \otimes \mathbb{C}^2)$ , can be written as:

$$\frac{1}{4} \left( \mathbb{1}_{4 \times 4} + \vec{\sigma} \cdot \vec{a} \otimes \mathbb{1} + \mathbb{1} \otimes \vec{\sigma} \cdot \vec{b} + \sum_{n,m=1}^3 t_{n,m} \sigma_n \otimes \sigma_m \right), \quad (9)$$

with  $\vec{\sigma} = [\sigma_i]$ ,  $\sigma_i$ ,  $i = 1, 2, 3$ , the Pauli matrices,  $\vec{a} = [a_i]$  with  $a_i := \text{Tr}[(\sigma_i \otimes \mathbb{1})\rho]$ ,  $\vec{b} = [b_i]$  with  $b_i := \text{Tr}[(\mathbb{1} \otimes \sigma_i)\rho]$ , and  $t_{nm} := \text{Tr}[\rho(\sigma_n \otimes \sigma_m)]$  making the matrix  $T_\rho := [t_{nm}] \in M_{3 \times 3}(\mathbb{R})$ . We have the following characterisation in terms of the sum  $M(\rho) := \mu + \tilde{\mu}$  of the two biggest eigenvalues  $\mu, \tilde{\mu}$  of the matrix  $U_\rho := T_\rho^T T_\rho \in M_{3 \times 3}(\mathbb{R})$ :

$$\begin{aligned} & \rho \text{ violates} \\ & \text{the CHSH} \iff M(\rho) := \mu + \tilde{\mu} > 1. \end{aligned} \quad (10)$$

Inequality Eq. (8)

This is because, it is possible to show that  $\max B_\rho := |\max_{A_1, A_2, B_1, B_2} B_\rho| = 2\sqrt{M(\rho)}$ . Then, using the Cirelson's bound  $\max B_\rho \leq 2\sqrt{2}$  (Cirelson 1980), it follows  $0 \leq M(\rho) \leq 2$ , showing nonlocality in the interval  $1 < M(\rho) \leq 2$ . Instead of  $M(\rho)$ , we could work with  $B(\rho) := \sqrt{\max \{0, M(\rho) - 1\}}$  because for pure states we have that the former turns out to be equal to the concurrence already discussed, i. e.,  $C(|\psi\rangle) = B(|\psi\rangle)$  (Miranowicz 2004). However, in order to analyse nonlocality through the CHSH inequality and make a direct comparison with EoF, we will plot:

$$CHSH(\rho) = h \left( \frac{1 + \sqrt{1 - B(\rho)^2}}{2} \right), \quad (11)$$

being  $h$  the binary entropy defined in the previous section. It should be pointed out that, if  $\rho$  does not violate the CHSH inequality, it does not necessarily


imply that  $\rho$  is a local state; actually, it is possible that the state violates another inequality, for instance the state considered in [Collins & Gisin \(2004\)](#). On the other hand, proving locality is perhaps trickier than doing so for nonlocality ([Brunner et al. 2014](#)). To wrap up, we search for nonlocality by means of the CHSH inequality, however, there are entangled states that do not violate it and even more, there are entangled local states (states that do not violate any Bell inequality). Let us address the process of how to activate nonlocality on these last classes of states.

### Activation of Nonlocality: Scenarios

On the one hand, we have already pointed out that, although for pure states entanglement and nonlocality are equivalent, (in the sense that every entangled state is nonlocal and vice versa) this is not the case for general mixed states. This, in principle, would settle the question about the relationship between entanglement and nonlocality. However, in the mid nineties, Popescu was able to activate the *hidden nonlocality* of a class of entangled local states after a proper manipulation of them ([Popescu 1995](#)). This raised again the question about the relationship between entanglement and what we could call, new definitions (generalisations) of nonlocality. Therefore, from a fundamental point of view, it is interesting to explore the aforementioned new relationship.

On the other hand, and from a practical point of view, even though entanglement has been regarded as resource for many tasks ([Horodecki et al. 2009](#)), it is also well known that there are protocols that explicitly require either nonlocality or the violation of certain Bell inequalities ([Brunner et al. 2014](#), [Acin et al. 2006a](#)). Let us consider the following situation: suppose that an experimentalist is able to prepare entangled local states only, but he needs to implement protocols that require nonlocality. What could he do about it? These procedures are called *activation scenarios* and we address them in what follows ([Brunner et al. 2014](#)):

**Activation through Local Filtering (LF):** Popescu, in Ref. ([Popescu 1995](#)), took Werner entangled local states  $\rho_W \in D(\mathbb{C}^d \otimes \mathbb{C}^d)$  and after applying a *local filter*, an operation that could be thought as a projection onto a two-qubit system, namely  $P \otimes Q$  with  $P := |1\rangle\langle 1| + |2\rangle\langle 2|$ , and  $Q := |1\rangle\langle 1| + |2\rangle\langle 2|$ , he found that the final state violated the CHSH inequality Eq. (8), with  $d > 5$  (See Appendix A for a detailed explanation). We can see a sketch of the procedure in **Figure 2(a)**. Soon after this result, Gisin ([Gisin 1996](#)) proposed another example, but this time with states that, even though it was not known whether they were local, at least, they did not violate the CHSH inequality. Recently, [Pal & Ghosh \(2015\)](#) derived a complete characterisation for two-qubit systems.



**Fig. 2.** (a) Activation of nonlocality through Local Filtering (Popescu 1995). Experimentalists Alice and Bob receive a part of a bipartite quantum state  $\rho$ . They make a local filtering (map onto a two-qubit system) before a standard Bell test. (b) Activation of nonlocality through Tensoring (Navascués & Vértesi 2011). Experimentalists Alice and Bob receive a part of a quantum state  $\rho_1$  and part of another quantum state  $\rho_2$ , then, they make a standard Bell test. (c) Activation of nonlocality through the combination of Tensoring and Local Filtering.

**Activation through Tensoring (T):** Here, the question is: Is it possible to find  $\rho_1$  and  $\rho_2$  entangled local states such that  $\rho_1 \otimes \rho_2$  is an entangled nonlocal state? This procedure is illustrated in **Figure 2(b)**. If  $\rho_2 = \rho_1$ , or in general if  $\rho \otimes^k$  is nonlocal, the phenomenon is called *superactivation*. Similarly, we could also say that the state  $\rho$  is *k-copy nonlocal* or that we have activation through *k-tensoring*. It is also natural to think about the combination of this procedure with the previous one as we can see in **Figure 2(c)**. Historically, the first tensoring procedure appeared as a combination of *k*-tensoring with LF (Peres 1996a, Masanes 2006). Then, general tensoring and LF would arise (Masanes et al. 2008, Liang et al. 2012). The protocol of activation through tensoring alone (without LF) was reported in Navascués & Vértesi (2011), later, superactivation was put forward (Palazuelos 2012, Cavalcanti et al. 2013).

We next address the following three scenarios. First, hidden nonlocality (Pal & Ghosh 2015). Second, *k*-copy nonlocality (Palazuelos 2012, Cavalcanti et al. 2013). Third, tensoring with local filtering (first tensoring, then local filtering) (Masanes et al. 2008, Liang et al. 2012). We shall focus on numerical approaches for two-qubit systems.

## Local Filtering: Hidden Nonlocality (HN)

Recently, [Pal & Ghosh \(2015\)](#) derived a complete characterisation of hidden nonlocality for two-qubit systems.

Given  $\rho \in D(\mathbb{C}^2 \otimes \mathbb{C}^2)$ ,  $\rho$  possesses hidden nonlocality (or  $\rho$  can be transformed into a CHSH-violating state  $\rho'$  by means of local filters) if and only if  $\lambda_\rho^1 + \lambda_\rho^2 > \lambda_\rho^0$ , where the  $\lambda_\rho^i$ 's are the eigenvalues of the matrix  $C_\rho := \eta T_\rho \eta T_\rho^T$  organized in decreasing order, with  $\eta := \text{diag}(1, -1, -1, -1)$  and  $T_\rho := [t_{nm}]$  a matrix with elements  $t_{nm} := \text{Tr}[(\sigma_m \otimes \sigma_n)\rho]$  as defined in the CHSH nonlocality section. The CHSH-violation of that new state  $\rho'$  is given by  $M'(\rho) := M(\rho') = \frac{\lambda_\rho^1 + \lambda_\rho^2}{\lambda_\rho^0}$ . Instead of  $M'(\rho)$ , we could work with  $B'(\rho) := \sqrt{\max\{0, M'(\rho) - 1\}}$  because for pure states we have that the former turns out to be equal to the concurrence already discussed, i. e.,  $C(|\psi\rangle) = B(|\psi\rangle)$  ([Miranowicz 2004](#)). Then, in order to analyse hidden nonlocality, we will plot:

$$HN(\rho) = h \left( \frac{1 + \sqrt{1 - B'(\rho)^2}}{2} \right), \quad (12)$$

being  $h$  the binary entropy defined in the EoF and CHSH sections.

## Tensoring: $k$ -copy Nonlocality (Superactivation of Nonlocality SA)

In this section we focus on the case of the activation of nonlocality only through tensor product; in particular, when the state is capable of activating nonlocality by itself, i. e., superactivating ([Palazuelos 2012](#), [Cavalcanti et al. 2013](#)). We shall specialise our results on two-qubit systems.

**Main Theorems:** In order to talk about  $k$ -copy nonlocality, the teleportation protocol ([Bennett et al. 1993](#)) will come in handy. Given a two-qubit state  $\rho \in D(\mathbb{C}^2 \otimes \mathbb{C}^2)$ , we have the next hierarchy of properties depicted by the following chain of implications:

$$\begin{aligned} & \rho \text{ violates the CHSH inequality Eq. (8).} \\ \Downarrow & \text{ (Horodecki et al. 1996b)} \\ & \rho \text{ is useful for teleportation.} \\ \Downarrow & \text{ (Palazuelos 2012, Cavalcanti et al. 2013)} \\ & \rho \text{ is } k\text{-copy nonlocal.} \end{aligned} \quad (13)$$

The proofs of both implications can be found in [Horodecki et al. \(1996b\)](#), [Palazuelos \(2012\)](#), and [Cavalcanti et al. \(2013\)](#), respectively. Here, we comment on a couple of remarks: The first implication is restrictive in the sense

that there exist entangled local states (states that do not violate any Bell inequality, in particular, the CHSH inequality), although useful for teleportation (Popescu 1994). The second implication actually holds true even for general two-qudit systems  $\rho \in D(\mathbb{C}^d \otimes \mathbb{C}^d)$  (Palazuelos 2012, Cavalcanti et al. 2013). Third, usefulness for teleportation does not cover all the entangled states set, i. e., there exist entangled states not useful for teleportation (Horodecki et al. 1996b). Fourth, it remains open both, the existence of an entangled never  $k$ -copy nonlocal state and a  $k$ -copy nonlocal, not useful for teleportation state. Finally, since usefulness for teleportation is a property which is possible to search for numerically, it consequently allows us to enquire about  $k$ -copy nonlocality. In what follows, we discuss how to numerically calculate usefulness for teleportation.

The teleportation protocol (Bennett et al. 1993), in a nutshell, works as follows. A quantum pure state  $|\phi\rangle \in \mathbb{C}^d$  can be teleported by means of a channel built with another quantum state  $\rho \in D(\mathbb{C}^d \otimes \mathbb{C}^d)$ . In order to check how useful  $\rho$  is to the protocol, a function called *Fidelity of Teleportation* (FoT) was proposed (Popescu 1994). A seminal result regarding this function (Horodecki et al. 1999) establishes that:

$$F_d(\rho) = \frac{df_d(\rho) + 1}{d + 1}, \quad (14)$$

where  $f_d$  is the so called *Fully Entangled Fraction* (FEF) (sometimes also called *Entanglement Singlet Fraction*) given by:

$$f_d(\rho) := \max_{\psi \in ME} \langle \psi | \rho | \psi \rangle, \quad (15)$$

where  $ME$  stands for the set of maximally entangled states, i. e., the states  $:= |\psi\rangle \langle \psi|$ , defined as:  $|\psi\rangle := (U_1 \otimes U_2) \frac{1}{\sqrt{d}} \sum_{i=0}^{d-1} |ii\rangle$ , with  $U_1, U_2$  local unitary operations. Taking a look at Eq. (15), we have  $\frac{1}{d+1} < F_d < 1$ . If we use separable states in the FEF Eq. (15), we obtain the value  $f_d(\rho_{sep}) = \frac{1}{d}$  (Horodecki et al. 1999). Consequently, we have the characterisation:

$$\rho \text{ is useful for teleportation} \iff f_d(\rho) > \frac{1}{d}. \quad (16)$$

In particular, for a two-qubit state  $\rho$  ( $d = 2$ ), with  $\frac{1}{3} < F_d(\rho) < 1$ ,  $\rho$  is useful for teleportation if and only if  $f_2(\rho) > \frac{1}{2}$  or  $F_2(\rho) > \frac{2}{3}$ . Therefore, we can search for  $k$ -copy nonlocality through the usefulness for teleportation by means of the calculation of the FoT Eq. (14). A natural question that arises is: what is the actual  $k$  value necessary for this superactivation? It turns out that, from the second implication in Eq. (13), it is possible to extract this value (Palazuelos 2012, Cavalcanti et al. 2013). This  $k$  value is a minimum because it is necessary only one nonlocal state in order to activate nonlocality through tensoring (Popescu & Rohrlich 1992). We next show how to calculate this  $k$ .

**The superactivation tensoring factor  $k(d, f_d)$ :** By means of the above-mentioned theorem, the second implication in Eq. (13) it is possible to extract the explicit values of  $k$  in terms of the dimension  $d$  of the two-qudit system and their fully entangled fraction  $f_d$ . From the theorem, those  $k$ 's must satisfy the following relation:

$$\left[ \frac{C'}{C(\ln d)^2} \right] \frac{(f_d d)^k}{k^2} > 1, \quad (17)$$

with constants  $C := e^4$ ,  $C' := 4$ . Since in Eq. (17), we have a function to the power of  $k$  over a polynomial (quadratic) function of  $k$ , we see that the activation goes from  $f > 1/d$ . Numerically, in **Figure 3(a)**, we see the behaviour of  $k(d, f_d)$ . For most of the  $d - f$  region,  $k < 10$  is enough (as it is depicted by the white curve in the left); just when  $f_d$  is close to the boundary  $\frac{1}{d}$ , it becomes asymptotically more difficult to superactivate it. In **Figure 3(b)**, we see cuts for  $d = 2, 3, 4$  and  $5$ . Next, we will analyse how to calculate  $f_d$  for two qubits ( $d = 2$ ) (Grondalski et al. 2002).

**Fidelity of Teleportation (FoT):** Following (Grondalski et al. 2002), we summarise the Fidelity of Teleportation (FoT) characterisation for two-qubit systems. Given  $\rho \in D(\mathbb{C}^2 \otimes \mathbb{C}^2)$ , we have that  $f_2(\rho) = \max\{\eta_i, 0\}$ , where the  $\eta_i$ 's are the eigenvalues of the matrix  $M = [M_{mn}]$ , with elements  $M_{mn} = \text{Re}(\langle \psi_m | \rho | \psi_n \rangle)$ , and  $\{|\psi_n\rangle\}$  the so-called magic basis  $|\psi_{ab}\rangle := i^{(a+b)}(|0, b\rangle + (-1)^a |1, 1 \oplus b\rangle)/\sqrt{2}$ . The FoT is given by  $F_2(\rho) = \frac{2f_2(\rho)+1}{3}$ , and we have already seen that  $\rho$  is useful for teleportation if and only if  $f_2 > \frac{1}{2}$  or  $F_2(\rho) > \frac{2}{3}$ . It should be pointed out that, the bound  $f_2 = \frac{1}{2}$  is also a measure of usefulness for other two-qubit protocols (Grondalski et al. 2002). In order to enquire about this property, we shall plot:

$$SA(\rho) = F_2'^+, \quad \text{with} \quad F_2' := F_2 - \frac{2}{3}, \quad (18)$$


and  $^+$  the positive part of the function. For multipartite or high-dimensional systems, there is not a general explicit relation for the FEF, except for Werner and Isotropic states (Zhao et al. 2010).

### Tensoring and Local Filtering (T&LF)

In this Section, we deal with the activation of nonlocality through tensoring and local filtering (Masanes et al. 2008, Liang et al. 2012). Let us consider the set  $P_{\text{CHSH}}$  formed by states satisfying the CHSH inequality even after all possible local filtering (LF) operations (Masanes et al. 2008). In other words:  $P_{\text{CHSH}} \subset D(\mathbb{H})$ ,

$$\rho \in P_{\text{CHSH}} \iff \forall \Omega : D(\mathbb{H}) \longrightarrow D(\mathbb{C}^2 \otimes \mathbb{C}^2) \text{ LF} \quad (19)$$

it holds that,  $\Omega(\rho)$  does not violate CHSH.



**Fig. 3.** (a) Superactivation tensoring factor  $k(d, f_d)$  in terms of the fully entangled fraction (FEF)  $0 \leq f_d \leq 1$  and the dimension  $d \geq 2$  of the two-qudit system. The white curves represent the threshold for  $k < 10$  and  $k < 3$  from left to right respectively. (b) Transversal cuts of Fig. 3. (a) for  $d = 2, 3, 4$  and  $5$ .

Here, the local filter LF denotes a *separable map* of the form  $\Omega(\rho) = \sum_i (A_i \otimes B_i)\rho(A_i \otimes B_i)^\dagger$ , with  $A_i, B_i$  being Kraus operators (Masanes et al. 2008). This  $P_{\text{CHSH}}$  set has been characterised for two-qubit systems (Verstraete & Wolf 2002). If we define  $P'_{\text{CHSH}}$  as the set of states that do not violate CHSH even after  $k$  tensoring themselves or LF, also called *not asymptotically violation* (Masanes 2006), we have the relation  $P'_{\text{CHSH}} \subset P_{\text{CHSH}}$ . There is an equivalence between this asymptotically violation of CHSH (Masanes 2006) and another property called *distillability* (Bennett et al. 1996a), which will let us search numerically for states in  $P_{\text{CHSH}}$ .

**Main Theorem:** Since we are focused on two-qubit systems, we highlight the result for bipartite systems (Masanes et al. 2008). However, it should be said that, the theorem also works for general multipartite systems (Liang et al. 2012). It establishes the following:

$$\begin{array}{ccc}
\tau \in D(\mathbb{C}^{d_A} \otimes \mathbb{C}^{d_B}) & \exists \rho_\tau \in D \left[ \bigotimes_{i=1}^2 (\mathbb{C}^{d_i} \otimes \mathbb{C}^2) \right] \\
\text{is} & \iff & \text{s.t. } \rho_\tau \in P_{\text{CHSH}}, \\
\text{entangled} & & \tau \otimes \rho_\tau \notin P_{\text{CHSH}}.
\end{array} \tag{20}$$

A couple of remarks: First, since the result holds for any entangled state  $\tau$ , it also applies for entangled states  $\tau \in P_{\text{CHSH}}$ , which would imply that the CHSH nonlocality of  $\tau$  has been activated (in the sense of  $P_{\text{CHSH}}$ ) through tensoring (with  $\rho_\tau$ ) and LF. We could also call it, tensorial activation of hidden nonlocality. Second, even though the theorem guarantees the existence of the matrix  $\rho_\tau$ , it does not tell us explicitly how to calculate it.

**Numerical Approach:** Given  $\tau \in D(\mathbb{C}^{d_A} \otimes \mathbb{C}^{d_B})$  an entangled state, we would like to find the aforementioned respective density matrix  $\rho_\tau$ . In order to do so, we follow the approach reported in (Liang et al. 2012). From the proof of the main theorem, which can be found in (Liang et al. 2012), it is enough to look for a density matrix  $\rho_\tau \in D[(\mathbb{C}^{d_A} \otimes \mathbb{C}^2) \otimes (\mathbb{C}^{d_B} \otimes \mathbb{C}^2)]$  with the following characteristics. First,

$$\text{Tr} [\rho_\tau (\tau^T \otimes H_{\pi/4})] < 0, \tag{21}$$

with:  $H_\theta := \mathbb{1} \otimes \mathbb{1} - \cos \theta \sigma_x \otimes \sigma_x - \sin \theta \sigma_z \otimes \sigma_z$ .

Second, we have to check that  $\rho_\tau \in P_{\text{CHSH}}$ . In principle, it is not numerically possible to check if a state belongs or not to the set  $P_{\text{CHSH}}$  (with the exception of two-qubit systems (Pal & Ghosh 2015), however, our matrix  $\rho_\tau$  is not of this sort). Fortunately, we have some results that partially allow us to search for it: i) as we have already pointed out,  $P'_{\text{CHSH}} \subset P_{\text{CHSH}}$ ; ii)  $P'_{\text{CHSH}}$  is equivalent to *Bound Entanglement (BE)* (Masanes 2006); iii) there is a class of these bound entangled states, the so-called *PPT* (Positive Partial Transpose) states (Peres 1996b, Horodecki et al. 1996a). The former are states such that, their partial transpose respect to the first subsystem is positive or  $\rho^{T_1} \geq 0$ . In other words, given  $\rho \in D(\mathbb{C}^{d_A} \otimes \mathbb{C}^{d_B})$ , which is always possible to write it as  $\rho := \sum_{ijkl} \rho_{ijkl} |i\rangle\langle j| \otimes |k\rangle\langle l|$  with  $\rho_{ijkl}$  coefficients, the partial transpose respect to the first subsystem is defined as:

$$\begin{aligned}
\rho^{T_1} &:= \sum_{ijkl} \rho_{ijkl} (|i\rangle\langle j|)^T \otimes |k\rangle\langle l| \\
&= \sum_{ijkl} \rho_{ijkl} |j\rangle\langle i| \otimes |k\rangle\langle l|,
\end{aligned} \tag{22}$$

with  $T$  the standard transposition. The state  $\rho$  is then a *PPT* state, if its partial transpose remains positive or  $\rho^{T_1} \geq 0$ . Therefore, we have the hierarchy of properties:

$$PPT \longrightarrow BE \longleftrightarrow P'_{\text{CHSH}} \longrightarrow P_{\text{CHSH}}. \quad (23)$$

Hence, we could look for a matrix  $\rho_\tau$  with a positive partial transposition respect to the first subsystem  $\mathbb{C}^{d_A} \otimes \mathbb{C}^2$  in order to guarantee that it belongs to  $P_{\text{CHSH}}$ . Then, we have the following minimisation problem:

$$\begin{aligned} \text{minimise: } \quad & \sigma(\tau) := \text{Tr} [\rho_\tau (\tau^T \otimes H_{\pi/4})], \\ \text{over } \{ \rho_\tau \} \\ \text{and constraints: } \quad & \rho_\tau \geq 0, \quad \rho_\tau^{T_1} \geq 0. \end{aligned} \quad (24)$$


A problem with these characteristics is a Semidefinite Programming (SDP) problem which is numerically solvable (Liang et al. 2012, Vandenberghe & Boyd 1996). We have solved it by using MATLAB (MATLAB 2011b) with the YALMIP toolbox (Löfberg 2004) and the solvers SDPT3 (Toh et al. 1998) and SeDuMi (Sturm 1999). We next analyse all these properties on specialised examples.

## Results: Some Quantum Properties of States of Interest

In this section, we shall use the formalism and tools already described in the previous sections in order to analyse the following quantum properties of some specific states: entanglement, nonlocality,  $k$ -copy nonlocality, hidden nonlocality, activation T&LF, and locality. From now on, we will deal with quantum states in terms of a parameter  $p$ , i. e.,  $\rho = \rho(p)$ , with the following notation: if  $p > p_E$  then the state is entangled, if  $p > p_{NL}$  the state is nonlocal, if  $p > p_{SA}$  the state is useful for teleportation (and therefore  $k$ -copy nonlocal), if  $p > p_{HN}$  the state contains hidden nonlocality, if  $p > p_{T&LF}$  the program described in the previous section has found an ancillary state that helps to the activation T&LF, and if  $p \leq p_L$  the state is local. We will analyse these properties onto the Werner states reproducing the values reported in Liang et al. (2012). Additionally, we report new activation regions for Isotropic and Hirsch states.

**Werner-Isotropic (WI) States:** We first consider the so-called Werner states (Werner 1989). Particularly, the two-qubit version of them, in which we refer to them as the Werner-Isotropic (WI) states, and read

$$\begin{aligned} \tau_{WI}(p) &= p |\psi_-\rangle \langle \psi_-| + \frac{(1-p)}{4} \mathbb{1}, \\ 0 \leq p \leq 1, \quad |\psi_-\rangle &:= \frac{1}{\sqrt{2}} (|01\rangle - |10\rangle). \end{aligned} \quad (25)$$



**Fig. 4.** Quantum properties for the WI states Eq. (25). Entanglement-EoF (blue solid thick curve) Eq. (4). Superactivation-SA (green dotted curve) Eq. (18). Nonlocality-CHSH (red dashed-dotted curve) Eq. (11). Hidden Nonlocality - HN (magenta solid thin curve) Eq. (12). Activation T&LF through the minimisation procedure Eq. (24) (cyan dashed curve) for which we have plotted  $-5\sigma[\tau_{WI}(p)]$  in order to make it visible amongst the other properties. R (orange triply dashed curve), activation T&LF with the ancillary matrix Eq. (26), which was obtained by the minimisation procedure Eq. (24) at  $p_A = 0.6569$ , and Projective-Locality (black vertical solid line) at  $p_L = 0.6595$  according to the best known bound derived in Ref. (Acín et al. 2006b).

In **Figure 4**, we have plotted the nonlocality-related properties discussed throughout the paper for these WI states, namely, EoF, CHSH, SA, HN and activation T&LF. Regarding activation T&LF, we have plotted  $-5\sigma[\tau_{WI}(p)]$  in order to make it visible amongst the other measures. We were also able to extract the ancillary matrix for  $p = 0.6569$  Eq. (26), which turns out to be useful for all the  $0.6569 < p < 1$  region, as we show in the  $R(p)$  function plotted in Figure 4. Therefore,  $\tau_{WI}(p) \otimes \rho_\tau$  is CHSH-nonlocal after a LF with,

$$\rho_\tau = \frac{1}{16} \sum_{i,j=0}^3 R_{ij} \sigma_i \otimes \sigma_i \otimes \sigma_j \otimes \sigma_j, \quad (26)$$

where

$$R := \frac{1}{9} \begin{pmatrix} 9 & 3 & 3 & 3 \\ 1 & -1 & 3 & -1 \\ 1 & -1 & 3 & -1 \\ 1 & -1 & 3 & -1 \end{pmatrix}. \quad (27)$$

To complete the results reported in Figure 4, the following **Table 1** gives the limit bounds of the nonlocality-related properties beyond entanglement for the WI states.

$p_E$	$p_{SA}$	$p_{T&LF}$	$p_L$	$p_{HN}$	$p_{NL}$
0.3333	0.3333	0.6569	0.6595	0.7054	0.7054

**Table 1.** Limit values for the nonlocality-related properties beyond entanglement for WI States Eq. (25).

A few things should be pointed out: First, even though the nonlocality limit point obtained from the CHSH inequality is  $p_{NL} = \frac{1}{\sqrt{2}} \approx 0.7071$ , there exists a slightly better bound which reads  $p_{NL} = 0.7054$  (Hua et al. 2015). Second, CHSH-nonlocality coincides with hidden nonlocality because the WI states are already in a Bell-diagonal form (see (Verstraete & Wolf 2002) for a detailed explanation). We see that for the states in the entangled local region  $0.3333 < p < 0.6596$ , even though they do not present hidden nonlocality, they present superactivation of nonlocality and Activation T&LF.

**Werner States:** We next address the so-called two-qudit Werner states (Werner 1989):

$$\begin{aligned} \tau_W^d(p) &= \frac{p}{d(d-1)} 2P_{\text{anti}} + \frac{(1-p)}{d^2} \mathbb{1}, \\ 1 - \frac{2d}{d+1} &\leq p \leq 1, \end{aligned} \tag{28}$$

$$P_{\text{anti}} := \frac{1}{2} \left( \mathbb{1} - \sum_{ij}^d |i\rangle \langle j| \otimes |j\rangle \langle i| \right).$$

In **Figure 5**, we have plotted the activation T&LF by means of the minimisation of the  $\sigma(\tau_W^d)$  function, according to Eq. (24) and up to qudits of dimension  $d = 6$ . In **Table 2**, we report the limit values for the nonlocality-related properties discussed throughout the paper, also up to  $d = 6$ . The entanglement and locality limits come from Werner (1989). The superactivation limit by means of the FoT comes from Zhao et al. (2010), from which it turns out they (as soon as  $d > 2$ ) are not useful for teleportation. Therefore, it is unknown if they are superactivable or not, and so, we have marked them in the table with an X. The activation T&LF column follows the plots shown in the Figure 5 and also reported in Liang et al. (2012). Finally, in the last column, hidden nonlocality from Popescu (1995). In Appendix A, we give a more detailed description of these bounds.


$d$	$p_E$	$p_{SA}$	$p_{T\&LF}$	$p_L$	$p_{HN}$
2	0.3333	0.3333	0.6569	0.6595	0.7054
3	0.2500	X	0.6360	0.6667	0.7630
4	0.2000	X	0.6247	0.7500	0.7837
5	0.1429	X	0.6174	0.8000	0.7944
6	0.1667	X	0.6127	0.8333	0.8009

**Table 2.** Limit values for nonlocality-related properties beyond entanglement for Werner states Eq.(28) up to  $d = 6$ .

**Isotropic States:** We now consider the Isotropic states (Horodecki & Horodecki 1999):

$$\tau_I^d(p) = p |\psi_d\rangle \langle \psi_d| + \frac{(1-p)}{d^2} \mathbb{1},$$

$$0 \leq p \leq 1, \quad |\psi_d\rangle := \frac{1}{\sqrt{d}} \sum_{i=1}^d |ii\rangle. \quad (29)$$



**Fig. 5.** Activation T&NL for Werner states. Minimisation procedure Eq. (24) of the function  $\sigma[\tau_w^d(p)]$  vs parameter  $p$  for  $\tau_w^d(p)$ , high dimensional Werner states Eq. (28).  $d = 2$  (green dotted curve),  $d = 3$  (red dashed-dotted curve),  $d = 4$  (cyan triply dashed curve),  $d = 5$  (magenta doubly dashed curve),  $d = 6$  (yellow dashed curve). The inset shows a zoom of the functions close to the zero value. We reproduce the values reported in Liang et al. (2012), which we report in Table 2.

In **Figure 6**, we have plotted the activation T&LF by means of the minimisation of the  $\sigma(\tau_I^d)$  function, according to Eq. (24) and up to qudits of dimension  $d = 6$ . In **Table 3**, we report the limit values for the nonlocality-related properties discussed throughout the paper up to  $d = 6$ . The entanglement and locality limits are taken from Horodecki & Horodecki (1999). The superactivation limit is obtained from Zhao et al. (2010) (all of them are useful). Activation T&LF comes from plots shown in Figure 6. Finally, nonlocality limit values are obtained from the Collins-Gisin-Linden-Massar-Popescu (CGLMP) inequalities (Collins et al. 2002). In Appendix B, we give a more detailed description of these bounds.


**Hirsch States:** Finally, we analyse the two-qubit states studied by Hirsch, Quintino, Bowles and Brunner (Hirsch et al. 2013) which for simplicity we will call Hirsch states:

$$\begin{aligned} \tau_F(p, q, \sigma) = & p |\psi_-\rangle \langle \psi_-| \\ & + [1 - p] \left[ q\sigma + (1 - q)\frac{1}{2} \right] \otimes \frac{1}{2}, \\ 0 \leq p \leq 1, \quad 0 \leq q \leq 1, \quad |\psi\rangle := & \frac{1}{\sqrt{2}}(|01\rangle - |10\rangle), \end{aligned} \quad (30)$$

and  $\sigma$  an arbitrary one-qubit state. These states can also be thought as a generalisation of the two-qubit WI states Eq. (25); in fact, we recover them by putting  $\sigma = \frac{1}{2}$ , and  $q = 1$ . In **Figure 7(a)**, we have plotted the nonlocality-related properties for the two-parameter Hirsch states Eq. (30), using  $\sigma = |0\rangle \langle 0|$ .

$d$	$p_E$	$p_{SA}$	$p_{T&LF}$	$p_L$	$p_{HN}$
2	0.3333	0.3333	0.6569	0.6595	0.7054
3	0.2500	0.2500	0.5606	0.4167	0.6961
4	0.2000	0.2000	0.4890	0.3611	0.6905
5	0.1429	0.1429	0.4337	0.3208	0.6872
6	0.1667	0.1667	0.3895	0.2900	0.6849

**Table 3.** Limit values for nonlocality related properties beyond entanglement for Isotropic states Eq.(29) up to  $d = 6$ .




**Fig. 6.** Activation T&NL for Isotropic states. Minimisation procedure Eq. (24) of the function  $\sigma[\tau_I^d(p)]$  vs parameter  $p$  for  $\tau_I^d(p)$  high dimensional Isotropic states Eq. (29).  $d = 2$  (green dotted curve),  $d = 3$  (red dashed-dotted curve),  $d = 4$  (cyan triply dashed curve),  $d = 5$  (magenta doubly dashed curve),  $d = 6$  (yellow dashed curve). We report in **Table 3** the values we are interested in.

This plot should be understood as the projection of these properties upon the  $p - q$  plane. The white region is composed by separable states whilst the rest stands for entangled states and the nonlocality properties that lie within. These properties have been superposed following the hierarchy we expect. Interestingly, these states for  $\sigma = |0\rangle\langle 0|$ , and  $q = 1$  become

$$\tau_F(p) = p |\psi_-\rangle\langle\psi_-| + (1 - p) |0\rangle\langle 0| \otimes \frac{1}{2}, \quad (31)$$

for which the authors of Ref. (Hirsch et al. 2013) were able to prove locality, i. e., to build a local model, for all  $p > 1/2$ . In **Figure 7(b)**, we have plotted the aforementioned properties for the states Eq. (31). In **Table 4**, we have reported the limit values for the aforementioned properties for the states Eq. (31).



**Fig. 7. (a)** Quantum properties for the two-parameter Hirsch states Eq. (30) with  $\sigma = |0\rangle\langle 0|$ . Entanglement-EoF (blue) Eq. (4). Nonlocality-CHSH (red) Eq. (11). Superactivation-SA (green) Eq. (18). Hidden Nonlocality-HN (magenta) Eq. (12). Activation T&LF (cyan) Eq. (24). **(b)** Quantum properties for one-parameter Hirsch states Eq. (31) (or two-parameter Hirsch states with  $\sigma = |0\rangle\langle 0|$  and  $q = 1$ , previous plot (a) at  $q = 1$ ). Entanglement-EoF (blue solid thick curve) Eq. (4). Nonlocality-CHSH (red dashed-dotted curve) Eq. (11). Superactivation-SA (green dotted curve) Eq. (18). Hidden Nonlocality-HN (magenta solid thin curve) Eq. (12). Activation T&LF for which we have plotted  $-5\sigma[\tau_F(p, q)]$  in order to make it visible amongst the other properties (cyan dashed curve) Eq. (24). Projective-Locality (black vertical solid line).

## Discussion

In this work, we have dealt with 6 properties of quantum states, namely, entanglement, nonlocality, locality, and three generalisations of nonlocality by means of the activation scenarios: hidden nonlocality HN (or activation through local filtering),  $k$ -copy nonlocality (or superactivation of nonlocality SA), and activation through tensoring and local filtering (T&LF). We stress that there are more general setups, for instance, tensoring between different states or quantum networks. However, we have chosen the already discussed ones, because, at least for two-qubit systems, we can numerically enquire for these properties (with the exception of locality).

$p_E$	$p_{SA}$	$p_{T&LF}$	$p_L$	$p_{HN}$	$p_{NL}$
0	0	0.1716	0.333	0.5000	0.7071

**Table 4.** Limit values for the nonlocality-related properties beyond entanglement for the one-parameter Hirsch states Eq. (31).

The three activation scenarios we have worked with share the mechanisms of whether local filtering or tensoring. Whilst local filters for two qubits, which could be seen as operations that the experimentalists can locally implement, keep the dimension of the system, tensoring increases it. Additionally, both mechanisms take separable states into separable states (and consequently local states). Therefore, they cannot possibly activate nonlocality on separable states. Thus, they could only be useful for entangled local states.

We then have analysed these properties upon particular states of interest, namely, the so-called Werner, Isotropic, and Hirsch states. These particular choices were made because there exist local bounds for these states which let us consider entangled local states. We remark that unlike the others properties, locality cannot be yet approached numerically. Locality can only be investigated by means of the construction of local models which is not an easy task.

This paper sheds light into mainly two aspects. First, from a practical point of view, it could be seen as a reference guide to calculate nonlocality-related properties for two-qubit systems. In particular, regarding  $k$ -copy nonlocality, Figure 3(a,b) specifies the integer number  $k$  necessary for the superactivation. We have checked these properties upon the well known Werner states Figures (4, 5) and Tables (1,2). Second, we have reported new bounds for other states of interest (namely, Isotropic and Hirsch states). We have chosen these particular states, in the same vein as Werner states, because of their known bounds regarding locality. Even though nonlocality-related properties have already been reported for these states, activation T&LF has not been calculated yet. We have reported these bounds, filling this gap. Before going into the details of our findings, a note on the activation T&LF.

The main theorem regarding activation T&LF (20) guarantees that all entangled states can be activated in this way. Therefore, from a purely theoretical point of view, it could be seen as pointless to further study this scenario in relation with entanglement. However, the theorem in question does not provide us either the matrix or the filter necessary for the activation, which is

an important issue from a practical point of view. The numerical approach Eq (21) could give us the ancillary matrix in question, and it is open how to find the local filter that maximises the activation. We now proceed to discuss our findings for the Isotropic and Hirsch states.

For the so-called two-qudit Isotropic states, even though nonlocality-related properties have been studied, no bounds for activation T&LF have been reported yet. In Figure 6, we have calculated these bounds which we report in Table 3 column 4 ( $p_{T\&LF}$ ) in terms of the dimension  $d$  of the qudits. From these results, one can observe the following. Unlike the Werner states' bounds, these Isotropic states' bounds do not cover the local states (given  $d > 2$ , values in Table 3 column 4 ( $p_{T\&LF}$ ) are still greater than values in Table 3 column 5 ( $p_L$ )). However, these new bounds now extend the known nonlocality region (given  $d > 2$ , values in Table 3 column 6 ( $p_{NL}$ ) are greater than values in Table 3 column 4 ( $p_{T\&LF}$ )). Unfortunately, there is no two-qudit characterisation of the set  $P_{CHSH}$ , unlike two-qubit systems (Pal & Ghosh 2015), so we cannot say anything in this regard.

For the so-called two-qubit Hirsch states, even though nonlocality-related properties have been reported, neither bounds for hidden nonlocality nor activation T&LF have been reported yet. In Figure 7(a), we have calculated these bounds. First, we remark the hierarchy amongst these properties. All of these properties are inside entanglement and all of them cover the standard definition of nonlocality. Second, we are able to report a  $P_{CHSH}$  activation region for these qubits (which could also be called tensorial activation of hidden nonlocality), here depicted by the cyan region (activation T&LF) that is not covered by the magenta region (HN) in Figure 7(a).

We have depicted the particular case of Figure 7(a) for  $q = 1$  in Figure 7(b) and Table 4, because there is a locality bound for these states. First, the states within the locality region ( $p < p_L = 0.5$ ) are usually considered useless for quantum protocols based on nonlocality. However, they are now displaying a nice variety of generalised nonlocality-related properties. Actually, all of the three generalisations we are considering here, unlike Werner states in Figure 4 which present SA (all of the local region) and Act T&LF (a small region) but no HN. Second, comparing this again with the Werner states in Figure 4, we are numerically showing that there is not a trivial relation between these generalisations of nonlocality. In particular, there are Werner states with SA but no HN whilst there are Hirsch states with HN but no SA.

## Concluding Remarks

First, we have reviewed the so far known activation of nonlocality scenarios. We payed particular attention to the hidden nonlocality,  $k$ -copy nonlocality, and the activation through tensoring and local filtering, in particular, upon two-qubit systems. We have reviewed the numerical approaches required in order to establish a quantification of such quantum properties. For the particular case of two-qudit Werner states, we have reproduced the limit points of the above-mentioned properties.

Second, using the above tools we analysed the activation of nonlocality related properties now for Isotropic and Hirsch states. In particular, we reported limit points on the activation of nonlocality through tensoring and local filtering that, to the best of our knowledge, have not been reported so far. Additionally, due to the recent result in Pal & Ghosh (2015), we have calculated hidden nonlocality for two-qubit Hirsch states which has led us to report tensorial activation of hidden nonlocality.

## Acknowledgements

We gratefully acknowledge support from COLCIENCIAS Program “Jovenes Investigadores e Innovadores Virginia Gutierrez de Pineda” under contract No. RC-0618-2013, from COLCIENCIAS contract No. 71003, Universidad del Valle-internal project No. 7930, the Colombian Science, Technology and Innovation Fund-General Royalties System (Fondo CTeI-SGR) contract No. BPIN 2013000100007, and from CIBioFi.

## Conflict of interest

There are no conflicts of interest with funding sources or institutions.

## Appendix A: Werner States

Here we show the nonlocality-related properties limit points (except activation T&LF) for the two-qudit Werner states Eq. (28). They are entangled for  $p > p_E$  and local for  $p \leq p_L$  with (Werner 1989):

$$p_E = \frac{1}{d+1}, \quad p_L = \frac{d-1}{d}. \quad (32)$$

The explicit calculation of the fidelity of teleportation reported in Zhao et al. (2010), shows that all entangled Werner states are not useful for teleportation, i. e.,  $f_d(\rho_{\text{ent}}) < \frac{1}{d}$ . Their hidden nonlocality or nonlocality through LF (only) is checked as in Popescu (1995), which we detail in what follows. Applying upon the two-qudit Werner states Eq. (28) the local filtering operation given by  $P \otimes Q$ , with:

$$P = |0\rangle\langle 0|_A + |1\rangle\langle 1|_A, \quad Q = |0\rangle\langle 0|_B + |1\rangle\langle 1|_B, \quad (33)$$

we obtain the two-qubit state  $\rho(p) := [P \otimes Q]\rho_W^d(p)$ ,

$$\rho(p) = \frac{p}{d(d-1)} 2|\psi\rangle\langle\psi| + \frac{(1-p)}{d^2} \mathbb{1}_2 \otimes \mathbb{1}_2, \quad (34)$$

with  $|\psi\rangle := \frac{1}{\sqrt{2}}(|01\rangle - |10\rangle)$ . Normalizing we obtain:

$$\begin{aligned} \rho(p) &= \left[ \frac{d(d-1)d^2}{2pd^2 + (1-p)4d(d-1)} \right] \\ &\quad \times \left[ \frac{p}{d(d-1)} 2|\psi\rangle\langle\psi| + \frac{(1-p)}{d^2} \mathbb{1}_2 \otimes \mathbb{1}_2 \right]. \end{aligned}$$

Now, checking its CHSH maximal violation by means of the criterion Eq. (10), we have that the second part vanishes, whilst the first part achieves maximum violation, i. e.,  $2\sqrt{2}$ , then:

$$\begin{aligned} \max B_{\rho(p)} &= \left[ \frac{d(d-1)d^2}{2pd^2 + (1-p)4d(d-1)} \right] \\ &\quad \times \left[ \frac{p}{d(d-1)} 2(2\sqrt{2}) \right]. \end{aligned} \quad (35)$$

We have CHSH violation when  $\max B_{\rho_W^d(p)} > 2$ , Eq. (8). After some algebra we obtain that this holds for  $p \geq p_{NL}$ , with:

$$p_{NL} = \frac{4(d-1)}{2d(\sqrt{2}-1) + 4(d-1)}. \quad (36)$$

For instance, the values we are interested in:

- $d = 3, \quad p_{NL} = \frac{4}{17}(3\sqrt{2}-1) \approx 0.7630.$
- $d = 4, \quad p_{NL} = \frac{3}{7}(2\sqrt{2}-1) \approx 0.7837.$
- $d = 5, \quad p_{NL} = \frac{8}{41}(5\sqrt{2}-3) \approx 0.7944.$
- $d = 6, \quad p_{NL} = \frac{5}{14}(3\sqrt{2}-2) \approx 0.8009.$

Which are the values reported in Table 2 and in Liang et al. (2012). In Liang et al. (2012) they also reported that, after numerical optimisations over possible filters, filters given by Eq. (33) are optimal, in the sense that, the obtained two-qubit system violates CHSH at its best.

## Appendix B: Isotropic States

Here, we show the nonlocality-related properties limit points (except activation T&LF) for the two-qudit Isotropic states Eq. (29). They are entangled for  $p > p_E$  and local for  $p \leq p_L$ , with (Horodecki & Horodecki 1999):

$$p_E = \frac{1}{d+1}, \quad p_L = \frac{-1 + \sum_{k=1}^d \frac{1}{k}}{d-1}. \quad (37)$$

The explicit calculation of the fidelity of teleportation reported in Zhao et al. (2010), shows that all entangled Isotropic states are useful for teleportation, i. e.,  $f_d(\rho_{\text{ent}}) > \frac{1}{d}$ . Finally, nonlocality is checked through the violation of the CGLMP inequalities (Collins et al. 2002).

## References

Acín A, Gisin N, Masanes L. From Bell's Theorem to Secure Quantum Key Distribution, *Physical Review Letters*, 97(12):120405, 2006a  
doi:10.1103/PhysRevLett.97.120405.

Acín A, Gisin N, Toner B. Grothendieck's Constant and Local Models for Noisy Entangled Quantum States, *Physical Review A*, 73(6):062105, 2006b  
doi:10.1103/PhysRevA.73.062105.

Batle J, Casas M. Nonlocality and Entanglement in the XY Model, *Physical Review A*, 82(6):062101, 2010  
doi:10.1103/PhysRevA.82.062101.

Batle J, Casas M. Nonlocality and Entanglement in Gubit Systems, *Journal of Physics A: Mathematical and Theoretical*, 44(44):445304, 2011  
doi:10.1088/1751-8113/44/44/445304.

Bell JS. On the Einstein-Podolsky-Rosen Paradox, *Physics*, 1:195200, 1964

Bennett CH, Brassard G, Crépeau C, Jozsa R, Peres A, et al. Teleporting an Unknown Quantum State Via Dual Classical and Einstein-Podolsky-Rosen Channels, *Physical Review Letters*, 70(13): 18951899, 1993  
doi:10.1103/PhysRevLett.70.1895.

Bennett CH, Brassard G, Popescu S, Schumacher B, Smolin JA, et al. Purification of Noisy Entanglement and Faithful Teleportation via Noisy Channels, *Physical Review Letters*, 76(5):722725, 1996a  
doi:10.1103/PhysRevLett.76.722.

Bennett CH, Di Vincenzo DP, Smolin JA, Wootters WK. Mixed-state entanglement and quantum error correction. *Physical Review A*, 54(5):38243851, 1996b  
doi:10.1103/PhysRevA.54.3824.

Brunner N, Cavalcanti D, Pironio S, Scarani V, Wehner S. Bell Nonlocality, *Review of Modern Physics*, 86(2):419478, 2014  
doi:10.1103/RevModPhys.86.419.

Cavalcanti D, Acín A, Brunner N, Vértesi T. All Quantum States Useful for Teleportation are Nonlocal Resources, *Physical Review A*, 87(4):042104, 2013  
doi:10.1103/PhysRevA.87.042104.

Cirelson B. Quantum Generalizations of Bell's Inequality, *Letters in Mathematical Physics*, 4(2):93100, 1980  
doi:10.1007/BF00417500.

Clauser JF, Horne MA, Shimony A, Holt RA. Proposed Experiment to Test Local Hidden-Variable Theories, *Physical Review Letters*, 23(15):880884, 1969  
doi:10.1103/PhysRevLett.23.880.

Collins D, Gisin N. A Relevant Two Qubit Bell Inequality Inequivalent to the CHSH Inequality, *Journal of Physics A: Mathematical and General*, 37(5):1775, 2004  
doi:10.1088/0305-4470/37/5/021.

Collins D, Gisin N, Linden N, Massar S, Popescu S. Bell Inequalities for Arbitrarily High Dimensional Systems, *Physical Review Letters*, 88(4):040404, 2002  
doi:10.1103/PhysRevLett.88.040404.165.

Einstein A, Podolsky B, Rosen N. Can Quantum-Mechanical Description of Physical Reality Be Considered Complete? *Physical Review*, 47(10):777780, 1935  
doi:10.1103/PhysRev.47.777.

Ekert AK. Quantum Cryptography Based on Bell's Theorem, *Physical Review Letters*, 67(6):661663, 1991  
doi:10.1103/PhysRevLett.67.661.

Gisin N. Bell's Inequality Holds for All Non-Product States, *Physics Letters A*, 154(56):201202, 1991  
doi:10.1016/0375-9601(91)90805-I.

Gisin N. Hidden Quantum Nonlocality Revealed by Local Filters, *Physics Letters A* 210(3):151156, 1996  
doi:10.1016/S0375-9601(96)80001-6.

Gisin N, Peres A. Maximal Violation of Bell's Inequality for Arbitrarily Large Spin, *Physics Letters A*, 162(1):1517, 1992  
doi:10.1016/0375-9601(92)90949-M.

Grondalski J, Etlinger D, James D. The Fully Entangled Fraction as an Inclusive Measure of Entanglement Applications, *Physics Letters A*, 300(6):573580, 2002 doi:10.1016/S0375-9601(02)00884-8.

Hirsch F, Quintino MT, Bowles J, Brunner N. Genuine Hidden Quantum Nonlocality, *Physical Review Letters*, 111(16):160402, 2013 doi:10.1103/PhysRevLett.111.160402.

Horodecki M, Horodecki P. Reduction Criterion of Separability and Limits for a Class of Distillation Protocols, *Physical Review A*, 59(6):42064216, 1999 doi:10.1103/PhysRevA.59.4206.

Horodecki M, Horodecki P, Horodecki R. Separability of Mixed States: Necessary and Sufficient Conditions, *Physics Letters A*, 223(1-2):18, 1996a doi:10.1016/S0375-9601(96)00706-2.

Horodecki M, Horodecki P, Horodecki R. General Teleportation Channel, Singlet Fraction, and Quasidistillation, *Physical Review A*, 60(3):18881898, 1999 doi:10.1103/PhysRevA.60.1888.

Horodecki R, Horodecki M, Horodecki P. Teleportation, Bell's Inequalities and Inseparability, *Physics Letters A*, 222(12):2125, 1996b doi:10.1016/0375-9601(96)00639-1.

Horodecki R, Horodecki P, Horodecki M. Violating Bell Inequality by Mixed Spin-1/2 States: Necessary and Sufficient Condition, *Physics Letters A*, 200(5):340-344, 1995 doi:10.1016/0375-9601(95)00214-N.

Horodecki R, Horodecki P, Horodecki M, Horodecki K. Quantum Entanglement, *Review of Modern Physics*, 81(2):865942, 2009 doi:10.1103/RevModPhys.81.865.

Hua B, Li M, Zhang T, Zhou C, Li-Jost X, et al. Towards Grothendieck Constants and LHV Models in Quantum Mechanics, *Journal of Physics A: Mathematical and Theoretical*, 48(6):065302, 2015 doi:10.1088/1751-8113/48/6/065302.

Liang YC, Masanes L, Rosset D. All Entangled States Display Some Hidden Nonlocality, *Physical Review A*, 86:052115, 2012 doi:10.1103/PhysRevA.86.052115.

Löfberg J. YALMIP : A Toolbox for Modeling and Optimization in MATLAB. in: CCA/ISIC/CACSD, 2004 doi:10.1109/CACSD.2004.1393890.

Masanes L. Asymptotic Violation of Bell Inequalities and Distillability, *Physical Review Letters*, 97(5):050503, 2006  
doi:10.1103/PhysRevLett.97.050503.

Masanes L, Liang YC, Doherty AC. All Bipartite Entangled States Display Some Hidden Nonlocality, *Physical Review Letters*, 100(9):090403, 2008  
doi:10.1103/PhysRevLett.100.090403.

MATLAB version 7.13.0.564 (R2011b). The MathWorks Inc., Natick, Massachusetts, 2011.

Miranowicz A. Violation of Bell Inequality and Entanglement of Decaying Werner States, *Physics Letters A*, 327(4):272283, 2004  
doi:10.1016/j.physleta.2004.05.001

Navascués M, Vértesi T. Activation of Nonlocal Quantum Resources, *Physical Review Letters*, 106(6):060403, 2011  
doi:10.1103/PhysRevLett.106.060403.

Nielsen MA, Chuang IL. Quantum Computation and Quantum Information. Cambridge University Press, Cambridge, UK. 2000

Pal R, Ghosh S. Non-locality Breaking Qubit Channels: The Case for CHSH Inequality, *Journal of Physics A: Mathematical and Theoretical*, 48(15):155302, 2015  
doi:10.1088/1751-8113/48/15/155302.

Palazuelos C. Superactivation of Quantum Nonlocality, *Physic Review Letters*, 109(19):190401, 2012  
doi:10.1103/PhysRevLett.109.190401.

Peres A. Collective Tests for Quantum Nonlocality, *Physical Review A*, 54(4):2685-2689, 1996a  
doi:10.1103/PhysRevA.54.2685.

Peres A. Separability Criterion for Density Matrices, *Physical Review Letters*, 77(8):14131415, 1996b  
doi:10.1103/PhysRevLett.77.1413.

Popescu S. Bell's Inequalities Versus Teleportation: What is Nonlocality?, *Physical Review Letters*, 72:797799, 1994  
doi:10.1103/PhysRevLett.72.797.

Popescu S. Bell's Inequalities and Density Matrices: Revealing Hiden Nonlocality, *Physical Review Letters*, 74(14):26192622, 1995  
doi:10.1103/PhysRevLett.74.2619.

Popescu S, Rohrlich D. Generic Quantum Nonlocality, *Physics Letters A*, 166(5-6):293297, 1992  
doi:10.1016/0375-9601(92)90711-T.

Sturm J. Using SeDuMi 1.02, a MATLAB toolbox for optimization over symmetric cones, *Optimization Methods and Software*, 1112:625653, 1999  
doi:10.1080/10556789908805766.

Terhal BM, Vollbrecht KGH. Entanglement of Formation for Isotropic States, *Physical Review Letters*, 85(12):26252628, 2000  
doi:10.1103/PhysRevLett.85.2625.

Toh KC, Todd M, Tutuncu RH. SDPT3 a MATLAB software package for semidefinite programming, *Optimization Methods and Software*, 11(1-4):545581, 1999  
doi:10.1080/10556789908805762.

Vandenberghe L, Boyd S. Semidefinite Programming, *SIAM Review*, 38(1):4995, 1996.

Verstraete F, Wolf MM. Entanglement Versus Bell Violations and Their Behavior Under Local Filtering Operations, *Physical Review Letters*, 89(17):170401, 2002  
doi:10.1103/PhysRevLett.89.170401.

Vollbrecht KGH, Werner RF. Entanglement Measures Under Symmetry, *Physical Review A*, 64(6):062307, 2001  
doi:10.1103/PhysRevA.64.062307.

Werner RF. Quantum States With Einstein-Podolsky-Rosen Correlations Admitting a Hidden- Variable Model, *Physical Review A*, 40(8):42774281, 1989  
doi:10.1103/PhysRevA.40.4277.

Wootters WK. Entanglement of Formation of an Arbitrary State of Two Qubits, *Physical Review Letters*, 80(10):22452248, 1998  
doi:10.1103/PhysRevLett.80.2245.

Wootters WK. Entanglement of Formation and Concurrence, *Quantum Information and Computation*, 1(1):2744, 2001.

Zhao MJ, Li ZG, Fei SM, Wang ZX. A Note on Fully Entangled Fraction, *Journal of Physics A: Mathematical and Theoretical*, 43(27):275203, 2010  
doi:10.1088/1751-8113/43/27/275203.

### Sobre la activación de no-localidad cuántica

**Resumen.** Se reportan algunas propiedades cuánticas de sistemas físicos, a saber, entrelazamiento, no-localidad, no-localidad a k-copias (superactivación de la no-localidad), no-localidad oculta (activación de la no-localidad a través de filtrado local) y activación de la no-localidad a través de producto tensorial y filtrado local. Este trabajo tiene dos propósitos: en primer lugar, proporcionar una reseña de los procedimientos numéricos que deben seguirse con el fin de calcular las propiedades mencionadas, en particular para cualquier estado de dos qubits, así como reproducir las cotas para los estados Werner de dos qudits. En segundo lugar, se utilizan estas herramientas numéricas para calcular nuevas cotas de estas propiedades para los estados isotrópicos de dos qudits y los estados de Hirsch de dos qubits.

**Palabras clave:** qubits; información cuántica, nolocalidad cuántica, entrelazamiento.

### Sobre a Ativação da Não-localidade Quântica

**Resumo.** Apresentamos um estudo sobre as propriedades quânticas de sistemas físicos, especialmente, o emaranhamento, a não-localidade, a não-localidade k-copy (superativação da não-localidade) e a ativação de não-localidade através do uso de tensores e filtragem local. O objetivo deste trabalho foi duplo. Primeiro, fornecemos uma revisão do procedimento numérico a ser seguido a fim de calcular as propriedades acima mencionadas, em particular, para qualquer sistema de dois-qubits, e que reproduza a ligação para dois-qudit em Estado Werner. Em segundo lugar, utilizamos estas ferramentas numéricas para calcular novas ligações destas propriedades para estados isotrópicos e Hirsch de dois-qudit.

**Palavras-chave:** qudits; informação quântica; não-localidade quântica; emaranhamento.

#### Andrés Felipe Ducuara

Is a MSc-Physics student at the Universidad del Valle, Cali-Colombia. His current work focuses on quantum information theory and quantum foundations.

#### Javier Madroñero

Is Professor of the Physics Department at the Universidad del Valle, Cali-Colombia. His current research activities focus on quantum optics, quantum information theory and quantum control.

#### John Henry Reina

Is a Professor of Physics at Universidad del Valle, Cali, Colombia. He is the Head of the Quantum Technologies, Information and Complexity Group, and the Founder Director of the recently created Centre for Bioinformatics and Photonics. He is an Oxonian Theoretical Physicist (D.Phil., 2002). His main research interests focus on the Physics of Information, Quantum Optics, Macroscopicity, Decoherence, and Experimental Quantum Information and Ultrafast Molecular Dynamics.